Title	Production of equol from daidzein by gram-positive rod-shaped bacterium isolated from rat intestine.
Author(s)	Minamida, Kimiko; Tanaka, Michiko; Abe, Ayumi; Sone, Teruo; Tomita, Fusao; Hara, Hiroshi; Asano, Kozo
Citation	Journal of Bioscience and Bioengineering, 102(3), 247-250 https://doi.org/10.1263/jbb.102.247
Issue Date	2006-09
Doc URL	http://hdl.handle.net/2115/14823
Type	article (author version)
File Information	JB&B102-3.pdf

[Instructions for use](#)

NOTES

Production of Equol from Daidzein by Gram-Positive Rod-Shaped Bacterium Isolated
from Rat Intestine

Kimiko Minamida,^{1*} Michiko Tanaka,¹ Ayumi Abe,¹ Teruo Sone,¹ Fusao Tomita,¹
Hiroshi Hara,¹ and Kozo Asano¹

*Division of Applied Bioscience, Graduate School of Agriculture, Hokkaido University,
Kita 9, Nishi 9, Kita-ku, Sapporo, Hokkaido 060-8589, Japan¹*

Received 11 May 2006 / Accepted 27 June 2006

[Key words: isoflavone, equol, equol-producing bacterium]

*Corresponding author. e-mail: kimiko-m@chem.agr.hokudai.ac.jp

phone: +81-(0)11-706-2493 fax: +81-(0)11-706-4961

1 Isoflavones (mainly daidzein and genistin) belong to the flavonoid group of
2 compounds and are classified as phytoestrogens. In the intestine, daidzin is converted to
3 daidzein by β -glucosidase, and then daidzein is converted to *O*-desmethylangolensin
4 (*O*-DMA) or equol via dihydrodaidzein by enzymes of intestinal bacteria. We isolated,
5 for the first time, an anaerobic gram-positive rod-shaped strain capable of producing
6 equol from daidzein. Its 16S rDNA gene sequence (1428 bp) showed 99% similarity
7 with that of the human intestinal bacterium SNU-Julong 732 (AY310748) and 93%
8 similarity with that of *Eggerthella lenta* ATCC 25559^T (AF292375). This strain
9 converted daidzein to equol via dihydrodaidzein in an equol-assay medium
10 anaerobically. The addition of butyric acid and arginine increased the conversion ratio
11 of daidzein to equol 4.7- and 4.5-fold, respectively.

1 Isoflavones are flavonoids present in various plants, particularly in soybean germ.
2 They are classified as phytoestrogens because their structures resemble that of estrogen
3 and they have a weak affinity for the estrogen receptor. Epidemiologic and experimental
4 studies showed that they had preventive effects on breast cancer, prostate cancer,
5 cardiovascular disease, osteoporosis and menopausal symptoms (1, 2). Isoflavones exist
6 as glycosides in some plants, mainly as daidzin and genistin. In the intestine, daidzin is
7 converted to daidzein by β -glucosidase. Then, daidzein is converted via dihydrodaidzein
8 to *O*-desmethylangolensin (*O*-DMA) or equol by enzymes of intestinal bacteria (3; Fig.
9 1). Equol has a stronger estrogenic activity than daidzein and *O*-DMA (4, 5). Humans
10 capable of producing equol from daidzein (equol producers) have a lower risk of
11 developing breast and prostate cancers than non-equol producers (6, 7). Several animals,
12 particularly rodents, produce adequate concentrations of equol. However, in humans,
13 only 30-50% of the population can produce equol owing to differences in intestinal
14 microbiota among individuals (8, 9). Daidzein-metabolizing phenotypes are stable in
15 individuals over time (7) because the intestinal microbiota of such individuals are stable.
16 Therefore, non-equol producers excrete no equol, even when they ingest soy protein
17 powder (34 g/d) for one month (10).

18 There are few reports on daidzein-metabolizing intestinal bacteria. An anaerobic

1 gram-positive strain HGH 6 that converts daidzein to dihydrodaidzein (11), and a
2 *Clostridium* sp. strain HGH 136 (12) and *Eubacterium ramulus* (13) that converts
3 daidzein to *O*-DMA were isolated from humans. The human intestinal bacterium
4 SNU-Julong 732 that converts dihydrodaidzein to equol was also isolated from humans
5 (14). A mixture of *Bacteroides ovatus*, *Ruminococcus productus* and *Streptococcus*
6 *intermedius* (15) or *Lactobacillus mucosae*, *Enterococcus faecium*, *Finnegoldia magna*
7 and *Veillonella* sp. (16) produces equol from daidzein. However, no bacterium that
8 produces equol from daidzein alone (an equol-producing bacterium) has yet been found.
9 Therefore, we intended to isolate an equol-producing bacterium from rat cecal contents,
10 because rats are good producers of equol.

11 Standards for daidzein and equol were purchased from LC Laboratories (Woburn,
12 MA, USA). Dihydrodaidzein was purchased from Toronto Research Chemicals. (North
13 York, ON, Canada). The equol-assay medium contained 29.5 g of GAM broth (Nissui
14 Pharmaceutical, Tokyo), 10 g of CaCO₃ and 2 g of Fujiflavone P10 (Fujicco, Kobe) per
15 liter of distilled water. After autoclaving, the medium was stored in an anaerobic
16 chamber (Coy Laboratory Products, Grass Lake, MI, USA) under an 85% N₂, 10% CO₂
17 and 5% H₂ atmosphere. A frozen glycerol stock of the cecal content of a male
18 Sprague-Dawley rat (SLC Japan, Tokyo) fed a casein diet for 3 weeks according to the

AIN-93G formulation (17) was added to the equol-assay medium and incubated anaerobically at 37°C for 2 d. The culture broth was spread on a plate containing 14.75 g of GAM broth, 2 g of Fujiflavone P10 and 15 g of agar per liter of distilled water, and then incubated anaerobically at 37°C for 2 d. A number of colonies that developed on the plate were selected and inoculated into the equol-assay medium, incubated anaerobically at 37°C for 2 d, extracted and analyzed by HPLC as described below.

After isolating an equol-producing bacterium, a precultured GAM broth containing 1% L-arginine at 37°C for 28 h was added to an equol-assay medium for quantitative determination containing 59 g of GAM broth and daidzein (final concentration: 200 µM) per liter of distilled water. Then, the medium was incubated anaerobically at 37°C, extracted and analyzed by HPLC as described below. To investigate the effects of arginine and butyric acid on equol production, 1% arginine and/or butyric acid (final concentration: 50 mM) was added to the equol-assay medium (arginine, before autoclaving; butyric acid, after autoclaving) and the resulting solutions were analyzed similarly. The pH of the medium containing 1% arginine was adjusted to 7.0 before autoclaving, whereas the pH of the arginine-free medium was not adjusted (approximately pH 7.0). Absorbance at 600 nm (OD₆₀₀) was measured using Spectronic 20D+ spectrophotometer (Thermo Electron, Waltham, MA, USA) and culture broth pH

1 was measured using an ISFET pH meter KS-701 (Shindengen Electric, Tokyo).

2 Aliquots of the assay media were extracted three times with ethyl acetate of 1.5
3 volume of the media and evaporated using a rotary evaporator. Then, the aliquots were
4 dissolved in methanol and filtered using a 0.45- μ m filter (Millex-LH; Millipore, Tokyo).
5 Each sample was injected into HPLC (Jasco, Tokyo) equipped with a Mightysil RP-18
6 GP 250-3.0 column (3.0 $\times$ 250 mm; 3 μ m; Kanto Chemical; Tokyo) and a UV detector
7 (280 nm; Jasco). The mobile phase was a solution of water : acetonitrile : acetic acid,
8 75 : 25 : 0.1 (V/V/V), the flow rate was 0.4 ml/min and the column temperature was
9 60°C. Metabolites were identified by comparing their retention times with those of
10 standards.

11 Cell morphology after anaerobic cultivation for 2 d at 37 °C in GAM broth was
12 examined using phase-contrast microscopy (ECLIPSE E600; Nikon, Tokyo). The Gram
13 staining solution used was neo-B&M Wako (Wako, Osaka). The isolated bacterium
14 was identified by 16S rDNA gene sequence analysis (18). Homology searches were
15 performed in the GenBank database using the BLAST search program. Some 16S rDNA
16 sequences were retrieved from the DDBJ, EMBL and GenBank databases for
17 comparison in the phylogenetic analysis. Sequence data were aligned with the
18 CLUSTAL X package program and corrected by manual inspection. Nucleotide

substitution rates were calculated, and a phylogenetic tree was constructed using the neighbor-joining method.

An anaerobic gram-positive rod-shaped strain capable of producing equol was isolated from a rat cecal content. This strain is referred to as the Gram-positive bacterium do03 (AB266102). Its 16S rDNA gene sequence (1428 bp) showed 99% similarity with the human intestinal bacterium SNU-Julong 732 (AY310748), 94% similarity with *Eggerthella sinensis* HKU14 (AY321958), 94% similarity with *Eggerthella hongkongensis* HKU10 (AY288517) and 93% similarity with *Eggerthella lenta* ATCC 25559^T (AF292375). The phylogenetic tree showed that the isolated strain does not belong to the genus *Eggerthella* (Fig. 2). The strains do03 and Julong 732 occupy the same cluster. Therefore, these strains may belong to a new genus. Moreover, the strain Julong 732 was isolated from a fecal sample of a healthy female human and the strain do03 was isolated from a rat cecal content. Hence, these strains are indigenous intestinal bacteria.

The strain do03 converted 200 μ M daidzein to equol via dihydrodaidzein for 4 d at 37°C anaerobically (Fig. 3). For 2 and 4 d, distinct peaks were observed at 4 min. There have been no reports on equol being metabolized by intestinal bacteria. Despite the fact that different amounts of equol were produced, the areas of the peaks were about the

same (data not shown). Therefore, these peaks do not correspond to equol split products but to other metabolic products.

In GAM broth (control), the conversion ratio of daidzein to equol (equol ratio: amount of equol production/amount of supplemented daidzein) was 0.15 ± 0.01 . In the medium containing butyric acid, the equol ratio increased to 0.71 ± 0.03 , although OD₆₀₀ and culture broth pH did not change compared with those of the control. In the medium containing arginine, the equol ratio increased to 0.67 ± 0.01 with increases in OD₆₀₀ and culture broth pH. In the medium containing butyric acid and arginine, the equol ratio increased to 0.58 ± 0.01 with a slight increase in culture broth pH (Table 1). Because butyric acid stimulates equol production (16), the equol ratio in the medium containing butyric acid was considered to have increased. Moreover, for the growth of some bacteria such as *E. lentum*, arginine is required because they obtain energy for growth using the arginine dihydrolase pathway (19). The bacterial metabolism of arginine produces NH₃, which caused the increase in culture broth pH. Arginine supplementation increased OD₆₀₀; thus, the strain do03 uses arginine for growth. Therefore, the increase in equol ratio may be attributed to an increase in the number of do03 cells. The supplementation of butyric acid and arginine decreased the equol ratio by approximately 10%. Because culture broth pH increased more than that of the

control, the strain do03 seemed to use arginine; however, OD₆₀₀ did not increase. Butyric acid supplementation caused a decrease in OD₆₀₀. The mechanism of equol production stimulated by butyric acid supplementation has not yet been reported. Antagonist action seemed to occur by the supplementation of butyric acid and arginine.

In the human intestine, when the strains HGH 6 and Julong 732 are present, the microbial community is able to produce equol from daidzein (16). However, the strain do03 converted daidzein to equol via dihydrodaidzein without any other strains (Fig. 3).

The authors thank Fujicco Co., Ltd. for providing us with Fujiflavone P10 and Mr. Ongol Martin Patrick for assistance with the preparation of this manuscript.

REFERENCES

1. **Adlercreutz, H.:** Phyto-oestrogens and cancer. *Lancet Oncol.*, **3**, 364-373 (2002).
2. **Duncan, A. M., Phipps, W. R., and Kurzer, M. S.:** Phyto-oestrogens. *Best Pract. Res. Clin. Endocrinol. Metab.*, **17**, 253–271 (2003).
3. **Hirayama, K:** Metabolism of soy isoflavones by intestinal flora. *J. Intestinal Microbiol.*, **19**, 17-23 (2005). (in Japanese)
4. **Schmitt, E., Dekant, W., and Stopper, H.:** Assaying the estrogenicity of phytoestrogens in cells of different estrogen sensitive tissues. *Toxicol. In Vitro*, **15**,

433-439 (2001).

5. **Sathyamoorthy, N. and Wang, T. T.:** Differential effects of dietary phyto-oestrogens daidzein and equol on human breast cancer MCF-7 cells. *Eur. J. Cancer*, **33**, 2384-2389 (1997).

6. **Duncan, A. M., Merz-Demlow, B. E., Xu, X., Phipps, W. R., and Kurzer, M. S.:** Premenopausal equol excretors show plasma hormone profiles associated with lowered risk of breast cancer. *Cancer Epidemiol. Biomark. Prev.*, **9**, 581-586 (2000).

7. **Akaza, H., Miyanaga, N., Takashima, N., Naito, S., Hirano, Y., Tsukamoto, T., Fujioka, T., Mori, M., Kim, W. J., Song, J. M., and Pantuck, A. J.:** Comparisons of percent equol producers between prostate cancer patients and controls: case-controlled studies of isoflavones in Japanese, Korean and American residents. *Jpn. J. Clin. Oncol.*, **34**, 86-89 (2004).

8. **Setchell, K. D., Brown, N. M., Desai, P. B., Zimmer-Nechimias, L., Wolfe, B., Jakate, A. S., Creutzinger, V., and Heubi, J. E.:** Bioavailability, disposition, and dose-response effects of soy isoflavones when consumed by healthy women at physiologically typical dietary intakes. *J. Nutr.*, **133**, 1027-1035 (2003).

9. **Arai, Y., Uehara, M., Sato, Y., Kimira, M., Eboshida, A., Adlercreutz, H., and Watanabe, S.:** Comparison of isoflavones among dietary intake, plasma concentration

and urinary excretion for accurate estimation of phytoestrogen intake. J. Epidemiol., **10**, 127–135 (2000).

10. **Lampe, J. W., Skor, H. E., Li, S., Wahala, K., Howald, W., N., and Chen, C.:**

Wheat bran and soy protein feeding do not alter urinary excretion of the isoflavan equol in premenopausal women. J. Nutr., **131**, 740-744 (2001).

11. **Hur, H. G., Lay, J. O., Jr, Beger, R. D., Freeman, J. P., and Rafii, F.:** Isolation of human intestinal bacteria metabolizing the natural isoflavone glycosides daidzin and genistin. Arch. Microbiol., **174**, 422–428 (2000).

12. **Hur, H. G., Beger, R. D., Heinze, T. M., Lay, J. O., Jr., Freeman, J. P., Dore, J., and Rafii, F.:** Isolation of an anaerobic intestinal bacterium capable of cleaving the C-ring of the isoflavonoid daidzein. Arch. Microbiol., **178**, 8-12 (2002).

13. **Schoefer, L., Mohan, R., Braune, A., Birringer, M., and Blaut, M.:** Anaerobic C-ring cleavage of genistein and daidzein by *Eubacterium ramulus*. FEMS Microbiol. Lett., **208**, 197-202 (2002).

14. **Wang, X. L., Hur, H. G., Lee, J. H., Kim, K. T., and Kim, S. I.:** Enantioselective synthesis of *S*-equol from dihydrodaidzein by a newly isolated anaerobic human intestinal bacterium. Appl. Environ. Microbiol., **71**, 214-219 (2005).

15. **Ueno, T., Uchiyama, S., and Kikuchi, N.:** The role of intestinal bacteria on

- biological effects of soy isoflavones in humans. J. Nutr., **132**, 594S (2002).
16. **Decroos, K., Vanhemmens, S., Cattoir, S., Boon, N., and Verstraete, W.:**
Isolation and characterisation of an equol-producing mixed microbial culture from a
human faecal sample and its activity under gastrointestinal conditions. Arch. Microbiol.,
183, 45-55 (2005).
17. **Reeves, P. G., Nielsen, F. H., and Fahey, G. C., Jr.:** AIN-93 purified diets for
laboratory rodents: final report of the American Institute of Nutrition ad hoc writing
committee on the reformulation of the AIN-76A rodent diet. J. Nutr., **123**, 1939-1951
(1993).
18. **Minamida, K., Shiga, K., Sujaya, I. N., Sone, T., Yokota, A., Hara, H., Asano, K.,
and Tomita, F.:** Effects of difructose anhydride III (DFA III) administration on rat
intestinal microbiota. J. Biosci. Bioeng., **99**, 230-236 (2005).
19. **Sperry, J. F. and Wilkins, T. D.:** Arginine, a growth-limiting factor for
Eubacterium lentum. J. Bacteriol., **127**, 780-784 (1976).

1 FIG. 1. Metabolism of daidzein to equol in intestine.

2 FIG. 2. 16S rDNA-based phylogenetic tree showing relationship between newly isolated
3 strain do03 and other closely related species. The numbers are bootstrap values
4 calculated from 1000 trees. GenBank accession numbers are shown in
5 parentheses.

6 FIG. 3. HPLC elution profiles of the supernatant of daidzein conversion by newly
7 isolated strain do03 at 0, 2 and 4 d under anaerobic condition in equol-assay
8 medium containing 50 mM butyric acid. The concentration of daidzein was
9 192.7 μ M at 0 d; those of daidzein, dihydrodaidzein and equol were 62.3, 45.3
10 and 63.7 μ M, respectively, at 2 d; and that of equol was 138.7 μ M at 4 d.

Minamida *et al.* FIG. 3

1

2

TABLE 1. Effects of arginine and butyric acid on equol production

	GAM (control)	GAM + butyric acid	GAM+ arginine	GAM + butyric acid and arginine
Conversion ratio of daidzein to equol	0.15 ± 0.01	0.71 ± 0.03	0.67 ± 0.01	0.58 ± 0.01
OD ₆₀₀	0.277 ± 0.005	0.222 ± 0.003	0.431 ± 0.004	0.203 ± 0.004
pH	7.2	7.2	7.9	7.5

3

Values are means $\pm$ SD.

4

5