



Title	On the Correlation between the Satellite of Chromosome and the Leaf Shape in Parideae from Hokkaido
Author(s)	STOW, Isamu
Citation	Journal of the Faculty of Science, Hokkaido Imperial University. Ser. 5, Botany, 4(1), 31-46
Issue Date	1935
Doc URL	http://hdl.handle.net/2115/26246
Type	bulletin (article)
File Information	4(1)_P31-46.pdf



Instructions for use

On the Correlation between the Satellite of Chromosome and the Leaf Shape in *Parideae* from Hokkaido

By

ISAMU STOW

The correct basic number of chromosomes of *Parideae* was determined as five by GOTOH and STOW (1930) who found that these five chromosomes of each species are different in size and shape from each other. Moreover, four of these five types of chromosomes are like not only among the different species of the same genus, but also in the two different genera, *Paris* and *Trillium*. The fifth chromosome type of *Paris*, however, differs in each species respectively, while in *Trillium* it is always almost similar in shape. GOTOH (1933) studied further this relation with an American species of *Trillium* and found that its basic number of chromosomes is also five and that the types of these five chromosomes are similar to those of the Japanese *Trillium*. HAGA (1934) made a very close investigation with Japanese *Parideae* and confirmed the results of GOTOH and STOW (1930).

Satellites were first reported by S. NAWASCHIN (1912) in *Galtonia candicans*, and afterwards many other plants¹⁾ were found, which carry satellites of various types.²⁾ The asymmetry³⁾ of nuclei in connection with the occurrence of satellites was also ascertained by the same author (1912) in *Galtonia* and in *Muscati* by DELAUNAY (1915). Such asymmetry or elimination of satellites is often accompanied by deviation of some physiological or morphological characters as seen in *Galtonia* (S. NAWASCHIN 1915), *Muscati* (S. NAWASCHIN 1915, DELAUNAY 1915) or *Crepis* (M. NAVASHIN 1925, 1926) etc.

Although there have been many cytological investigations on *Parideae*⁴⁾, the relation between the chromosome types and the morphological characters of these plants have been studied very

1) See KUHN (1928).

2) See DELAUNAY (1929).

3) The term "asymmetry and symmetry" of nuclei is used in the same sense as by S. NAWASCHIN.

4) See HAGA (1934), HUSKINS and SMITH (1935), SMITH (1935) and MATSUURA (1935).

rarely. In a joint work with GOTOH (1930), the present writer reported that the diploid plant of *Paris* of the asymmetrical type with regard to the satellite grows more vigorously than the symmetrical ones, as confirmed in *Galtonia* (S. NAWASCHIN 1912) or *Muscari* (S. NAWASCHIN 1915, DELAUNAY 1915) and that the ratios of the breadth to the length of the laminae of the asymmetrical type are greater than those of the symmetrical ones. By a re-examination of the results of the former investigation and by further studies, the writer has found a noticeable relation between the satellites and an outer morphological character of the leaf.

Material and method

The materials used in the present investigation are as follows: *Paris hexaphylla* CHAM., *Paris tetraphylla* A. GRAY, *Trillium kamtschaticum* PALL., *Trillium Tschonoskii* MAXIM. and *Trillium Smallii* MAXIM.¹⁾. The majority of these materials were collected in the vicinity of Sapporo, and some of them in the Botanical Garden of the university, where they grew under the wild condition.

FLEMMING-Bonn and NAVASHIN's²⁾ fluids were employed for the fixation of the root-tip, both normal and chloralized materials, after SAKAMURA's procedure (1920). The microtome sections were stained with HEIDENHAIN's iron-alum hematoxylin or with MACCLINGSTOCK's iodine crystal violet; in both cases the results were satisfactory. For the fixation of the anther, in which the reduction division in the pollen-mother cells occurred, FLEMMING-Bonn solution was chiefly employed and the sections were stained with HEIDENHAIN's iron-alum hematoxylin with fairly good results. The most satisfactory result, however, was obtained with the iron-acetocarmine method.

The size of the angle between the two edges of the leaf margin at the base was calculated in a simple geometrical way. No doubt the leaf margin does not show a straight, but a curved line. In order to simplify the calculation, the writer treated it as a straight line, for its curvature is not remarkably great. Now let us consider a triangle $\triangle OBC$ (Fig. 1) whose base BC is the maximum width of laminae and the apex O is the leaf base. It is easy to measure the size of the angle between the two edges of the leaf margin at the base, which will be

1) In the previous paper (GOTOH & STOW 1930) the synonymous name *T. apetalon* MAKINO was used.

2) The formalin is reduced to 1/4 of the original receipt.

indicated hereafter by θ for simplifying the description, assuming it to be approximately equal to the angle $\angle BOC$. Measuring the length BC, the maximum width of the laminae, and the height OD, the vertical line from O to BC, one can obtain the value of θ from the following formula:

$$\tan \frac{\theta}{2} = \frac{BC}{2OD}$$

In some materials the mean values of both BC and OD from many leaves were used for the calculation of θ , while in the others the calculation was made with each leaf separately and then the mean value of θ was obtained¹⁾.

The θ value thus obtained is somewhat smaller than the value of the angle between two tangent lines (OB' and OC') and this error becomes larger, the wider the laminae. Nevertheless there is no marked objection to assuming $\angle BOC$ as equal to θ , because this error does not extend beyond the limit of the purpose of the present investigation.




Fig. 1.

Results

The chromosome set of *Parideae* consists of five chromosomes having different size and shape (Fig. 2)²⁾.

a-chromosome: The largest V-shaped chromosome with a median attachment (52% of the total length from one end).

b-chromosome: A V-shaped chromosome about 2/3 as long as the a-chromosome with a submedian attachment (58%–62% of the total length from one end).

c-chromosome: A J-shaped chromosome with an atelomitic attachment (70%–74% of the total length from one end), somewhat shorter than the b-chromosome.

1) The size of the leaves was measured with the materials in bloom except *T. Smallii*, which was measured after bloom. The angle θ becomes somewhat smaller, when the material bears the berry.

2) See also GOTOH and STOW (1930), GOTOH (1933), HAGA (1934 a and b), MATSUURA (1934), HUSKINS and SMITH (1935).




Fig. 2. Chromosomal elements of *Paris* and *Trillium*; chloralized root-tip, 300×. Ph. *Paris hexaphylla* Pt, *Paris tetraphylla*. Tk, *Trillium kamtschaticum*. TT, *Trillium Tschonoskii*. TS, *Trillium Smallii*. *a, b, c, d* and *e*, chromosome type.

d-chromosome: The shortest V-shaped chromosome, about half as long as the a-chromosome, with a submedian attachment (60%–64% from one end).

e-chromosome: A J-shaped chromosome with a subterminal attachment, of median length between the c- and d-chromosome. In the case of *Paris*, it carries a satellite on the proximal knob, while in *Trillium* it has no satellite.

There are slight differences between the chromosomes of the same type among the different species except in the e-chromosome, so the chromosomes can be distinguished by precise measurement and observation. In the case of the e-chromosome, especially in *Paris*, however, there are characteristic differences between one species and another. The e-chromosome of *Paris hexaphylla* has a somewhat large oblong satellite on the proximal knob with a short connecting filament, and that of *P. tetraphylla* has a relatively large stick shaped satellite, which is thicker than a mere connecting filament. In the first nuclear division of the pollen grain, one can find a spiral with long pitch, deeply stained with iron-acetocarmine, and embedded in the weak stained ground matrix substance¹⁾ (Figs. 11a and b). Therefore, the stick shaped portion may be considered as an elongated satellite²⁾. The satellite of the e-chromosome of *P. japonica* is of a characteristic tandem shape (HAGA 1934 a). In the case of *Trillium*, however, the e-chromosome has only a knob, but no satellite, and only slight differences in the size of the knob or the limb can be recognized between the different species. The e-chromosome of




Fig. 3. *Trillium kamtschaticum*, first nuclear division in the pollen grain, anaphase, e-chromosomes indicated by solid black. 1000×

1) At the kinetic constriction of the chromosome of *Paris* this deeply stained spiral structure disappears, and it seems that it is broken at this portion. The connecting filament of the satellite is constructed with a deeply staining thin thread, and the spiral seems to be stretched to a thread without any matrix.

2) HAGA (1934 a) has studied the somatic chromosomes especially in the root-tip fixed with modified FLEMMING's solution and stained with iodine-gentian violet after NEWTON. He stated that the e-chromosome of *P. tetraphylla* (dt after HAGA) has a small satellite on the proximal knob with a long connecting filament, and that on some occasions its form changes into a long thread, especially at anaphase.

T. kamtchaticum has a knob of almost the same size as *Paris*, and the part of its limb nearer to the knob is somewhat more slender than the other part. This character does not appear in other species of *Trillium* and *Paris*, and it is clearly visible in the first nuclear division of the pollen grain especially at anaphase (Figs. 3 and 15). The knob of the e-chromosome of *T. Smallii* is somewhat larger than in the other species.

In some individuals of *Paris*, the satellite of the e-chromosome is reduced; in the materials studied, one of two e-chromosomes of the diploid plant of *P. hexaphylla* and *P. tetraphylla*, and one or two of the e-chromosomes of the triploid plant of *P. hexaphylla* do not carry a satellite. Now we can suppose the existence of several caryotypes in *Paris* and *Trillium* from Hokkaido regarding the elimination of the satellite of the e-chromosome as follows:

The chromosome value of a nucleus is indicated by Roman numerals II or III..., and the number of satellites of the e-chromosome in a nucleus is indicated by arabic numerals 0, 1, 2,....

<i>Paris hexaphylla</i> CHAM.	
"	II ₂
"	II ₁ *
"	II ₀
"	III ₃
"	III ₂
"	III ₁ *
"	III ₀
<i>Paris tetraphylla</i> A. GRAY	II ₂
"	II ₁ *
"	II ₀
<i>Trillium kamtchaticum</i> PALL.	II ₀
<i>T. Tschonoskii</i> MAXIM.	IV ₀ ¹⁾
<i>T. Smallii</i> MAXIM.	VI ₀

Among these caryotypes, *P. hexaphylla* II₀, III₀ and *P. tetraphylla* II₀, those marked with an *, have not been found, but are expected to exist. The plant of each caryotype, which was found actually, has a different morphological appearance, especially in the leaf character (Figs. 6-10). The writer reported in a joint work with GOTOH (1930) that the ratios of breadth to length of the laminae of *P. hexaphylla* II₂, II₁ and III₁²⁾ are different from each other.

1) *T. Tschonoskii* f. *violaceum* MAKINO has two kinds of the c- and e-chromosomes (Fig. 4), and it seems to be a hybrid plant between *T. Tschonoskii* and *T. Smallii*.

2) Described as types 1, 2 and 3 respectively in the paper of GOTOH and STOW (1930).




Fig. 4. *Trillium Tschonoskii* f. *violaceum*; root-tip, showing two kinds of c-and e-chromosomes, c-chromosome dotted and e-chromosome solid black. 1000 \times

Similar studies with other caryotypes, *P. hexaphylla* III₃, III₂ and *P. tetraphylla* II₂ and II₁, show the results indicated in Table 1. As the differences between the above mentioned ratios in respect to the leaf character in each caryotype are larger than 5 times their probable errors, it is not probable that such differences are a matter of chance.

Table I.
Breadth, length and the ratio of breadth to length of laminae
of the different caryotypes of *Paris*.

	breadth (mm)	length (mm)	ratio
<i>Paris hexaphylla</i> II ₂	16.975±0.168	63.422±0.819	0.2781±0.0025
„ II ₁	24.315±0.343	66.489±0.639	0.3651±0.0016
„ III ₃	31.061±0.394	78.073±1.000	0.3991±0.0038
„ III ₂	31.780±0.332	74.870±0.705	0.4271±0.0035
„ III ₁	35.076±0.302	77.557±0.430	0.4521±0.0033
<i>Paris tetraphylla</i> II ₂	38.645±0.479	68.458±0.842	0.5650±0.0042
„ II ₁	32.875±0.816	67.209±1.294	0.4950±0.0056

The most probable reason for such differences of the ratios is the elimination of the satellite of the e-chromosome. By a further investigation, the writer ascertained that the elimination of the satellite is more closely related to the angle between the two edges of the leaf margin at the base (θ) than it is to the ratio of the breadth to length of the laminae. The angle θ in *Paris* was measured and the results obtained are shown in Table 2.

Table II.

Angle between the two edges of leaf margin at the base (θ), quotient
 $360^\circ/\text{leaf number } (A)$ and degree of decrease of θ (λ) of
the different caryotypes of *Paris*.

	ratio of breadth to length of lamina (%)	θ	$A = \frac{360^\circ}{n}$	$\lambda = \frac{A-\theta}{A}$
<i>P. hexaphylla</i> II ₂	27.81 ± 0.245	28°20'	$\frac{360^\circ}{n} = 45^\circ$	0.372
	" II ₁	36.51 ± 0.159	36°21'	0.189
	" III ₃	39.91 ± 0.375	38°18'	0.149
	" III ₂	42.71 ± 0.345	41°12'	0.088
	" III ₁	45.21 ± 0.331	43°24'	0.085
<i>P. tetraphylla</i> II ₂	56.50 ± 0.420	52°35'	$\frac{360^\circ}{4} = 90^\circ$	0.355
	" II ₁	49.50 ± 0.560	52°31'	0.183
		$\theta' = 88^\circ31/*$	$\frac{360^\circ}{4} = 90^\circ*$	0.183*

It is a remarkable fact that the differences of the values of θ between the different caryotypes III₁, III₂ and III₃ of the triploid *Paris* are nearly equal:

$$\text{III}_2 - \text{III}_3 = 41^\circ12' - 38^\circ18' = 2^\circ54',$$

$$\text{III}_1 - \text{III}_2 = 43^\circ24' - 42^\circ12' = 2^\circ12'.$$

As the caryological difference of these plants is seen in the number of the eliminated satellites, it may be suggested that the difference between the III₁ and III₂ plant or between the III₂ and III₃ plant indicates the degree of increase of the θ value, when a satellite is eliminated. If all e-satellites were eliminated, as in the case of the assumed III₀ plant, the θ value is expected to become equal to the sum of the θ value of the III₁ plant and the average of the difference of the θ value between the III₂ and III₃ plant, or between the III₁ and III₂ plant. It indicates:

$$43^\circ24' + \frac{2^\circ12' + 2^\circ54'}{2} = 45^\circ57$$

In the same way the θ value of the assumed II₀ plant is expected to be equal to the sum of the θ value of the II₁ plant and the difference of the θ value between the II₁ and II₂ plant, thus:

$$36^{\circ}30' + (36^{\circ}30' - 28^{\circ}20') = 44^{\circ}40'.$$

In both cases, diploid and triploid, the θ values of the plants without any satellite approximate to 45° . The leaves of *Parideae* are verticillate and arranged on a horizontal plane, and in *Paris hexaphylla* the leaf number of the well developed plant is ¹⁾ 8. Therefore, 45° is the maximum value of θ , assuming that 8 leaves grow radiating from an axis and are arranged on a plane without covering each other, because 45° is the quotient $\frac{360^{\circ}}{8}$, where 360° is the angle surrounding a point on a plane and 8 is the leaf number.

In the case of *P. tetraphylla*, the leaf number of the II₂ plant is always 4 and its mean θ value is $58^{\circ}35'$. Comparing these results with those of *P. hexaphylla* II₂, we can find that the leaf number of the former is just a half of the latter, while the θ value of the former is almost twice that of the latter. In the case of the II₁ plant of *P. tetraphylla*, however, the leaf number varies from 4 to 7 and averages 5.62. In the same plant the mean value of θ is $52^{\circ}21'$, which is smaller than the θ value of the II₂ plant. On the other hand, the mean value of θ of *P. hexaphylla* with less leaves is larger than that of the normal eight leaved plant. Moreover, in case of one individual²⁾ of *P. tetraphylla*, the leaf number varies from 4 to 6, so the mean value of θ in the four leaved plant is $83^{\circ}18'$, while it is $68^{\circ}40'$ in the five leaved one³⁾. In this case, we can find that the ratio of θ , $\frac{83^{\circ}18'}{68^{\circ}40'} = 1.214$, to be approximately equal to the inverse ratio of the leaf number $\frac{5}{4} = 1.25$. From these facts, it follows that the θ value varies proportionally to the reciprocal of the leaf number. Applying this idea to the case of *P. tetraphylla* II₁ plant, the calculated value of θ ⁴⁾ of the 4-leaved plant can be obtained, multiplying $52^{\circ}21'$ (the θ value of the II₁ plant with an average leaf number 5.62) by the inverse ratio of the leaf number; thus

1) Well developed *P. hexaphylla* has 8 leaves, and in the young or poorly developed ones there are less than 8, sometimes even 4. When the leaf number is less than 8 especially in young ones, the width of laminae is larger than in the 8 leaved.

2) Caryological study is not yet finished.

3) The θ value of the 6 leaved plant is omitted, because only one came to the writer's observation.

4) This is indicated by θ' .

$$\theta' = 52^\circ 21' \times \frac{5.62}{4} = 73^\circ 33'$$

this calculated value of θ' for the 4 leaved *P. tetraphylla* II₁ plant is almost twice that of the II₁ plant of *P. hexaphylla* ($36^\circ 21'$), while the leaf number of the former is a half that of the latter. Now we can calculate the θ value of the assumed *P. tetraphylla* II₀ plant with four leaves, in which all satellites are eliminated, according to the following formula:

$$73^\circ 33' + (73^\circ 33' - 58^\circ 35') = 88^\circ 31',$$

where $58^\circ 35'$ is θ of the four leaved II₂ plant, and $73^\circ 33'$ is θ of the four leaved II₁ plant. This θ value of *P. tetraphylla* II₀ plant is almost twice that of *P. hexaphylla* II₀ and III₀ plant, and it is nearly equal to the quotient $\frac{360^\circ}{4} = 90^\circ$, where 4 is the leaf number of this plant.

From the above mentioned results in *Paris*, it can be expected that the θ value becomes larger and approximates to the quotient $\frac{360^\circ}{\text{leaf number}}$, proportionally to the increase of the satellite elimination. If all satellites were eliminated, it will become equal to the quotient, which will be indicated hereafter by Δ for simplifying the description. So one can say that the θ value is limited below Δ by the presence of the satellite and then the degree of decrease of θ (indicated by λ) will be expressed by the following formula:

$$\lambda = \frac{\Delta - \theta}{\Delta} = 1 - \frac{\theta}{\Delta}$$

$$\text{or } \frac{\frac{360^\circ}{n} - \theta}{\frac{360^\circ}{n}} = 1 - \frac{n\theta}{360^\circ},$$

where n is the leaf number.

A remarkable similarity of the λ value between the same kinds of caryotype of both *P. hexaphylla* and *P. tetraphylla* is recognized (Table 2). The λ value of the II₂ plant in both *Paris* species is 0.372 and 0.355 respectively, and in the II₁ plants it is 0.189 and 0.183.

Therefore, it may be assumed that the λ value is closely related with the elimination of the satellite equally among the plants of different species used. On the other hand, as no decrease of the θ value would take place in the case of the II_0 or III_0 plant without satellite, one can expect that the θ value of these plants will become equal to Δ , namely that the λ value becomes equal to zero. This relation is easily recognized in a graph in which the λ values are taken as the co-ordinate, and the quotient, that is the number of eliminated




Fig. 5. Close relation between the angle of the two edges of the leaf margin at the base and the elimination of the satellite as indicated by a graph. Ph, *P. hexaphylla*. Pt, *P. tetraphylla*. Tk, *Trillium kamtschaticum*. TT, *T. tschonoskii*. TS. *T. Smallii*.

satellites is divided by the number of the e-chromosomes, as the abscissa. Two lines of *P. hexaphylla* (diploid and triploid) and the line of *P. tetraphylla* converge near the point A, where $\lambda = 0$ and the quotient is 1 (Fig. 5). But, as no material without a satellite, neither diploid nor triploid of both *Paris* species, came within the observation

of the writer, it is impossible to ascertain the presence of the point of the coincidence in *Paris*. *Trillium* has not only very similar chromosomal elements to those of *Paris*, but also the e-chromosome of the former genus is closely similar to the e-chromosome without satellite of *Paris*. So it may not be impossible to confirm the assumed relation, that the λ values become zero in the absence of a satellite, by the use of *Trillium* instead of *Paris*. In reality it was found that the λ values of *T. kamtschaticum*, *T. Tschonoskii* and *T. Smallii* are 0.032, 0.025 and 0.178 respectively (Table 3), where Δ of all *Trillium*

Table III.

Ratio of breadth to length of laminae, value of θ and degree of decrease of θ (λ) in *Trillium*.

	Ratio of breadth to length of lamina (%)	θ	$\Delta = \frac{360^\circ}{n}$	$\lambda = \frac{\Delta - \theta}{\Delta}$
<i>T. kamtschaticum</i> II ₀	106.90 \pm 0.558	116° 3' \pm 16'.5	$\frac{360^\circ}{3} = 120^\circ$	0.032
<i>T. Tschonoskii</i> VI ₀	106.22 \pm 0.622	116° 26' \pm 54'.9	„	0.027
<i>T. Smallii</i> IV ₀	106.28 \pm 0.467	98° 37' \pm 21'.1	„	0.178

plants is equally 120°. Thus the results obtained with *T. kamtschaticum* and *T. Tschonoskii* agree almost with the expectation, but in the case of *T. Smallii* the result is entirely contrary to it and the λ value is nearly equal to that of the II₁ plant of *Paris* (0.189 or 0.183). But it is a noticeable fact that the mean values of the ratio of the breadth to the length of the laminae of these three *Trillium* species nearly agree with each other, that is 106.90 ± 0.558 and 106.22 ± 0.622 , 106.28 ± 0.467 , respectively (Table 3). So the writer inclines to assume that a certain unknown factor, instead of the satellite, may exist, which is related with the value of θ at least in the case of *T. Smallii*.

Conclusion

With the recent advance of cytological investigations nuclear polymorphism was found in many plants and sometimes an interesting relation between the caryotypes and morphological character was ascertained. Certainly, this character can be distinguished both in the qualitative and the quantitative sense. The former is recogni-

zable rather easily, but in the latter case it is a matter of difficulty, because it needs many statistical studies even for a single character and sometimes it is rather difficult to find out its mode. Therefore, the investigations, undertaken hitherto in this direction, have been mostly of qualitative nature. The case of *Paris* is an example of the latter; the different caryotypes have different quantitative morphological characters¹⁾. The θ value, the angle between the two edges of the leaf margin at the base, is closely related to the satellite of the e-chromosome, and the λ value, the decrease of the θ value, is proportional to the presence of the satellite. It can be expected that θ reaches the maximum value and becomes equal to Δ , when all satellites are eliminated, where Δ is the quotient; $\frac{360^\circ}{\text{leaf number}}$. This expectation is supported powerfully by the fact, that θ of two species of *Trillium* are nearly equal to their Δ , where the chromosomal elements are similar to those of *Paris*, especially when there is an e-chromosome without satellite.

It is a noticeable fact that the maximum value of θ becomes equal to Δ , because the latter is the maximum size of the angle between the two edges of the leaf margin at the base, assuming that they do not cover each other, radiating from an axis and arranged on a plane. As the θ value will never become larger than Δ , even if all satellites were eliminated, the presence of the satellite may be considered as the secondary factor related to the decrease of the θ values, and one should probably assume a certain unknown primary factor which restricts the θ value below Δ . If one could suppose the primary existence of a satellite, it is impossible to explain why the λ values of those plants, in which all the e-chromosomes have satellites, are different and those of the plants with no satellite converge nearly to a point. So it is rather natural to expect that the satellite of the e-chromosome of *Paris* has been derived secondarily. It is not difficult to consider that the e-chromosome of *Paris* might be derived from a chromosome without satellite as in *Trillium*, having obtaining a mass of chromatin at the proximal end, as in the case of *Crepis*²⁾ where a fragment of a chromosome or satellite alone translocated to another chromosome. One might suppose that an e-chromosome with

1) It is not necessary to consider that these characters develop from the genes located on the chromosome.

2) M. NAVASHIN (1931, b and 1932), SCHKWARNIKOW u. M. NAVASHIN (1934).

satellite had been derived from a J-shaped chromosome, a constriction being newly produced in its shorter arm and its terminal portion being changed into a satellite. But it is not probable that such a change, which does not accompany the loss or gain of chromatin substance, is related with the change of the morphological character.

Summary

1) The satellite of the e-chromosome of *Paris hexaphylla* and *P. tetraphylla* is sometimes eliminated. The size of the angle between the two edges of the leaf margin at the base (θ) is closely related with this elimination of the satellite; the increase of the former is proportionate to that of the latter. The θ value approaches to the quotient: $\frac{360^\circ}{\text{leaf number}} = \Delta$ in accordance with the elimination of the satellite.

2) No plant of *Paris*, in which all satellites of the e-chromosomes are eliminated, is found. But it is expected that the angle between the two edges of the leaf margin at the base of such a plant will become equal to the quotient Δ , which is the maximum value of the angle between the two edges of the leaf margin at the base.

3) The chromosomal elements of *Trillium* are closely similar respectively to those of *Paris*, and especially the e-chromosome of the former strikingly resembles the e-chromosome without satellite of the latter. In *T. kamtschaticum* and *T. Tschoroskii*, the θ values are nearly equal to the calculated value $\Delta = \frac{360^\circ}{3} = 120^\circ$, where 3 is the leaf number. This fact is a powerful support for the expectation that the θ value of *Paris* without satellite will become equal to the quotient Δ .

4) It may be considered that the e-chromosome of *Paris* was derived from a chromosome without satellite, such as that of *Trillium*, by the addition of a chromatid to its proximal end.

It is the writers pleasure to express his hearty thanks to Prof. T. SAKAMURA under whose kind guidance and encouragement the present investigation was carried out.

Literature

- DELAUNAY, L. N. (1915 & 1925): (cited after DELAUNAY, 1929).
- (1929): Kern und Art. *Planta*, **7**. 110–112.
- GOTOH, K. (1933): Karyologische Studien an *Paris* und *Trillium*. *Jap. Journ. Gene.*, **8**. 197–203.
- GOTOH, K. and STOW, I. (1930): Caryological studies on *Paris* and *Trillium*. (preliminary report in Japanese). *Jap. Journ. Gene.*, **5**. 114–117.
- HAGA, T. (1934): The comparative morphology of the chromosome complement in the tribe *Parideae*. *Journ. Fac. Sci. Hokkaido Imp. Univ. Ser. V*, **3**. 1–32.
- (1935): On the karyotypes and their gametes of *Paris quadrifolia* L. var. *obovata* REGEL et TIL. (A preliminary note in Japanese with English résumé.). *Bot. Mag.*, **48**. 241–248.
- HUSKINS, C. L. and SMITH, S. G. (1935): Meiotic chromosome structure in *Trillium erectum*. *Ann. Bot.*, **69**. 119–150.
- KUHN, E. (1928): Zur Zytologie von *Thalictrum*. *Jahrb. f. wiss. Bot.*, **68**. 382–430.
- MATSUURA, H. (1934): On the number of spiral gyres in the chromonemata. (A preliminary note in Japanese). *Jap. Journ. Gene.*, **9**. 143–149.
- (1935): Chromosome studies on *Trillium kamtschaticum* PALL. I. The number of coils in the chromonema of the normal and abnormal meiotic chromosomes and its relation to the volume of chromosomes. *Cytologia*, **6**. 270–280.
- NAVASHIN, M. (1925): Morphologische Kernstudien der *Crepis*-Arten in bezug auf die Artbildung. *Zeitschr. f. Zellforsch. u. mikr. Anat.*, **2**. 98–111.
- (1926): Variabilität des Zellkerns bei *Crepis*-Arten in bezug auf die Artbildung. *Zeitsch. f. Zellforsch. u. mikr. Anat.*, **4**, 171–215.
- (1931 a and b): (cited after NAVASHIN, 1932).
- (1932): The dislocation hypothesis of evolution of chromosome numbers. *Zeitschr. f. ind. Abst.- u. Vererbgr.*, **63**. 224–234.
- NAWASCHIN, S. (1912): (cited after SHARP, 1934).
- (1915): (cited after DELAUNAY, 1929).
- SAKAMURA, T. (1920): Experimentelle Studien über die Zell- und Kernteilung mit besonderer Rücksicht auf Form, Grösse und Zahl der Chromosomen. *Journ. Coll. Sci. Imp. Univ. Tokyo.*, **39**. Art 11.
- SCHKWARNIKOW, P. K. u. NAWASCHIN, M. S. (1934): Über die Beschleunigung des Mutationsvorganges in ruhenden Samen unter dem Einflus von Temperaturhöhung. *Planta*, **22**. 720–736.
- SMITH, S. G. (1935): Chromosome fragmentation produced by crossing-over in *Trillium erectum* L. *Journ. Gene.*, **30**. 227–232.


Explanation of Plates

Plate I


- Figs. 6-9. Direct print of leaves. Fig. 10. Photograph. Reduced size $\frac{1}{2} \times$.
Fig. 11 a & b. Photomicrograph, iron-acetocarmine preparation, ca. 500 \times .
- 6. *Paris hexaphylla* II₂.
 - 7. " " II₁.
 - 8. " " III₁.
 - 9. *Paris tetraphylla* II₂.
 - 10. " " II₁.
 - 11. *Paris tetraphylla*, first nuclear division in the pollen grain, showing the stick shaped satellite. 750 \times .

Plate II


- Figs. 12-14. Direct prints of leaves, reduced size 1/3 \times .
Fig. 15. Photomicrograph, iron-acetocarmine preparation.
- 12. *Trillium kamtschaticum*.
 - 13. *Trillium Tschonoskii*.
 - 14. *Trillium Smallii*.
 - 15. *Trillium kamtschaticum*, first nuclear division in the pollen grain. 750 \times .
-




Stow : Correlation between Satellite and Leaf Shape.




12



13



14



15

Stow : Correlation between Satellite and Leaf Shape.