



Title	Two Metamorphic Events of the Nepal Himalayas Prior and Posterior to India-Eurasia Collision
Author(s)	Arita, Kazunori; Gautam, Pitambar; Ganzawa, Yoshihiro
Citation	北海道大学理学部紀要, 22(4), 519-528
Issue Date	1990-08
Doc URL	http://hdl.handle.net/2115/36763
Type	bulletin (article)
File Information	22_4_p519-528.pdf



[Instructions for use](#)

TWO METAMORPHIC EVENTS OF THE NEPAL HIMALAYAS PRIOR AND POSTERIOR TO INDIA-EURASIA COLLISION

by

Kazunori Arita, Pitambar Gautam* and Yoshihiro Ganzawa**

(with 3 text-figures and 2 tables)

Abstract

K-Ar ages were newly determined on biotite from a pelitic gneiss and on hornblende from a calcareous gneiss both from the Higher Himalayas in Nepal. The corresponding ages, 56.7 ± 2.8 Ma and 61.5 ± 3.1 Ma, respectively, are quite old compared with their fission-track ages.

We made a synthesis of the various isotopic ages from the Nepal Himalayas available so far, and attempted an analysis on the age versus closure temperature basis. The findings suggest that the Higher Himalayas have undergone two major reheating episodes, during the Himalayan orogeny, which are reflected in two different cooling trajectories roughly corresponding to late Cretaceous to Eocene and Eocene-Oligocene to the present times. A possible two-stage model involving two metamorphic events, which occurred prior and posterior to the India-Eurasia collision, is proposed, and the subsequent cooling histories are discussed. Implication of the (re-) magnetization events revealed by paleomagnetic data on the Nepal Lesser Himalayas to the model is also considered.

Introduction


The Himalayas represent a collisional product between Indian and Eurasian continents (Gansser, 1964; Powell and Conaghan, 1973; Le Fort, 1975). Most of the metamorphic and deformational features found in the Himalayas are considered to have been formed during the Himalayan orogeny, which resulted from the collision started in Eocene age (Frank et al., 1973; Le Fort, 1975), although some, especially in the Higher and Lesser Himalayas, show relics of an extensive pre-Himalayan tectono-metamorphism. At least three major intracontinental thrusts can be traced all over the Himalayas. From north to south, they are: Main Central Thrust (Gansser, 1964), Main Boundary Thrust (Gansser, 1964) and Himalayan Frontal Thrust (Nakata, 1972) with the related thrust activities progressively becoming younger in age towards the south. The Main Central Thrust is considered to be Miocene or older in age.

Isotopic geochronological studies on the Himalayan rocks reveal that the deter-

Contribution from the Department of Geology and Mineralogy, Faculty of Science, Hokkaido University, No. 1994

* Present address: Central Department of Geology, Tribhuvan University, Kathmandu, Nepal

** Institute of Earth Science, Hakodate College, Hokkaido University of Education, Hakodate, 040 Japan



Text-fig. 1 Histogram of isotopic ages determined by various methods in the Himalayas. Data are cited from the papers which are shown in references with star or double stars.

mined ages are concentrated mainly into the Tertiary, especially Miocene age (Text-fig. 1). At the same time, they also prove the Himalayas to have undergone a poly-metamorphism or poly-orogeny since the Precambrian (Crawford, 1981). The poly-metamorphism and poly-orogeny were inferred from the geological and petrological points of view (Hashimoto et al., 1973; Arita, 1983; Caby et al., 1983; Brunel and Kienast, 1986).


Mehta (1980) compiled the isotopic dating ages, and divided the Cretaceous-Tertiary Himalayan orogeny into three phases: (1) 75–50 Ma; folding and metamorphism, (2) 40–25 Ma; uplift, and (3) 25–10 Ma; major uplift, thrusting and regional retrogression.

The present paper discusses two events of metamorphism and the subsequent cooling history during Cretaceous to Tertiary age in the Nepal Himalayas based on newly determined K-Ar isotopic ages and zircon fission-track ages (Arita and Ganzawa, unpub. data) together with other isotopic ages already reported by various methods. Findings of the paleomagnetic research into the Lesser Himalayan rocks in the Tansen area (Gautam, 1989a; 1989b) are also discussed to support, though indirectly, the two-stage model.

Radiometric ages in the Nepal Himalayas

In Nepal, more than 160 age determinations measured by Rb/Sr and K/Ar methods have so far been reported mainly from the Higher Himalayas. A histogram of these ages, except for 36 older ages corresponding to Paleozoic to Precambrian time, is shown in Text-fig. 2. There are three distinct concentrations around 100–85 Ma, 70–45 Ma and 40–5 Ma. It is noteworthy that the trends of the histogram for the Nepal Himalayas closely resemble to those recognized by Mehta (1980) for the Himalayas as a whole.

Fifteen zircon fission-track ages, which were newly obtained from the Himalayan gneisses collected from the Higher Himalayas of central Nepal, are



Text-fig. 2 Histogram of isotopic ages from Cretaceous to Quaternary time determined by various methods in the Nepal Himalayas. Data are cited from the papers which are shown in references with double stars.

very young and range between 2.1 Ma and 0.6 Ma, except for one sample which is 7.7 Ma (Arita and Ganzawa, unpub. data).

Further, in an effort to establish the thermal history during the Tertiary, K-Ar age determinations were carried out on biotite from a pelitic gneiss and hornblende from a calcareous gneiss both collected from the upper stream of the Modi Khola river in central Nepal. The pelitic gneiss, which occurs in the upper part of the lower argillo-arenaceous unit of the Himalayan gneiss (Arita et al., 1982), consists mainly of kyanite, garnet, biotite, plagioclase and quartz. The calcareous gneiss,

Table 1 K-Ar ages on hornblende and biotite of the Himalayan gneisses in the Higher Himalayas, central Nepal.

Sample No.	Mineral	K (%)	$^{40}\text{Ar}^*$ (10^{-5}sc/gm)	Atm. ^{40}Ar (%)	Age (Ma)
HA80101624	Hornblende	1.29	0.322	76.6	61.5 \pm 3.1
		1.29	0.305	90.4	
HA7831510B	Biotite	7.32	1.63	84.5	56.7 \pm 2.8
		7.24	1.63	91.3	

$^{40}\text{Ar}^*$ is radiogenic argon. Ages were calculated with $\lambda_\beta = 4.962 \times 10^{-10}\text{yr}^{-1}$, $\lambda_\epsilon = 0.581 \times 10^{-10}\text{yr}^{-1}$, $^{40}\text{K}/\text{K} = 1.167 \times 10^{-4}\text{atom}\%$ and $^{40}\text{Ar}/^{36}\text{Ar}$ atmosphere = 295.5. Determined by Teledyne Isotopes, N. J., United States.

exposed 1 km upward from the pelitic gneiss, is collected from the lower part of the middle calcareous unit of the Himalayan gneiss (Arita et al., 1982), and is mostly composed of garnet, hornblende and plagioclase with small amount of calcite. These gneisses which gave fission-track ages of 2.1 and 2.2 Ma, respectively, yield K-Ar ages of 56.7 ± 2.8 Ma and 61.5 ± 3.1 Ma (Table 1). These ages are far older than those expected to be 20 to 10 Ma.

Closure temperatures of various geochronological methods

The closure temperature below which an isotopic clock of mineral or rock ceases from working is one of the most important factors in deciding the cooling history of rock, and is dependent on kinds of the age determination method and mineral. Many theoretical, experimental and geological studies have been carried out to settle the closure temperatures of different chronometries and different minerals (e. g., Dodson, 1979; Dodson and McClelland-Brown, 1985; Harrison, 1981; Harrison and McDougall, 1980). These studies show that the closure temperature is variable with variations of even up to 100°C or more due to the influence of many factors such as differences in kinematic behavior and grain size of minerals, and to hydraulic effects of pore fluids. However, the estimates of closure temperatures shown in Table 2 are commonly adopted (Jäger, 1979; Harrison and McDougall, 1980). The temperature of 230°C is used for the closure temperature of the zircon fission-track dating.


Table 2 Adopted closure temperatures (°C) for whole rock and various minerals by K-Ar and Rb-Sr methods.

	K-Ar method	Rb-Sr method
Whole rock	—	650
Hornblende	530	—
Muscovite	350	500
Biotite	280	320

Two metamorphic events and the subsequent cooling history in the Nepal Himalayas

The radiometric ages from the Nepal Himalayas are plotted against their closure temperatures (Text-fig. 3). These ages are obtained from central and eastern Nepal, and distributed in a wide area mainly of the Higher Himalayas. However, in the Himalayas it is well known that the rocks have undergone similar types of metamorphism and show monotonous and similar lithology especially in the Higher Himalayas (Gansser, 1964; Hashimoto et al., 1973; Le Fort, 1975).

This suggests that the Himalayas as a whole have suffered similar thermal



Text-fig. 3 Plot of radiometric ages derived by various methods and minerals from the Nepal Himalayas versus their closure temperatures. Components B and C show magnetization and remagnetization events in the Tansen area of the Lesser Himalayas (Gautam, 1989b). The age bar for the Rajmahal Traps is after McDougall and McElhinny (1971) and Baksi et al. (1987). The closure temperature levels are shown as horizontal lines, and height of columns is proportional to the number of ages corresponding to the particular interval. For clarity, however, the exact number is indicated inside the bar if it exceeds five.

history throughout the region. Therefore, the ages from various areas in Nepal which, during the late Eocene to Oligocene through the Miocene to the present, show younging with a corresponding decrease in closure temperature, are supposed to represent a trajectory of cooling history. A rather broad width of the trajectory can be attributed to the minor regional variation in cooling history. Another trajectory, although not so clear as the former, is also traceable from the late Cretaceous through the Paleocene to the Eocene. Thus, two cooling history passes are appreciable in the Nepal Himalayas. Some K-Ar ages on muscovite and biotite are found between the two trends. They can be considered to be rejuvenized ages of the older trajectory. Most of the K-Ar ages older than 85 Ma are obtained from detrital sericite and muscovite in sandstone and calcareous schist (Krummenacher, 1966; Krummenacher et al., 1978). The newly determined K-Ar ages for hornblende and biotite belong to the older trajectory. The estimated cooling rate based on these ages is 52°C/Ma. This value is twice as much as that of the Alps (Jäger, 1979). The cooling rate of the younger cooling pass is estimated to be almost the same to that of the older cooling pass.

In this paper, we consider the fission-track ages of 2.1 Ma and 2.2 Ma for gneisses to represent a later stage of the young cooling history. However, judging from the estimated cooling rate of about 50°C/Ma for the cooling trajectory, the ages appear to be young. In this context, it is worth noting that Gautam and Koshimizu (1988) suggested a possible strong thermal event around 5 Ma on the

basis of paleomagnetic and fission-track studies on the Ampipal alkaline massif in the Nepalese Lesser Himalayas.

Tectonic significance of two cooling trajectories

The metamorphism in the Himalayas, apart from that in Paleozoic and Precambrian times, is generally designated in age to the Tertiary, especially to the Miocene. The metamorphism in the pre-Tertiary is not yet proved geologically. However, two episodes of metamorphism have been petrologically recognized in the Nepal Himalays (Arita, 1983). Caby et al. (1983) and Brunel and Kienast (1986) inferred that the main metamorphism of barrovian type had occurred prior to thrusting on the Main Central thrust zone. We infer that the thrust movement gave metamorphism to the rocks in and around the Main Central Thrust zone, and brought about granitic magma in the depth. The young cooling trajectory is indicative of the metamorphism related to the thrusting during Oligocene to Miocene age. The old one seems to represent the cooling history of the metamorphism prior to the thrusting. It is noteworthy that these tectono-metamorphic events took place before and after the collision between India and Eurasia which occurred during 50 to 40 Ma.

The events visualized by two cooling trajectories are supported by the paleomagnetic data on the Nepalese Lesser Himalayan rocks. In the Tansen area of central Nepal, complex multi-component magnetic remanence consisting of at least four components, designated as components A, B, C and D, was recognized in the Aulis volcanics (Gautam, 1989a). The first three components could also be isolated from the sedimentary rocks in the same area (Gautam, 1989b). A mean direction of component A magnetization is consistent with the present dipole field at Tansen, and is of recent field origin. Timing of other components was inferred by comparing the paleolatitude values derived from the mean inclination with the calculated paleolatitude pattern at Tansen (27.9°N, 83.5°E) expected from the simulated Indian Apparent Polar Wander Path.

Component B, before any bedding tilt-correction, yields a most probable paleolatitude of 17°N suggesting its acquisition around 23 Ma. In volcanics, this component rests on a low Curie-temperature (360°C) mineral phase interpreted as (titano-) maghemite which represents a low-temperature (200–300°C) oxidation product of a Ti-rich magnetite. In red siderite and hematite-bearing sediments, this component rests on hematite which was probably formed due to oxidation of siderite during the time of the remanence acquisition. Thus, the Tansen area seems to have undergone a secondary moderate heating in the early Miocene, whereas the major folding or tectonic disturbance occurred prior to that time. This tectono-thermal picture might be a manifestation of the events responsible for the younger cooling trajectory in the Higher Himalayas.

Component C, of normal polarity, with thermally distributed unblocking temperatures up to 550–560°C suggests a paleolatitude of about 50°S after bedding tilt-

correction, Component D, almost antipodal to component C and hence of reverse polarity, with thermally discrete unblocking temperatures close to 550–560°C or above could be recognized only in few samples and so its quantitative estimation was rather poor. A high Curie-temperature (ca. 565°C) phase of Ti-poor magnetite carries the remanence corresponding to these components.

Sakai (1983) correlated the Aulis volcanics with the Rajmahal Traps and the sedimentary rocks of the Taltung Formation with the intertrappean sediments in the eastern part of the Indian shield north of Calcutta on geological grounds. Recent K-Ar dating of the Rajmahal Traps and associated volcanics by Bakshi et al. (1987) suggests that they were erupted ca. 115 Ma rather than 100–105 Ma based on previously determined K-Ar ages (McDougall and McElhinny, 1970). The high southern paleolatitude derived from component C and the occurrence of dual polarity components (i.e. component C probably acquired during the early time of Cretaceous Normal Polarity Superchron and component D acquired prior to it?) suggest that the volcanic activities in Peninsular India and Nepal Lesser Himalayas are coeval. At this point, we speculate that such volcanic activity in the Nepal Lesser Himalayas and the tectono-metamorphic event in the Higher Himalayas initiating the older cooling trajectory might be genetically interrelated.

Concluding remarks

An analysis of the Rb-Sr, K-Ar and fission-track zircon ages from the Nepal Himalayas (including the newly obtained K-Ar isotopic ages from two gneiss samples) and study on the age versus closure temperature relationship reveal that:

1) The isotopic ages, except those reflecting Quaternary uplift, are chiefly concentrated around 100–85 Ma, 70–45 Ma and 40–5 Ma.

2) These ages reflect two major cooling histories: the old one of late Cretaceous to Eocene, and the young one of late Eocene-Oligocene to the present. Two metamorphic events occurring prior and posterior to the India-Eurasia collision are suggested, accordingly.

The younger metamorphism and the resultant cooling history were undoubtedly caused by the activities related to an intracontinental thrust (the Main Central Thrust) and the following uplift. As evidenced by the secondary remagnetization episode, the latest maximum reheating event in the Lesser Himalayas dates back to the early Miocene. The initiation of the tectono-thermal event responsible for the older metamorphism is possibly coeval with the Aulis volcanic activity in the Lesser Himalayas. It is, however, unclear how such a metamorphism took place on the northern margin of the northward drifting Indian continent before the India-Eurasia collision.

Acknowledgements

This study was financially supported by a Grant-in-Aid for Overseas Scientific Survey (57041045) and a Grant-in-Aid for Scientific Research (61540550) of the

Ministry of Education, Science and Culture, Japan.

References⁺

- Arita, K., 1983. Origin of the inverted metamorphism of the Lower Himalayas, central Nepal. *Tectonophysics*, 95: 43-60.
- Arita, K., Hayashi, D. and Yoshida, M., 1982. Geology and structure of the Pokhara-Piuthan area, central Nepal. *J. Nepal Geol. Soc.*, 2 (Special Issue): 5-29.
- Baksi, A. K., Barman, T. R., Paul, D. K. and Farrar, E., 1987. Widespread Early Cretaceous flood basalt volcanism in eastern India: Geochemical date from the Rajmahal-Bengal-Sylhet Traps. *Chem. Geol.*, 63: 133-141.
- Bhandari, A. K. and Singh, K. N., 1979*. Granitic rocks of N.W. Himalaya and their geochronological status. *Geol. Surv. India, Misc. Pub.*, 41, Part I, 155-169.
- Bhanot, V. B., Goel, A. K., Singh, V. P. and Kawatra, S. K., 1975*. Rb-Sr radiometric studies for the Dalhousie and Rohtang areas, Himachal Pradesh. *Current Sci.*, 44: 219-220.
- Bhanot, V. B., Pandey, B. K., Singh, V. P. and Thakur, V. C., 1977a*. Rb-Sr whole-rock age of the granitic-gneiss from Askot area, eastern Kumaun and its implication on tectonic interpretation of the area. *Himal. Geol.*, 7: 118-122.
- Bhanot, V. B., Singh, V. P., Kansal, A. K. and Thakur, V. C., 1977b*. Early Proterozoic Rb-Sr whole-rock age for Central Crystalline Gneiss of Higher Himalaya, Kumaun. *J. Geol. Soc. India*, 18: 90-91.
- Bhanot, V. B., Kawatra, S. K., Kansal, A. K. and Pandey, B. K., 1978*. Rb-Sr whole rock age for Chail series of northwestern Himalaya. *J. Geol. Soc. India*, 19: 224-225.
- Bhanot, V. B., Bhandari, A. K., Singh, V. P. and Kansal, A. K., 1979*. Geochronological and geological studies on a granite of Higher Himalaya, northeast of Manikaran, H. P. *J. Geol. Soc. India*, 20: 90-94.
- Bhanot, V. B., Pandey, B. K., Singh, V. P. and Kansal, A. K., 1980*. Rb-Sr ages for some granitic and gneissic rocks of Kumaun and Himachal Himalaya. In: K. S. Valdiya and S. B. Bhatia (Editors), *Stratigraphy and correlations of Lesser Himalayan formations*. Hindustan Publishing Co., Delhi, 139-142.
- Bhattacharya, A. K., Bhatnagar, G. S., Narayan Das, G. R., Gupta, J. N., Chhabria, T. and Bhalala, N. S., 1982*. Rb/Sr dating and geological interpretation of sheared granitic-gneisses of Brijranigad-Ingedinala, Bhilangana valley, Tehri District, U. P. *Him. Geol.*, 12: 212-224.
- Brunel, M. and Kienast, J.-R., 1986. Étude pétro-structurale des chevauchements ductiles himalayens sur la transversale de l'Everest-Makalu (Népal oriental). *Can. J. Earth Sci.*, 23: 1117-1137.
- Caby, R., Pêcher, A. and Le Fort, P., 1983. Le grand chevauchement central himalayen: nouvelles données sur le métamorphisme inverse à la base de la Dalle du Tibet. *Revue de géologie dynamique et de géographie physique*, 24: 89-100.
- Crawford, A. R., 1981. Isotopic age data for the eastern half of the Alpine-Himalayan belt. In: H. K. Gupta and F. M. Delany (Editors). *Zagros, Hindu Kush, Himalaya, Geodynamic Evolution*. Geodynamic Series 3. American Geophysical Union, pp. 189-204.
- Dodson, M. H., 1979. Theory of cooling ages. In: E. Jäger and J. C. Hunziker (Editors), *Lectures in Isotope Geology*. Springer-Verlag, Berlin, pp. 194-202.
- Dodson, M. H. and McClelland-Brown, E., 1985. Isotopic and palaeomagnetic evidence for rates of cooling, uplift and erosion. *Mem. Geol. Soc.*, 10: 315-325.
- Ferrara, G., Lombardo, B. and Tonarini, S., 1983**. Rb/Sr geochronology of granites and gneisses from the Mount Everest region, Nepal Himalaya. *Geol. Rundschau*, 72: 119-136.
- Frank, W., Hoinkes, G., Miller, G., Purtscheller, F., Richter, W. and Thöni, M., 1973. Relations between metamorphism and orogeny in a typical section of the Indian Himalayas. *Tscherm. Min. Petr. Mitt.*, 20: 303-332.
- Frank, W., Thöni, M. and Purtscheller, F., 1977*. Geology and petrology of Kulu-south-Lahul area. *Colloques internat. C. N. R. C.*, No. 268, 147-172.

* * and ** represent the articles cited in Fig. 1 and Figs. 1 and 2, respectively.

- Gansser, A., 1964**. *Geology of the Himalayas*. Wiley-Interscience, New York, 289 pp.
- Gautam, P., 1989a. Multi-component remanent magnetization in the Aulis volcanics, the Lesser Himalaya, Nepal. *J. Geomag. Geoelectr.*, 41: 101-117.
- Gautam, P., 1989b. Magnetic properties of some late Paleozoic to Tertiary sedimentary rocks of Tansen area, Lesser Himalaya, Nepal. *J. Fac. Sci., Hokkaido Univ.*, Ser. IV, 22: 467-487.
- Gautam, P. and Koshimizu, S., 1989. Remanent magnetism and fission-track ages of some rocks from the Ampipal massif, the Lesser Himalaya, Nepal. *Abstracts of 1989 Annual Meeting of the Geological Society of Japan*. 640.
- Hamet, J. and Allègre, C. J., 1978**. Rb-Sr systematics in granite from central Nepal (Manaslu): Significance of the Oligocene age and high $^{87}\text{Sr}/^{86}\text{Sr}$ ratio in Himalayan orogeny: reply. *Geology*, 6: 197.
- Harrison, T. M., 1981. Diffusion of ^{40}Ar in hornblende. *Contrib. Mineral. Petrol.* 78: 324-331.
- Harrison, T. M. and McDougall, I., 1980. Investigations of an intrusive contact, northwest Nelson, New Zealand — I. Thermal, chronological and isotopic constraints. *Geochim Cosmochim Acta*, 44: 1985-2003.
- Hashimoto, S., Ohta, Y. and Akiba, C., 1973. *Geology of the Nepal Himalayas*. Saikon Publishing Co., Tokyo, 286pp.
- Jäger, E., 1979. Introduction to geochronology. In E. Jäger and J. C. Hunziker (Editors), *Lectures in Isotopic Geology*. Springer-Verlag, Berlin, pp. 1-12.
- Jäger, E., Bhandari, A. K. and Bhanot, V. B., 1971*. Rb-Sr age determinations on biotite and whole rock samples from the Mandi and Chor granites, Himachal Pradesh, India. *Eclogae geol. Helv.*, 64: 521-527.
- Kai, K. and Hayase, I., 1976**. Rb-Sr age and initial Sr isotopic ratio of some Himalayan rocks. *Mass Spectrometry*, 24: 181-188.
- Kai, K., 1981a**. Rb-Sr ages of the biotite and muscovite of the Himalayas, eastern Nepal: its implication in the uplift history. *Geochem. J.*, 15: 66-68.
- Kai, K., 1981b**. Rb-Sr geochronology of the rocks of the Himalaya, eastern Nepal. Part I: the metamorphic age of the Himalayan gneiss. *Mem. Fac. Sci., Kyoto Univ.*, Ser. Geol. & Mineral., 47: 135-148.
- Kai, K., 1981c**. Rb-Sr geochronology of the rocks of the Himalaya, eastern Nepal. Part II: the age and the origin of the granite on the Higher Himalayas. *Mem. Fac. Sci., Kyoto Univ.*, Ser. Geol. & Mineral., 47: 149-157.
- Khan, R. H. and Tater, J. M., 1970**. *Radiometric dates of some Nepalese rocks*. Nepal Geological Survey, Kathmandu, 6 pp.
- Krummenacher, D., 1961**. Déterminations d'âge isotopique faites sur quelques roches de l'Himalaya du Népal par la méthode potassium-argon. *Bull. Suisse Min. Petrogr.*, 41: 273-283.
- Krummenacher, D., 1966**. Népal central: géochronométrie des séries de l'Himalaya. *Bull. Suisse Min. Petrogr.*, 46: 43-54.
- Krummenacher, D., Kingery, F. A., Basett, A. M. and Layne, H. F., 1978**. Petrology, metamorphism and K/Ar age determinations in eastern Nepal. In: P. S. Saklani (Editor), *Tectonic Geology of the Himalaya*. Today and Tomorrow's Printers & Publishers, New Delhi, pp. 151-166.
- Laboratory of Isotopic Geology, Institute of Geochemistry, Academia Sinica, 1973**. Isotopic dating of the metamorphic rocks from the Mt. Jolmo Lungma area, China. *Scientia Sinica*, 16: 385-395.
- Le Fort, P., 1975. Himalayas: the collided range. *Am. J. Sci.*, 275A: 1-44.
- Le Fort, P., Debon, F. and Sonet, J., 1980*. The "Lesser Himalayan" cordierite granite belt typology and age of the pluton of Manserah (Pakistan). *Sp. Iss. Geol. Bull. Univ. Peshawar*, vol. 13, 51-61.
- McDougall, I. and McElhinny, M. W., 1970. The Rajmahal Traps of India. K-Ar ages and palaeomagnetism. *Earth Planet. Sci. Lett.*, 9: 371-378.
- Mehta, P. K., 1977*. Rb-Sr geochronology of the Kulu-Mandi Belt: its implications for the Himalayan tectogenesis. *Geol. Rundschau*, 66: 156-175.
- Mehta, P. K. and Rex, D. C., 1977*. K-Ar geochronology of the Kulu-Mandi Belt, NW Himalaya,

- India. *N. Jb. Miner. Mh.*, 343-355.
- Mehta, P. K., 1980. Tectonic significance of the young mineral dates and the rates of cooling and uplift in the Himalaya. *Tectonophysics*, 62: 205-217.
- Nakata, T., 1972. Geomorphologic history and crustal movements of the foot-hills of the Himalayas. *Sci. Rep. Tohoku Univ.*, 7th Ser. (Geogr.), 22: 39-177.
- Pande, I. C. and Kumar, S., 1974*. Absolute age determinations of crystalline rocks of Manali-Jaspa region, northwestern Himalaya. *Geol. Rundschau*, 63: 539-548.
- Powell, C. McA. and Conaghan, P. J., 1973. Plate tectonics and the Himalayas. *Earth Planet. Sci. Lett.*, 20: 1-12.
- Powell, C. McA., Crawford, A. R., Armstrong, R. L., Prakash, R. and Wynne-Edwards, H. R., 1979*. Reconnaissance Rb-Sr dates for the Himalayan Central Gneiss, northwest India. *Indian J. Earth Sci.*, 6: 139-151.
- Raju, B. N. V., Chabria, T., Prasad, R. N., Mahadevan, T. M. and Bhalla, N. S., 1982*. Early Proterozoic Rb-Sr isochron age for Central Crystalline rocks, Bhilangana valley, Garhwal Himalaya. *Him. Geol.*, 12: 196-205.
- Sakai, H., 1983. Geology of the Tansen Group of the Lesser Himalaya in Nepal. *Mem. Fac. Sci., Kyushu Univ.*, ser. D, Geol., 25: 27-74.
- Sinha, A. K., 1977*. Geochronology, petrography, petrochemistry and tectonic significance of basic rock suites of north-western Himalaya with special reference to Himachal Himalaya, India. In: B. S. Tewari and V. J. Gupta (Editors), *Recent Researches in Geology*, 3. Hindustan Publishing Co., Delhi, pp. 478-494.
- Sinha, A. K. and Bagdasarian, G. P., 1977*. Potassium-Argon dating of some magmatic and metamorphic rocks from Tethyan and Lesser zones of Kumaun and Garhwal Indian Himalaya and its implication in the Himalayan tectogenesis. *Colloques internat. C. N. R. S.*, no. 286. pp. 387-394.
- Trivedi, J. R. Gopalan, K. and Valdiya, K. S., 1984*. Rb-Sr ages of granitic rocks within the Lesser Himalayan Nappes, Kumaun, India. *J. Geol. Soc. India*. 25: 641-654.
- Wager, L. R., 1965**. Injected granite sheets of the Rongbuk valley and the North Face of Mount Everest. *Min. Geol. Met. Inst. India*, D. N. Wadia commemorative volume, 358-380.

(Manuscript received on May 29, 1989; and accepted on Oct. 27, 1989).