

HOKKAIDO UNIVERSITY

Title	On the commutativity of fundamental groups of complements to plane curves
Author(s)	SHIMADA, ICHIRO
Citation	Mathematical Proceedings of the Cambridge Philosophical Society, 123, 49-52 https://doi.org/10.1017/S0305004197002107
Issue Date	1998
Doc URL	http://hdl.handle.net/2115/5794
Rights	Copyright © 1998 Cambridge University Press
Туре	article
File Information	MPCPS123.pdf



# On the commutativity of fundamental groups of complements to plane curves

#### By ICHIRO SHIMADA

Department of Mathematics, Faculty of Science, Hokkaido University, Sapporo 060 e-mail: shimada@math.hokudai.ac.jp

(Received 14 February 1996; revised 7 June 1996)

#### Introduction

In this paper, we prove the following:

THEOREM. Let  $C \subset \mathbb{P}^3$  be a complex reduced irreducible space curve which is nondegenerate (i.e., C is not contained in any plane). Let  $p: C \to \mathbb{P}^2$  be a general projection. Then the topological fundamental group  $\pi_1(\mathbb{P}^2 \setminus p(C))$  is abelian.

When C is non-singular, the image of the general projection has only nodes as its singularities, and hence the complement has an abelian fundamental group thanks to Fulton-Deligne's theorem on Zariski conjecture ([1, 2]). The point of our theorem is that we make no assumptions on the singularities of the space curve C.

We apply this theorem to the problem of commutativity of fundamental groups of complements to plane curves of a given degree with prescribed numbers of nodes and cusps. Here a cusp means a germ of curve singularity which is analytically isomorphic to a small neighborhood of the singular point of the affine curve  $x^2 + y^3 = 0$ . The strongest result concerned with this problem is Nori's result [5, proposition 6.5]. (Nori considered not only plane curves with only nodes and cusps but curves on arbitrary surfaces with any kind of singularities.) Let  $C \subset \mathbb{P}^2$  be a reduced irreducible plane curve of degree d whose singular locus consists of n nodes and k cusps.

THEOREM (Nori). Suppose that  $2n + 6k < d^2$ . Then  $\pi_1(\mathbb{P}^2 \setminus C)$  is abelian.

We will prove a proposition which enlarges slightly the region of (d, n, k) on which the commutativity of  $\pi_1(\mathbb{P}^2 \setminus C)$  is guaranteed.

PROPOSITION 1. Suppose that  $2n \ge d^2 - 5d + 8$ . Then  $\pi_1(\mathbb{P}^2 \setminus C)$  is abelian.

## 1. Proof of Theorem

Let G denote the Grassmannian variety Grass  $(\mathbb{P}^1, \mathbb{P}^3)$  of all lines in  $\mathbb{P}^3$ , and let  $(\mathbb{P}^3)^*$  be the dual projective space Grass  $(\mathbb{P}^2, \mathbb{P}^3)$  of  $\mathbb{P}^3$ . For a plane  $\Pi \in (\mathbb{P}^3)^*$ , we put

$$\Sigma(\Pi) \coloneqq \{l \in G; \ l \subset \Pi\}$$

We also put

$$U\!\!\coloneqq\!\{l\in G;\ C\cap l=\emptyset\}.$$

When  $\Pi \in (\mathbb{P}^3)^*$  is general, no three points among  $C \cap \Pi$  are co-linear. Indeed, let  $B \subset (\mathbb{P}^3)^*$  be the Zariski open dense subset consisting of the planes which intersect C

## ICHIRO SHIMADA

at its non-singular points transversely. Let  $\Pi_0 \in B$  be a plane which serves as a base point of B. Since C is irreducible, the monodromy group of the action of  $\pi_1(B, \Pi_0)$ on the set  $C \cap \Pi_0$  is the full symmetric group, because the action is 2-transitive and the monodromy group contains a simple transposition (cf. [4, Uniform Position Lemma]). Hence if there exist co-linear three points among  $C \cap \Pi$  for a general  $\Pi \in$  $(\mathbb{P}^3)^*$ , then all points in  $C \cap \Pi$  are on a line for a general  $\Pi$ . Note that this condition is closed for the choice of  $\Pi$  in  $(\mathbb{P}^3)^*$  because C is non-degenerate. Thus all points in  $C \cap \Pi$  are on a line for every  $\Pi$ , which is absurd. Consequently,  $\Sigma(\Pi) \setminus U$  is a union of lines on  $\Sigma(\Pi) \cong \mathbb{P}^2$ , which is a normal crossing divisor when the plane  $\Pi$  is generally chosen. Hence  $\pi_1(\Sigma(\Pi) \cap U)$  is abelian for a general  $\Pi$  because of Fulton–Deligne's theorem ([1-3]). Let  $I \subset (\mathbb{P}^3) * \times G$  be the closed subvariety  $\{(\Pi, l); \Pi \supset l\}$ . The second projection  $\rho_2: I \to G$  is a fibre bundle with fibres isomorphic to  $\mathbb{P}^1$ . Hence  $\rho_2$ induces an isomorphism  $\pi_1(\rho_2^{-1}(U)) \cong \pi_1(U)$ . Consider the projection  $\rho_1: \rho_2^{-1}(U) \to (\mathbb{P}^3)^*$ . The fibre  $\rho_1^{-1}(\Pi) \subset \rho_2^{-1}(U)$  over  $\Pi \in (\mathbb{P}^3)^*$  is isomorphic to  $\Sigma(\Pi) \cap U$ , which is non-empty and non-singular for every  $\Pi$ . We apply Nori's Lemma [5, lemma 1.5 (C)] to conclude that the inclusion of the fibre induces a surjection  $\pi_1(\Sigma(\Pi) \cap U) \rightarrow$  $\rightarrow \pi_1(\rho_2^{-1}(U))$  for a general  $\Pi$ . Hence  $\pi_1(\rho_2^{-1}(U))$  is abelian, and thus  $\pi_1(U)$  is also abelian.

We fix a plane  $\Pi_0 \subset \mathbb{P}^3$  and denote by A the complement  $\mathbb{P}^3 \setminus \Pi_0$ . Let  $J_A \subset A \times G$  be the closed subvariety  $\{(P, l); P \in l\}$ . Consider the projection  $\tau_2: J_A \to G$ . If  $l \notin \Pi_0$ , then  $\tau_2^{-1}(l)$  is isomorphic to  $\mathbb{A}^1$ , while if  $l \subset \Pi_0$ , then  $\tau_2^{-1}(l)$  is empty. Thus  $\tau_2$  induces an isomorphism  $\pi_1(\tau_2^{-1}(U)) \cong \pi_1(U \setminus \Sigma(\Pi_0))$ . Since  $\Sigma(\Pi_0)$  is of codimension 2 in G, we have  $\pi_1(U \setminus \Sigma(\Pi_0)) \cong \pi_1(U)$ . Therefore,  $\pi_1(\tau_2^{-1}(U))$  is isomorphic to  $\pi_1(U)$ . We have an isomorphism  $J_A \cong A \times \Pi_0$  given by  $(P, l) \mapsto (P, l \cap \Pi_0)$ . Let  $\tilde{\tau}_2: A \times \Pi_0 \to G$  be the morphism corresponding to  $\tau_2: J_A \to G$  via this isomorphism. Let  $D \subset G$  be the Zariski closed subset  $G \setminus U$ . We provide it with the reduced structure. For  $P \in A$ , consider the scheme theoretic intersection

$$D_P \coloneqq (\{P\} \times \Pi_0) \cap \tilde{\tau}_2^{-1}(D),$$

which is considered as a sub-scheme of  $\Pi_0$ . If  $P \in C$ , then  $D_P = \Pi_0$ . Suppose that  $P \notin C$ . Let  $p\langle P \rangle \colon C \to \Pi_0$  denote the projection with the centre P. Then we see that the reduced part  $(D_P)_{\text{red}}$  of  $D_P$  coincides with the reduced part  $p\langle P \rangle(C)_{\text{red}}$  of the image of the projection  $p\langle P \rangle$ . Hence  $D_P$  is a reduced divisor of  $\Pi_0$  if deg  $p\langle P \rangle(C) = \deg C$ .

Claim. There are at most finitely many points  $P \in A \setminus C$  such that the degree of  $p\langle P \rangle(C)_{\text{red}}$  is less than the degree of C.

Since  $C \cap A \subset A$  is of codimension 2, Claim implies that the locus of all points  $P \notin A$  such that the scheme theoretic intersection  $D_P$  is *not* a reduced divisor of  $\Pi_0$  is of codimension 2 in A. Let  $P \in A$  be generally chosen. Applying [6, theorem 1], we get  $\pi_1(\Pi_0 \setminus p \langle P \rangle(C)) \cong \pi_1(\tilde{\tau}_2^{-1}(U)) \cong \pi_1(U)$ . Therefore  $\pi_1(\Pi_0 \setminus p \langle P \rangle(C))$  is abelian, and Theorem is proved.

Proof of Claim. Suppose that there exists an irreducible curve  $\Xi \subset A \setminus C$  such that deg  $p\langle P \rangle(C)_{\text{red}} < \deg C$  for all  $P \in \Xi$ . Let  $P_0$  be a general point of  $\Xi$  and  $Q_0$  a general point of C. Since  $P_0$  is not on C, the projection  $p\langle P_0 \rangle: C \to \Pi_0$  with the centre  $P_0$ must be of mapping degree  $\geq 2$  onto its image. Thus there exists a point  $R_0 \in C$  such

## Fundamental groups of complements to plane curves

that  $Q_0 \neq R_0$  and  $p\langle P_0 \rangle(Q_0) = p\langle P_0 \rangle(R_0)$ . Since  $Q_0$  is chosen generally,  $p\langle P_0 \rangle(Q_0)$  is a non-singular point of  $p\langle P_0 \rangle(C)_{\text{red}}$ , and the morphism  $p\langle P_0 \rangle: C \to p\langle P_0 \rangle(C)_{\text{red}}$  is étale at  $Q_0$  and at  $R_0$ . This implies that the tangent lines  $l(Q_0)$  and  $l(R_0)$  of C at  $Q_0$  and at  $R_0$ , respectively, intersect each other in  $\mathbb{P}^3$ . Since  $P_0$  is also chosen generally on  $\Xi$ , there exist small open neighbourhoods (in the sense of complex analytic geometry)  $V \subset \Xi$  of  $P_0$  in  $\Xi$  and  $W \subset C$  of  $R_0$  in C, and an isomorphism  $f: V \xrightarrow{\sim} W$  such that  $p\langle P \rangle(Q_0) = p\langle P \rangle(f(P))$  for all  $P \in V$ . This implies that the tangent line l(R) of C at R intersects  $l(Q_0)$  for all  $R \in W$ . Therefore  $l(R) \cap l(Q_0) \neq 0$  for every non-singular point R of C. However, consider the projection  $\lambda: C \to \mathbb{P}^1$  with the centre  $l(Q_0)$ . Since C is non-degenerate,  $\lambda$  is surjective, and hence, by Sard's theorem,  $\lambda$  is smooth at a general point  $R \in C$ . This implies that  $l(R) \cap l(Q_0) = 0$  for a general  $R \in C$ , and we get a contradiction.

# 2. Proof of Proposition 1

First note that if  $2n \ge d^2 - 5d + 8$  and  $2n + 2k < d^2 - 4d + 3$ , then 2n + 6k is automatically less than  $d^2$ . Therefore, taking Nori's theorem into account, we may assume that the inequality  $2n + 2k \ge d^2 - 4d + 3$  also holds.

Let  $\nu: \tilde{C} \to C$  be the normalization of C. The genus of  $\tilde{C}$  is given by

$$g = (d-1)(d-2)/2 - n - k.$$

The assumed inequalities imply that

$$d \ge \max(2g+1, g+3), \quad \text{and} \quad k \le d-g-3. \tag{2.1}$$

Let L be the pull-back of  $\mathcal{O}_C(1)$  by  $\nu$ , which is an invertible sheaf of degree d on  $\tilde{C}$ . By the first inequality of (2.1), L is very ample and dim |L| = d - g. Let C' be the image of the embedding of  $\tilde{C}$  in  $\mathbb{P}^{d-g}$  by L. Then C is obtained by a certain projection  $C' \to \mathbb{P}^2$ . Let  $\Gamma \subset \mathbb{P}^{d-g}$  be the centre of this projection, which is a linear subspace of codimension 3. Let  $T \subset \mathbb{P}^{d-g}$  be the union of all lines tangent to C'. Then  $\Gamma$  intersects T at distinct k points  $P_1, \ldots, P_k$ , each of which corresponds to a cusp of C bijectively. By the second inequality of (2.1), there exists a linear subspace  $\Lambda \subset \Gamma$ of codimension 1 which contains  $P_1, \ldots, P_k$ . Let  $p(\Lambda): C' \to \mathbb{P}^3$  be the projection with the centre  $\Lambda$ . The image  $C'' \subset \mathbb{P}^3$  is a non-degenerate curve, which has at least k singular points  $Q_1, \ldots, Q_k$  corresponding to  $P_1, \ldots, P_k \in \Lambda \cap T$ . The projection  $p\langle \Gamma \rangle : C' \to C$  with the centre  $\Gamma$  factors as  $C' \to C'' \to C$ , and  $C'' \to C$  maps  $Q_1, \ldots, Q_n$  $Q_k$  to the cusps of C bijectively. Therefore, C'' has k cusps and some nodes (possibly none) as its only singularities. Thus the image p(C'') of the general projection of C''to  $\mathbb{P}^2$  has the same numbers and types of singular points as the original curve C. Note that C and p(C'') are contained in an irreducible equisingular family of plane curves, every member of which is obtained as an image of a projection from  $C'' \subset \mathbb{P}^3$ to  $\mathbb{P}^2$ . Thus our main theorem implies that  $\pi_1(\mathbb{P}^2 \setminus C)$  is abelian.

#### 3. Examples

Let  $\mathscr{A}$  denote the set of triples (d, n, k) with  $d \in \mathbb{Z}_{>0}$ ,  $n \in \mathbb{Z}_{\geq 0}$  and  $k \in \mathbb{Z}_{\geq 0}$  for which the following hold:

(i) There exists a reduced irreducible plane curve  $C \subset \mathbb{P}^2$  of degree d with n nodes and k cusps as its only singularities.

# Ichiro Shimada

(ii) For any plane curve C as in (i),  $\pi_1(\mathbb{P}^2 \setminus C)$  is abelian.

Nori's result says that if (d, n, k) satisfies (i) and  $2n + 6k < d^2$ , then  $(d, n, k) \in \mathscr{A}$ . Using Proposition 1, we shall enlarge the known region of  $\mathscr{A}$ .

Let  $C \subset \mathbb{P}^n$  be a non-degenerate irreducible curve of degree d whose singular locus consists of only nodes and cusps. Let k be the number of cusps. We write by Tan  $(C) \subset \mathbb{P}^n$  the union of all lines in  $\mathbb{P}^n$  tangent to C at its non-singular points. The following proposition is elementary:

PROPOSITION 2. If P is a general point of Tan (C), then the image of the projection  $C \to \mathbb{P}^{n-1}$  with the centre P is of degree d, has only nodes and cusps as its singularities, and the number of cusps is k + 1.

Let C be a non-singular curve of genus g, and L a line bundle on C of degree  $d \ge \max (2g + 1, g + 3)$ . We embed C into  $\mathbb{P}^{d-g}$  by |L|. Applying Proposition 2 to  $C \subset \mathbb{P}^{d-g}$  repeatedly, we obtain a non-degenerate space curve  $C' \subset \mathbb{P}^3$  of degree d with certain number of nodes (possibly none) and d - g - 3 cusps as its only singularities. Let  $C'' \subset \mathbb{P}^2$  be the image of a general projection  $C' \to \mathbb{P}^2$ . Then C'' is an irreducible plane curve of degree d with

$$n = (d^2 - 5d + 8)/2$$
 nodes and  $k = d - g - 3$  cusps

as its only singularities. Since  $2n \ge d^2 - 5d + 8$ , Proposition 1 implies that  $(d, n, k) \in \mathscr{A}$ . Thus, choosing a line bundle of degree  $\ge 6g + 10$ , we obtain a region of  $\mathscr{A}$  not covered by Nori's theorem.

Note that there is another known region of  $\mathscr{A}$ , which is given by the following theorem due to Zariski [7]. Let  $R(d,k) \subset \mathbb{P}_*(\Gamma(\mathbb{P}^2, \mathcal{O}(d)))$  be the locus of all *rational* plane curves C of degree d with n nodes and k cusps as its only singularities. Since the genus of the normalization of C is 0, we have n = (d-1)(d-2)/2 - k.

THEOREM (Zariski). (1) The locus R(d, k) is non-empty if and only if  $k \leq 3(d-2)/2$ . (2) For a member C of R(d, k),  $\pi_1(\mathbb{P}^2 \setminus C)$  is abelian unless d is even and k = 3(d-2)/2.

Thus, if n + k = (d - 1)(d - 2)/2 and  $k \leq (3d - 7)/2$ , we have  $(d, n, k) \in \mathscr{A}$ .

Acknowledgement. I would like to thank Max-Planck-Institut für Mathematik in Bonn for providing me with the stimulating research environment. This work was done during my stay at the Institute in 1995.

#### REFERENCES

- P. DELIGNE. Le groupe fondamental du complément d'une courbe plane n'ayant que des points doubles ordinaires est abélien. Sém. Bourbaki, no. 543, 1979/80, Lecture Notes in Math., vol. 842 (Springer-Verlag 1981), 1–10.
- [2] W. FULTON. On the fundamental group of the complement of a node curve. Ann. Math. 111 (1980), 407–409.
- [3] W.FULTON and R. LAZARSFELD. Connectivity and its applications in algebraic geometry. Lecture Notes in Math., vol. 862 (Springer-Verlag, 1981), 26–92.
- [4] J. HARRIS. The genus of space curves. Math. Ann. 249 (1980), 191–204.
- [5] M. NORI. Zariski's conjecture and related problems. Ann. Sci. École Norm. Sup. (4), 16 (1983), 305–344.
- [6] I. SHIMADA. Fundamental groups of open algebraic varieties. Topology, 34 (1995), 509-532.
- [7] O. ZARISKI. On the Poincaré group of rational plane curves. Amer. J. Math., 58 (1936), 607–619; Collected Papers, Volume 3 (The MIT Press, 1978), 266–278.

52