

| Title            | Motion of a Graph by R-Curvature   |
|------------------|--|
| Author(s)        | Ishii, Hitoshi; Mikami, Toshio   |
| Citation         | Archive for Rational Mechanics and Analysis, 171(1), 1-23<br>https://doi.org/10.1007/s00205-003-0294-1 |
| Issue Date       | 2004-01  |
| Doc URL          | http://hdl.handle.net/2115/5879  |
| Rights           | The original publication is available at www.springerlink.com  |
| Туре             | article (author version)   |
| File Information | ARMA171-1.pdf  |



# Motion of a graph by R-curvature

Нітозні Ізнії\*, Тозніо Мікамі\*\*

#### Abstract

We show the existence of weak solutions to the PDE which describes the motion by R-curvature in  $\mathbf{R}^d$ , by the continuum limit of a class of infinite particle systems. We also show that weak solutions of the PDE are viscosity solutions and give the uniqueness result on both weak and viscosity solutions.

### 1. Introduction

In [8] Firey proposed a mathematical model of the wearing process of a convex stone rolling on a beach. In his model a stone evolves according

<sup>\*</sup> supported in part by the Grant-in-Aid for Scientific Research, No. 12440044 and No. 12304006, JSPS.

<sup>\*\*</sup> supported in part by the Grant-in-Aid for Scientific Research, No. 13640096 and 12440044, JSPS.

to the Gauss curvature flow. (see, e.g., [1, 4, 5, 17] for the mathematical developments regarding the Gauss curvature flow).

The crystalline approximation of a simple closed convex curve which evolves according to the curvature flow was considered by Girão and Kohn and is useful in the numerical analysis (see [10, 11]). We refer to [9] and the references therein for the recent development of this topics.

In [14] we proposed and studied a two-dimensional random version of [10]. We also generalized Firey's argument to the case when the stone does not necessarily have a convex shape and when the boundary of the stone is given by the graph of an evolving function (see [15]).

In this paper we propose and study the stochastic approximations of evolving functions which are generalizations of Gauss curvature flow considered in [15].

For  $u \in C(\mathbf{R}^d)$  and  $x \in \mathbf{R}^d$ , the following set is called the subdifferential of u at x:

$$\partial u(x) \equiv \{ z \in \mathbf{R}^d : u(y) - u(x) \ge \langle z, y - x \rangle \text{ for all } y \in \mathbf{R}^d \}, \quad (1.1)$$

where  $\langle \cdot, \cdot \rangle$  denotes the inner product in  $\mathbf{R}^d$ . For a set  $A \subset \mathbf{R}^d$  and a function  $v : A \mapsto \mathbf{R}$ , let  $\operatorname{epi}(v)$  and  $\hat{v}$  denote, respectively, the epi-graph of v, i.e., the set  $\{(x, y) : x \in A, y \ge v(x)\}$ , and the convex envelope of v, i.e., the function whose epi-graph is the convex hull of  $\operatorname{epi}(v)$ , provided that it exists.

For  $R \in L^1(\mathbf{R}^d : [0, \infty), dx)$ , *R*-curvature (or a generalized Gauss curvature measure) which can be defined as follows plays a crucial role in this paper.

**Definition 1 (***R***-curvature).** Let  $R \in L^1(\mathbf{R}^d : [0,\infty), dx)$ . For  $u \in C(\mathbf{R}^d)$ , we define the *R*-curvature of (a graph of) u as the finite Borel measure w(R, u, dx) on  $\mathbf{R}^d$  given by (see e.g. [2, section 9.6]):

$$w(R, u, A) \equiv \int_{\bigcup_{x \in A} \partial u(x)} R(y) dy \quad \text{for all Borel } A \subset \mathbf{R}^d.$$
(1.2)

**Remark 1.** (i) w in (1.2) has another expression:

$$w(R, u, A) = \int_{\bigcup_{x \in A} \partial \hat{u}(x)} R(y) dy \quad \text{for all Borel } A \subset \mathbf{R}^d$$
(1.3)

since the Lebesgue measure of the set  $\cup_{x \in \mathbf{R}^d} \{ p \in \partial \hat{u}(x) : p \text{ is singular} \}$ is zero  $(p \in \partial \hat{u}(x)$  is called singular if  $\{(y, \hat{u}(x) + \langle p, y - x \rangle) : y \in \mathbf{R}^d\} \cap \operatorname{epi}(\hat{u})$  contains at least two different points.) (see [2, section 9.4]). (ii) w(1, u, dx) is also called the Monge-Ampère measure associated with u and is useful in the study of the Monge-Ampère equation (see [2, 12]).

By the continuum limit of a class of infinite particle systems, we first show the existence of the solution to the following equation (see Theorem 1 in Sect. 2).

**Definition 2 (Motion by** *R*-curvature). The graph of  $u \in C([0, \infty) \times \mathbf{R}^d)$  is called the motion by *R*-curvature if the following holds: for any  $\varphi \in C_o(\mathbf{R}^d)$  and any  $t \ge 0$ ,

$$\int_{\mathbf{R}^d} \varphi(x) u(t, x) dx - \int_{\mathbf{R}^d} \varphi(x) u(0, x) dx \qquad (1.4)$$
$$= \int_0^t ds \int_{\mathbf{R}^d} \varphi(x) w(R, u(s, \cdot), dx).$$

Roughly speaking, our infinite particle systems  $\{(Z_n(t,z))_{z\in \mathbb{Z}^d/n}\}_{t\geq 0}$ satisfy that for any  $t\geq 0$  and any  $z\in \mathbb{Z}^d/n$ ,

$$P(Z_n(t + \Delta t, z) - Z_n(t, z) > 0) \propto E[w(R, \hat{Z}_n(t, \cdot), \{z\})]\Delta t + o(\Delta t)$$

as  $\Delta t \to 0 \ (n \ge 1)$ , where  $\hat{Z}_n(t, \cdot)$  denotes a convex envelope of the function  $z \mapsto Z_n(t, z)$  (see Sect. 2).

We also show the uniqueness result on and elementary properties of the solutions to (1.4) (see Theorems 1 and 2 in Sect. 2).

Theorem 3 in Sect. 2 shows that a continuous solution to (1.4) is a viscosity solution of the following PDE:

$$\partial u(t,x)/\partial t = \chi(u, Du(t,x), t, x) \operatorname{Det}_+(D^2 u(t,x)) R(Du(t,x)), \qquad (1.5)$$

where  $Du(t,x) \equiv (\partial u(t,x)/\partial x_i)_{i=1}^d, D^2u(t,x) \equiv (\partial^2 u(t,x)/\partial x_i\partial x_j)_{i,j=1}^d,$ 

$$\chi(u, p, t, x) \equiv \begin{cases} 1 & \text{if } p \in \partial u(t, x), \\ 0 & \text{otherwise,} \end{cases}$$

 $\partial u(t,x)$  denotes the subdifferential of the function  $x \mapsto u(t,x)$ , and for a real  $d \times d$ -symmetric matrix X,

$$\operatorname{Det}_{+} X \equiv \begin{cases} \operatorname{Det} X & \text{if } X \text{ is nonnegative definite,} \\ 0 & \text{otherwise.} \end{cases}$$

A continuous viscosity solution u to (1.5) is the Gauss curvature flow when  $R(y) = (1 + |y|^2)^{-(d+1)/2}$  and  $u(0, \cdot)$  is convex.

We give the definition of the viscosity solution to (1.5) for the reader's convenience.

#### **Definition 3 (Viscosity solution).** (see [15] and also [6]).

(i). We say that  $u \in USC((0,\infty) \times \mathbf{R}^d)$  is a viscosity subsolution of (1.5) if the following holds: whenever  $\varphi \in C^2((0,\infty) \times \mathbf{R}^d)$  and  $u - \varphi$  attains its maximum at  $(t_o, x_o) \in (0,\infty) \times \mathbf{R}^d$ ,

$$\partial \varphi(t_o, x_o) / \partial t \le \chi(u, D\varphi(t_o, x_o), t_o, x_o) \text{Det}_+(D^2 \varphi(t_o, x_o)) R(D\varphi(t_o, x_o)).$$

(ii) We say that  $u \in LSC((0,\infty) \times \mathbf{R}^d)$  is a viscosity supersolution of (1.5) if the following holds: whenever  $\varphi \in C^2((0,\infty) \times \mathbf{R}^d)$  and  $u - \varphi$  attains its minimum at  $(t_o, x_o) \in (0,\infty) \times \mathbf{R}^d$ ,

$$\partial \varphi(t_o, x_o) / \partial t \ge \chi^-(u, D\varphi(t_o, x_o), t_o, x_o) \operatorname{Det}_+(D^2 \varphi(t_o, x_o)) R(D\varphi(t_o, x_o)).$$

Here  $\chi^{-}(v, p, t, x) = 1$  if  $p \in \partial v(t, x)$  and is not singular and if there exists  $\varepsilon > 0$  such that for all  $(s, y) \in (0, \infty) \times \mathbf{R}^{d}$  satisfying  $|y| > \varepsilon^{-1}$  and  $|s-t| < \varepsilon$ ,

$$v(s,y) > < p, y > +\varepsilon |y|,$$

and  $\chi^{-}(v, p, t, x) = 0$ , otherwise.

(iii) We say that a function  $u \in C((0, \infty) \times \mathbf{R}^d)$  is a viscosity solution of (1.5) if it is both a viscosity subsolution and supersolution of (1.5).

Theorem 4 in Sect. 2 shows that a continuous viscosity solution to (1.5) in the space of continuous functions  $v : [0, \infty) \times \mathbf{R}^d \mapsto \mathbf{R}$  for which

$$\sup\{|v(t,x) - v(0,x)| : (t,x) \in [0,T] \times \mathbf{R}^d\} < \infty \text{ for all } T > 0$$
(1.6)

is unique and is also a solution to (1.4).

Theorem 5 in Sect. 2 shows that a continuous viscosity solution, to (1.5) with  $R(y) = (1 + |y|^2)^{-(d+1)/2}$ , is unique in  $C([0,\infty) \times \mathbf{R}^d)$  under the stronger assumption on the initial value than that in Theorem 4.

In Sect. 2 we give our main result which will be proved in Sect. 4. In Sect. 3 we state and prove technical lemmas. Sect. 5 is the appendix.

We give the following notation. For any metric space A and B, the topology of C(A : B) is induced by the uniform convergence on every compact subsets of A, and for f and  $g \in C(\mathbf{R}^d)$ , we put

$$d_{C(\mathbf{R}^d)}(f,g) \equiv \sum_{m \ge 1} 2^{-m} \min(\sup_{|x| \le m} |f(x) - g(x)|, 1).$$

#### 2. Main Result

We fix a sequence  $\{\varepsilon_n\}_{n\geq 1}$  of positive real numbers which converge to zero as  $n \to \infty$ , and introduce assumptions.

(A.1).  $R \in L^1(\mathbf{R}^d : [0, \infty), dx), ||R||_{L^1} \equiv \int_{\mathbf{R}^d} R(y) dy > 0 \text{ and } h \in C(\mathbf{R}^d : \mathbf{R}).$ 

(A.2). The set  $\partial h(\mathbf{R}^d) \equiv \bigcup_{x \in \mathbf{R}^d} \partial h(x)$  has a positive Lebesgue measure, i.e.,

 $\partial h(\mathbf{R}^d)$  has a non-empty interior.

Under (A.1)-(A.2), for any  $n \ge 1$  and  $v : \mathbf{Z}^d/n \mapsto \mathbf{R}$ , put

$$\mu_{n,v}(d\beta) \equiv n^d \varepsilon_n^{-1} \{ \sum_{z \in \mathbf{Z}^d/n} w(R, \hat{v}, \{z\}) \delta_{v_{n,z}}(d\beta)$$

$$+ (||R||_{L^1} - w(R, \hat{v}, \mathbf{R}^d)) \delta_v(d\beta) \},$$
(2.1)

where  $\mathbf{Z}^d/n := \{z/n | z \in \mathbf{Z}^d\}$ , and

$$v_{n,z}(x) \equiv \begin{cases} v(x) + \varepsilon_n & \text{if } x = z, \\ v(x) & \text{if } x \in (\mathbf{Z}^d/n) \setminus \{z\}. \end{cases}$$

Notice that  $w(R, \hat{v}, \mathbf{R}^d \setminus (\mathbf{Z}^d/n)) = 0$  (see Remark 1, (i) in Sect. 1).

Put also

$$S_n \equiv \{ v : \mathbf{Z}^d / n \mapsto \mathbf{R} | \sum_{z \in \mathbf{Z}^d / n} (v(z) - h(z)) < \infty,$$

$$(v(z) - h(z)) / \varepsilon_n \in \mathbf{N} \cup \{0\} \text{ for all } z \in \mathbf{Z}^d / n \}.$$
(2.2)

Then  $S_n$  is countable and complete by the following metric: for u and  $v \in S_n$ ,

$$d_{S_n}(u,v) \equiv \sum_{z \in \mathbf{Z}^d/n} |u(z) - v(z)|.$$

For a bounded function  $f: S_n \mapsto \mathbf{R}$  and  $v \in S_n$ , put

$$A_n f(v) \equiv \int_{S_n} (f(\beta) - f(v)) \mu_{n,v}(d\beta).$$
(2.3)

Then  $A_n$  generates jump-type Markov processes on  $S_n$  under (A.1)-(A.2) (see e.g. [7, p. 162]).

Let  $\{Z_n(t, \cdot)\}_{0 \le t}$  denote a Markov process on  $S_n$ , with a generator  $A_n$ and with an initial condition  $Z_n(0, z) = h(z)$   $(z \in \mathbf{Z}^d/n)$ .

For  $t \ge 0$  and  $x \in \mathbf{R}^d$ , put

$$X_n(t,x) \equiv \max(\hat{Z}_n(t,x), h(x)).$$
(2.4)

We introduce additional assumptions and we state our first result. (A.3). The closure of the set  $\{x \in \mathbf{R}^d : \hat{h}(x) < h(x)\}$  does not contain any line which is unbounded in two different directions.

(A.4). For any  $p \notin \partial h(\mathbf{R}^d)$  and  $C \in \mathbf{R}$ ,

$$\int_{\mathbf{R}^d} \max(\langle p, x \rangle + C - h(x), 0) dx = \infty.$$
(2.5)

**Theorem 1.** Suppose that (A.1) and (A.3)-(A.4) hold. Then there exists a unique continuous solution u to (1.4) with  $u(0, \cdot) = h$ . Suppose in addition that (A.2) holds. Then the following holds: for any  $\gamma > 0$  and T > 0,

$$\lim_{n \to \infty} P(\sup_{0 \le t \le T} d_{C(\mathbf{R}^d)}(X_n(t, \cdot), u(t, \cdot)) \ge \gamma) = 0.$$
(2.6)

**Remark 2.** (i) (A.3) holds if h is convex. (ii) For any  $(p, C) \in (\mathbf{R}^d \setminus \partial h(\mathbf{R}^d)) \times \mathbf{R}$ , the set  $\{x \in \mathbf{R}^d : \langle p, x \rangle + C \rangle h(x)\}$  is unbounded.

Besides, if h is convex, then (A.4) holds. Indeed, the set  $\{x \in \mathbf{R}^d : < p, x > +C > h(x)\}$  is an open unbounded convex set.

The following theorem collects some of elementary properties of solutions to (1.4) and hence those of the motion by *R*-curvature.

**Theorem 2.** Suppose that (A.1) holds. Let  $u \in C([0, \infty) \times \mathbf{R}^d)$  be a solution of (1.4) with  $u(0, \cdot) = h$ . Then:

- (a) For each  $x \in \mathbf{R}^d$ , the function  $t \mapsto u(t, x)$  is nondecreasing in  $[0, \infty)$ .
- (b) If  $\hat{u}(t,x) < u(t,x)$  for some  $(t,x) \in (0,\infty) \times \mathbf{R}^d$ , then u(s,x) = h(x) for
- all  $s \in [0, t]$ . In particular,  $u = \max(\hat{u}, h)$  on  $[0, \infty) \times \mathbf{R}^d$ .
- (c) For any  $(t, x) \in [0, \infty) \times \mathbf{R}^d$ ,

$$u(t,x) - \hat{u}(t,x) \le h(x) - \hat{h}(x).$$
(2.7)

In particular, if  $h(x) = \hat{h}(x)$  for some  $x \in \mathbf{R}^d$ , then  $u(t, x) = \hat{u}(t, x)$ . Or equivalently, if  $\partial h(x) \neq \emptyset$  for some  $x \in \mathbf{R}^d$ , then  $\partial u(t, x) \neq \emptyset$ .

Suppose in addition that (A.4) holds. Then:

(d) For any t > 0,  $\partial u(t, \mathbf{R}^d) = \partial h(\mathbf{R}^d)$ . In particular,

$$\int_{\mathbf{R}^d} (u(t,x) - h(x))dx = t \cdot w(R,h,\mathbf{R}^d).$$
(2.8)

(e) Let  $\overline{u} \in C([0,\infty) \times \mathbf{R}^d)$  be the solution of (1.4) with  $\overline{u}(0,\cdot) = \hat{h}$ . If  $u(s,\cdot) - \hat{u}(s,\cdot) \neq h - \hat{h}$  for some  $s \in (0,\infty)$ , then  $\overline{u}(t,\cdot) - \hat{u}(t,\cdot) \neq 0$  for all  $t \geq s$ .

According to the above theorem, (a) any graph moves upward by Rcurvature, (b) those points on any graph moving by R-curvature do not move as far as they stay in its cavities, (c) the height between any graph moving by R-curvature and its convex envelope is nonincreasing as it evolves, (d) any graph moving by R-curvature sweeps in time t > 0 a region with volume given by  $t \cdot w(R, h, \mathbf{R}^d)$ , and (e) for the motion of a graph by Rcurvature, taking its convex envelope at time t > 0 and evolving up to time t starting with the convex envelope of the initial graph give different graphs in general, if the initial graph is not convex.

We introduce

(A.5). 
$$R \in C(\mathbf{R}^d : [0, \infty)),$$

and give the relation between the motion by R-curvature and the viscosity solution of (1.5).

**Theorem 3.** Suppose that (A.1) and (A.5) hold. Then a continuous solution u to (1.4) with  $u(0, \cdot) = h$  is a viscosity solution to (1.5).

We introduce more assumptions to show that the viscosity solution to (1.5) in the flamework of [15] is the motion by R-curvature.

(A.6).  $R(x) \ge R(rx)$  for any  $r \ge 1$  and  $x \in \mathbf{R}^d$ .

(A.7). 
$$\inf_{x \neq o} h(x)/|x| > 0$$

(A.8). There exists a constant C > 0 such that  $h(x+y)+h(x-y)-2h(x) \le C$ for all  $(x,y) \in \mathbf{R}^d \times U_1(o)$ , where  $U_1(o) \equiv \{y \in \mathbf{R}^d : |y| < 1\}$ .

**Theorem 4.** Suppose that (A.1) and (A.3)-(A.8) hold. Then there exists a unique continuous viscosity solution u to (1.5) with  $u(0, \cdot) = h$  in the space

of continuous functions v for which (1.6) holds. u is also a unique continuous solution to (1.4) with  $u(0, \cdot) = h$ .

We give the uniqueness result for the viscosity solution to (1.5) in a different flamework from that of [15], when the solution is a Gauss curvature flow. Put

(A.1)'.  $R(y) = (1 + |y|^2)^{-(d+1)/2}$  and  $h \in C(\mathbf{R}^d : \mathbf{R})$ .

For r > 0, define  $h^r : \mathbf{R}^d \mapsto \mathbf{R}$  by

$$h^{r}(x) = \inf\{y \in \mathbf{R} \mid U_{r}((x,y)) \subset \operatorname{epi}(h)\} \quad (x \in \mathbf{R}^{d}),$$
(2.9)

and we introduce

(A.2)'.

$$\inf_{r>0} \limsup_{\theta \downarrow 1} \sup_{\delta > 0} \liminf_{|x| \to \infty} \left( h(\theta x) - \delta |x| - h^r(x) \right) > 0,$$

$$\lim_{\theta \downarrow 1} \{ \sup_{x \in \mathbf{R}^d} (h(x) - h(\theta x)) \} = 0.$$

Then we have

**Theorem 5.** Suppose that (A.1)'-(A.2)' hold. Then for any viscosity subsolution u and supersolution v, of (1.5) in the space  $C([0, \infty) \times \mathbf{R}^d)$ , such that  $u(0, \cdot) \leq h \leq v(0, \cdot), u \leq v$ .

**Remark 3.** (A.2)' holds if there exists a convex function  $h_0 : \mathbf{R}^d \to \mathbf{R}$ such that  $h_0(x) \to \infty$  as  $|x| \to \infty$  and that

$$\lim_{|x| \to \infty} [h(x) - h_0(x)] = 0.$$
(2.10)

(see Sect. 5 for the proof).

## 3. Lemmas

In this section we state and prove technical lemmas.

**Lemma 1.** Suppose that  $f: U_{2m}(o) \mapsto [0, \infty)$  is convex for some  $m \ge 1$ . Then

$$\sup_{|x| < m} f(x) \le |U_m(o)|^{-1} \int_{|x| < 2m} f(x) dx.$$
(3.1)

(Proof). For x and  $y \in U_m(o)$ , by the convexity of f,

$$2f(x) \le f(x+y) + f(x-y).$$

Integrating the both sides on the set  $U_m(o)$  with respect to dy, the proof is over since f is nonnegative.

Q. E. D.

**Lemma 2.** Suppose that (A.1)-(A.2) hold. Then for any  $r \in (0, 1/2)$ ,  $n \ge d^{1/2}/r$ ,  $m \ge 1$  and s and t for which  $0 \le s \le t$ , the following holds almost surely:

$$\sup_{|x| \le m} (X_n(t, x) - X_n(s, x))$$
(3.2)

Motion of a graph by R-curvature

$$\leq 2\{4r \sup_{|x| \leq m+2, u=s,t} |\hat{Z}_n(u,x)| + \sup_{|x|, |y| \leq m+1, |x-y| < 2r} |h(x) - h(y)|\} + \sum_{z \in \mathbf{Z}^d/n, |z| \leq m+1} (Z_n(t,z) - Z_n(s,z))n^{-d}/|U_r(o)| + \varepsilon_n.$$

(Proof). For  $x = (x_i)_{i=1}^d \in \mathbf{R}^d$ , put  $[x] \equiv ([x_i])_{i=1}^d$ , where  $[x_i]$  is an integer such that  $[x_i] \leq x_i < [x_i] + 1$ . Then for any  $t \geq 0$  and  $x \in \mathbf{R}^d$ ,

$$0 \le Z_n(t, [nx]/n) - X_n(t, [nx]/n) < \varepsilon_n.$$
(3.3)

This is true, since

$$Z_n(t, [nx]/n) \begin{cases} \in [\hat{Z}_n(t, [nx]/n), \hat{Z}_n(t, [nx]/n) + \varepsilon_n) \\ & \text{if } \hat{Z}_n(t, [nx]/n) \ge h([nx]/n), \\ & = h([nx]/n) \quad \text{if } \hat{Z}_n(t, [nx]/n) < h([nx]/n). \end{cases}$$

For  $x \in U_m(o)$ ,

$$0 \leq X_{n}(t,x) - X_{n}(s,x)$$

$$\leq \sup_{|x-y| \leq r} \{ |X_{n}(t,x) - X_{n}(t,[ny]/n)| + |X_{n}(s,[ny]/n) - X_{n}(s,x)| \}$$

$$+ \int_{U_{r}(x)} (X_{n}(t,[ny]/n) - X_{n}(s,[ny]/n)) dy / |U_{r}(o)|.$$
(3.4)

Hence (3.2) holds by (2.4) and (3.3).

Indeed, for  $t \ge 0$ ,  $x \in U_m(o)$  and y for which  $|x - y| \le r$ , if  $X_n(t, x) = h(x)$  and  $X_n(t, [ny]/n) = h([ny]/n)$ , then

$$|X_n(t,x) - X_n(t, [ny]/n)| \le \sup_{|x|, |\tilde{y}| \le m+1, |x-\tilde{y}| < 2r} |h(x) - h(\tilde{y})|.$$

Hitoshi Ishii, Toshio Mikami

If  $X_n(t,x) = \hat{Z}_n(t,x)$  and  $X_n(t, [ny]/n) = \hat{Z}(t, [ny]/n)$ , then

$$|X_n(t,x) - X_n(t, [ny]/n)| \le 4r \sup_{|\overline{y}| \le m+2, u=s, t} |\hat{Z}_n(u, \overline{y})|$$

by the following: for a convex function  $f : \mathbf{R}^d \mapsto \mathbf{R}$  and r > 0,

$$\sup\{|f(x) - f(y)| / |x - y| : x \neq y, x, y \in U_r(o)\} \le 2 \sup_{|z| \le r+1} |f(z)|$$
(3.5)

(see e.g. [2, p. 20, Lemma 3.1]). If  $X_n(t, x) = \hat{Z}_n(t, x)$  and  $X_n(t, [ny]/n) = h([ny]/n)$ , then one can take a point  $\overline{\overline{y}}$  between x and [ny]/n so that  $\hat{Z}_n(t, \overline{\overline{y}}) = h(\overline{\overline{y}})$  by the intermediate value theorem, and

$$\begin{aligned} |X_n(t,x) - X_n(t, [ny]/n)| \\ &\leq |\hat{Z}_n(t,x) - \hat{Z}_n(t,\overline{y})| + |h(\overline{y}) - h([ny]/n)| \\ &\leq 4r \sup_{|\overline{y}| \leq m+2, u=s, t} |\hat{Z}_n(u,\overline{y})| + \sup_{|x|, |\tilde{y}| \leq m+1, |x-\tilde{y}| < 2r} |h(x) - h(\tilde{y})|. \end{aligned}$$
Q. E. D.

**Lemma 3.** Suppose that (A.1)-(A.2) hold. Then for any T > 0,  $n \ge 2d^{1/2}$ and  $m \ge 1$ ,

$$P(\sup_{0 \le t \le T} \int_{|x| \le m} \{\hat{Z}_n(t, x) + 2 \sup_{|y| \le m+2} |\hat{h}(y)|\} dx$$

$$> 2^{d+1} m(\sup_{|x| \le m+2} \{2|\hat{h}(x)| + |h(x)|\} |U_{m+2}(o)| + T||R||_{L^1}))$$

$$\le \varepsilon_n n^{-d} m^{-2} (\sup_{|x| \le m+2} \{2|\hat{h}(x)| + |h(x)|\} |U_{m+2}(o)| + T||R||_{L^1})^{-1}.$$
(3.6)

(Proof). For  $t \in [0, T]$ ,

$$\int_{|x| \le m} \{ \hat{Z}_{n}(t,x) + 2 \sup_{|y| \le m+2} |\hat{h}(y)| \} dx$$

$$\leq 2^{d} \sum_{z \in \mathbf{Z}^{d}/n, |z| \le m+1} \{ Z_{n}(t,z) + 2 \sup_{|y| \le m+2} |\hat{h}(y)| \} n^{-d}$$

$$\leq 2^{d} \{ \sum_{z \in \mathbf{Z}^{d}/n, |z| \le m+1} (Z_{n}(t,z) - Z_{n}(0,z)) n^{-d} - \int_{0}^{t} ds \sum_{z \in \mathbf{Z}^{d}/n, |z| \le m+1} w(R, \hat{Z}_{n}(s, \cdot), \{z\}) \}$$

$$+ 2^{d} (\sup_{|x| \le m+2} \{ 2|\hat{h}(x)| + |h(x)| \} |U_{m+2}(o)| + T||R||_{L^{1}}),$$

$$(3.7)$$

since  $w(R, \hat{Z}_n(s, \cdot), U_{m+2}(o)) \leq ||R||_{L^1}$ . Here we used the fact that a convex function takes its maximum on the boundary of the set where it is defined and that  $Z_n(t, z) + |\hat{h}(z)| \geq 0$ .

Hence by Doob-Kolmogorov's inequality (see [13, p. 34]), the following completes the proof: by Itô's formula (see [7, p. 162] or [13, p. 66]), for any s and t for which  $0 \le s \le t$  and any  $\varphi \in C_b(\mathbf{R}^d)$ ,

$$E[|\sum_{z \in \mathbf{Z}^d/n} \varphi(z)(Z_n(t,z) - Z_n(s,z))n^{-d}$$
(3.8)  
$$-\int_s^t \sum_{z \in \mathbf{Z}^d/n} \varphi(z)w(R, \hat{Z}_n(u, \cdot), \{z\})du|^2]$$
$$= E[\int_s^t \sum_{z \in \mathbf{Z}^d/n} \varphi(z)^2 w(R, \hat{Z}_n(u, \cdot), \{z\})du]\varepsilon_n n^{-d}.$$
Q. E. D.

**Lemma 4.** Suppose that (A.1)-(A.2) hold. Then for any  $\eta$  and T > 0, the following holds: for sufficiently small  $\delta > 0$ ,

$$\lim_{n \to \infty} P(\max_{1 \le i \le [T/\delta]+1} d_{C(\mathbf{R}^d)}(X_n(i\delta, \cdot), X_n((i-1)\delta, \cdot)) \ge \eta) = 0.$$
(3.9)

(Proof). Take m for which  $2^{-m} < \eta/2,$  and take also  $r \in (0,1/2)$  such that

$$r \leq (\eta/64)|U_{m+2}(o)|\{2^{d+2}(m+2)(\sup_{|x|\leq 2(m+2)+2}\{2|\hat{h}(x)| + |h(x)|\}|U_{2(m+2)+2}(o)| + (T+1)||R||_{L^1})\}^{-1},$$

$$2 \sup_{|x|,|y| \le m+1, |x-y| < 2r} |h(x) - h(y)| < \eta/8.$$

Then for any  $\delta \in (0, \min(1, \eta |U_r(o)|/(8||R||_{L^1})))$  and any  $n \ge d^{1/2}/r$ , by Lemmas 1-2,

$$P(\max_{1 \le i \le [T/\delta]+1} d_{C(\mathbf{R}^d)}(X_n(i\delta, \cdot), X_n((i-1)\delta, \cdot)) \ge \eta)$$
(3.10)  
$$\le P(\sup_{0 \le t \le T+1} \int_{|x| \le 2(m+2)} \{\hat{Z}_n(t, x) + 2 \sup_{|y| \le 2(m+3)} |\hat{h}(y)|\} dx > 2^{d+2}(m+2)$$
$$\times (\sup_{|x| \le 2(m+3)} \{2|\hat{h}(x)| + |h(x)|\} |U_{2(m+3)}(o)| + (T+1)||R||_{L^1}))$$
$$+ \sum_{1 \le i \le [T/\delta]+1} P(|\sum_{z \in \mathbf{Z}^d/n, |z| \le m+1} (Z_n(i\delta, z) - Z_n((i-1)\delta, z))n^{-d}$$
$$- \int_{(i-1)\delta}^{i\delta} \sum_{z \in \mathbf{Z}^d/n, |z| \le m+1} w(R, \hat{Z}_n(s, \cdot), \{z\}) ds| \ge |U_r(o)|(\eta/8 - \varepsilon_n)).$$

Indeed, from Lemma 1, for any  $t \ge 0$ ,

$$\sup_{|x| \le m+2} |\hat{Z}_n(t,x)| \le \sup_{|x| \le m+2} \{\hat{Z}_n(t,x) + 2 \sup_{|y| \le 2(m+3)} |\hat{h}(y)|\}$$
(3.11)

16

Motion of a graph by R-curvature

$$\leq \frac{1}{|U_{m+2}(o)|} \int_{|x| \leq 2(m+2)} \{\hat{Z}_n(t,x) + 2 \sup_{|y| \leq 2(m+3)} |\hat{h}(y)|\} dx,$$

since

$$\inf_{|x| \le 2(m+2)} \{ \hat{Z}_n(t,x) + 2 \sup_{|y| \le 2(m+3)} |\hat{h}(y)| \} \ge 0.$$
(3.12)

(3.10) together with Lemma 3 and (3.8) completes the proof.

Q. E. D

**Lemma 5.** Suppose that (A.1)-(A.2) hold. Then for any  $n \ge 1$ , the following holds almost surely: for any  $t \ge 0$  and  $x \in \mathbf{R}^d$ ,

$$X_n(t,x) = \max(\hat{X}_n(t,x), h(x)).$$
(3.13)

(Proof). From (2.4) and [16, p. 56, Prop. 2.31],

$$\hat{Z}_{n}(t,x) \leq \hat{X}_{n}(t,x)$$

$$= \inf\{\sum_{i=0}^{d} \lambda_{i} X_{n}(t,x_{i}) : \lambda_{i} \geq 0, x_{i} \in \mathbf{R}^{d} (i=0,\cdots,d)$$

$$, \sum_{i=0}^{d} \lambda_{i} = 1, \sum_{i=0}^{d} \lambda_{i} x_{i} = x\}$$

$$\leq \inf\{\sum_{i=0}^{d} \lambda_{i} Z_{n}(t,x_{i}) : \lambda_{i} \geq 0, x_{i} \in \mathbf{Z}^{d} / n (i=0,\cdots,d)$$

$$, \sum_{i=0}^{d} \lambda_{i} = 1, \sum_{i=0}^{d} \lambda_{i} x_{i} = x\} = \hat{Z}_{n}(t,x).$$
Q. E. D.

**Lemma 6.** Suppose that (A.1) and (A.3)-(A.4) hold. Take functions f and  $f_n \in C(\mathbf{R}^d)$   $(n \ge 1)$  such that  $f_n = \max(\hat{f}_n, h)$   $(n \ge 1)$ , that  $f_n \to f$  in

 $C(\mathbf{R}^d)$  as  $n \to \infty$  and that  $\int_{\mathbf{R}^d} (f(x) - h(x)) dx$  is finite. Then  $\partial f(\mathbf{R}^d) = \partial h(\mathbf{R}^d)$  and  $\hat{f}_n \to \hat{f}$  in  $C(\mathbf{R}^d)$  as  $n \to \infty$ . In particular,  $f = \max(\hat{f}, h)$  and the following holds: for any  $\varphi \in C_o(\mathbf{R}^d)$ ,

$$\lim_{n \to \infty} \int_{\mathbf{R}^d} \varphi(x) w(R, f_n, dx) = \int_{\mathbf{R}^d} \varphi(x) w(R, f, dx).$$
(3.15)

(Proof). We first show that  $\partial f(\mathbf{R}^d) = \partial h(\mathbf{R}^d)$ .

 $\partial h(\mathbf{R}^d) \subset \partial f(\mathbf{R}^d)$  since  $f \ge h$ .

 $\partial f(\mathbf{R}^d) \subset \partial h(\mathbf{R}^d)$  by (A.4) since for any  $x_0 \in \mathbf{R}^d$  for which  $\partial f(x_0) \neq \emptyset$ and any  $p \in \partial f(x_0)$ , the following implies that  $p \in \partial h(x_0)$ :

$$\int_{\mathbf{R}^d} \max(\langle p, x - x_0 \rangle + f(x_0) - h(x), 0) dx \le \int_{\mathbf{R}^d} (f(x) - h(x)) dx < \infty.$$

Since  $f_n \ge \hat{f}_n \ge \hat{h}$  and  $\{f_n\}_{n\ge 1}$  is uniformly bounded on every compact subset of  $\mathbf{R}^d$ , there exists a convergent subsequence  $\{\hat{f}_{n_k}\}_{k\ge 1}$  in  $C(\mathbf{R}^d)$  (see [2, p. 21, Theorem 3.2]). We denote by g the limit of  $\hat{f}_{n_k}$  as  $k \to \infty$ . It is easy to see that  $f = \max(g, h)$  and that  $f \ge \hat{f} \ge g \ge \hat{h}$ .

Suppose that  $\hat{f} \neq g$ . Then there exist  $x_1$  and  $x_2 \in \mathbf{R}^d$  and  $p \in \partial \hat{f}(x_1) \cap \partial g(x_2)$  such that

$$\langle p, x - x_1 \rangle + \hat{f}(x_1) \rangle \langle p, x - x_2 \rangle + g(x_2)$$
 for all  $x \in \mathbf{R}^d$  (3.16)

since  $\partial \hat{f}(\mathbf{R}^d) = \partial g(\mathbf{R}^d)$  from the above argument.

For sufficiently large  $k \geq 1$ , one can take  $(\lambda_{k,i}, x_{k,i})_{i=0}^d \in ([0,1] \times \mathbf{R}^d)^{d+1}$  such that  $\sum_{i=0}^d \lambda_{k,i} = 1$ ,  $\sum_{i=0}^d \lambda_{k,i} x_{k,i} = x_2$ ,  $h(\sum_{i=0}^d t_i x_{k,i}) > 0$ 

 $\hat{f}_{n_k}(\sum_{i=0}^d t_i x_{k,i})$  for all  $t_i \ge 0$   $(i = 0, \dots, d)$  for which  $\sum_{i=0}^d t_i = 1$  and for which  $t_i = 0$  if  $\lambda_{k,i} = 0$ , and that

$$0 \le \sum_{i=0}^{d} \lambda_{k,i} h(x_{k,i}) - \hat{f}_{n_k}(x_2) < 1/k$$

Indeed, since  $f(x_2) = h(x_2) \ge \hat{f}(x_2) > g(x_2)$  by (3.16),  $f_{n_k}(x_2) = h(x_2) > \hat{f}_{n_k}(x_2)$  for sufficiently large  $k \ge 1$ , and hence from [16, p. 56, Prop. 2.31],

$$\hat{f}_{n_k}(x_2) = \inf\{\sum_{i=0}^d \lambda_i f_{n_k}(y_i) : \lambda_i \ge 0, y_i \in \mathbf{R}^d (i = 0, \cdots, d)$$
(3.17)  
$$, \sum_{i=0}^d \lambda_i = 1, \sum_{i=0}^d \lambda_i y_i = x_2 \}$$
$$= \inf\{\sum_{i=0}^d \lambda_i h(y_i) : \lambda_i \ge 0, y_i \in \mathbf{R}^d (i = 0, \cdots, d)$$
$$, \sum_{i=0}^d \lambda_i = 1, \sum_{i=0}^d \lambda_i y_i = x_2, h(\sum_{i=0}^d t_i y_i) > \hat{f}_{n_k}(\sum_{i=0}^d t_i y_i)$$
$$(t_i \ge 0, \ t_i = 0 \ \text{if} \ \lambda_i = 0 (i = 0, \cdots, d), \sum_{i=0}^d t_i = 1) \}$$

by approximating the intersection of the graph of  $y = \hat{f}_{n_k}(x)$  and that of the supporting hyperplane of  $y = \hat{f}_{n_k}(x)$  at  $x = x_2$  from the inside.

In (3.17) we used the following argument. Suppose that there exist  $j \in \{1, \dots, d\}$  and  $\lambda_i > 0, y_i \in \mathbf{R}^d (i = 0, \dots, j)$  for which  $h(y_i) > \hat{f}_{n_k}(y_i)(i = 0, \dots, j), \sum_{i=0}^j \lambda_i = 1$  and  $\sum_{i=0}^j \lambda_i y_i = x_2$ . Suppose also that there exists  $t_i \ge 0 (i = 0, \dots, j)$  for which  $\sum_{i=0}^j t_i = 1, \ \tilde{y} \equiv \sum_{i=0}^j t_i y_i (\neq x_2)$  and  $\hat{f}_{n_k}(\tilde{y}) = f_{n_k}(\tilde{y})$ . Put  $\lambda \equiv \min_{i=0,\dots,j} (\lambda_i/t_i) \ (\in (0,1))$ . Then

$$\lambda + \sum_{i=0,\dots,j,\lambda_i/t_i > \lambda} (\lambda_i - \lambda t_i) = 1,$$

$$\lambda \tilde{y} + \sum_{i=0,\cdots,j,\lambda_i/t_i > \lambda} (\lambda_i - \lambda t_i) y_i = x_2$$

$$\lambda h(\tilde{y}) + \sum_{i=0,\dots,j,\lambda_i/t_i > \lambda} (\lambda_i - \lambda t_i) h(y_i) < \sum_{i=0}^j \lambda_i h(y_i)$$

by the convexity of  $\hat{f}_{n_k}$ , since

$$h(\tilde{y}) \le \hat{f}_{n_k}(\tilde{y}) = f_{n_k}(\tilde{y}), \quad \hat{f}_{n_k}(y_i) < h(y_i) = f(y_i) \quad (i = 0, \cdots, j).$$

Here one can assume, without loss of generality, that  $(\tilde{y}, \hat{f}_{n_k}(\tilde{y}))$  is on the supporting hyperplane of  $y = \hat{f}_{n_k}(x)$  at  $x = x_2$ . Therefore one can find a point  $\overline{y}$  between  $x_2$  and  $\tilde{y}$  such that  $h(\overline{y}) > \hat{f}_{n_k}(\overline{y})$  and that there exists positive numbers  $\lambda, \overline{\lambda}_i$   $(i = 0, \dots, j, \lambda_i/t_i > \lambda)$  for which the following holds:

$$\overline{\lambda} + \sum_{i=0,\cdots,j,\lambda_i/t_i > \lambda} \overline{\lambda}_i = 1,$$

$$\overline{\lambda}\overline{y} + \sum_{i=0,\cdots,j,\lambda_i/t_i > \lambda} \overline{\lambda}_i y_i = x_2,$$

$$\overline{\lambda}h(\overline{y}) + \sum_{i=0,\cdots,j,\lambda_i/t_i > \lambda} \overline{\lambda}_i h(y_i) < \sum_{i=0}^j \lambda_i h(y_i).$$

Since  $\hat{f}(x_2) > g(x_2)$ ,  $\{x_{k,i} : \lambda_{k,i} > 0, i = 0, \dots, d\}_{k \ge 1}$  is not bounded. Therefore, by (A.3), there exists a sequence  $\{m(k)\}_{k \ge 1} \subset \mathbf{N}$  such that  ${x_{m(k),i}, \lambda_{m(k),i}^{-1}}_{k\geq 1}$  is bounded and that  $|\sum_{j\neq i,j=0,\cdots,d,} \lambda_{m(k),j}x_{m(k),j}| \rightarrow \infty$  as  $k \rightarrow \infty$ , for some  $i \in \{0,\cdots,d\}$ . Hence  $\operatorname{epi}(g)$  contains a line which is infinite in one direction, which intersects with  $\operatorname{epi}(\hat{f})$  in another one and which contains  $(x_2, g(x_2))$ . This contradicts (3.16).

(3.15) can be proved by [2, p. 119, Th. 9.1].

Q. E. D.

**Lemma 7.** Suppose that  $R \in L^1(\mathbf{R}^d : [0, \infty), dx)$ . Take continuous solutions  $u_1$  and  $u_2$ , to (1.4), for which  $u_1(0, \cdot) \leq u_2(0, \cdot)$  and for which  $\partial u_1(t, \mathbf{R}^d) \subset \partial u_2(t, \mathbf{R}^d)$  for all  $t \geq 0$ . Then  $u_1 \leq u_2$ .

(Proof). Suppose that  $u_1 \leq u_2$  is not true. For  $\varepsilon > 0$ , put

$$V_{\varepsilon} = \{(t, x) \in (0, \varepsilon^{-1}) \times \mathbf{R}^d; u_2(t, x) + \varepsilon t < u_1(t, x)\}.$$

Then for sufficiently small  $\varepsilon > 0$ , the set  $V_{\varepsilon}$  is open and the volume  $|V_{\varepsilon}| > 0$ .

If  $(\tau,\xi) \in V_{\varepsilon}$  and  $p \in \partial u_1(\tau,\xi)$ , then  $p \in \partial u_2(\tau,\eta)$  for some  $\eta$ .

Besides,  $(\tau, \eta) \in V_{\varepsilon}$ . Indeed, put

$$l(x) := u_1(\tau, \xi) + \langle p, x - \xi \rangle \qquad \text{for all } x \in \mathbf{R}^d.$$

Then  $u_2(\tau, x) - l(x)$  attains the minimum over  $\mathbf{R}^d$  at a point  $\eta$ . If  $(\tau, x) \notin V_{\varepsilon}$ , then

$$u_2(\tau, x) \ge u_1(\tau, x) - \varepsilon \tau \ge l(x) - \varepsilon \tau, \qquad (3.18)$$

and

$$u_{2}(\tau,\eta) - l(\eta) - \varepsilon\tau$$

$$\leq u_{2}(\tau,\xi) - l(\xi) - \varepsilon\tau < u_{1}(\tau,\xi) - l(\xi) - 2\varepsilon\tau = -2\varepsilon\tau.$$
(3.19)

An immediate consequence is that

$$\int_{V_{\varepsilon}} w(R, u_1(t, \cdot), dx) dt \le \int_{V_{\varepsilon}} w(R, u_2(t, \cdot), dx) dt.$$
(3.20)

Take a nondecreasing sequence  $\{\eta_n\}_{n\geq 1}$  of nondecreasing  $C^1\text{-functions}$  such that

$$\eta_n(r) = 0$$
 for all  $r \le 0$ ,  $\eta_n(r) = 1$  for all  $r \ge \frac{1}{n}$ , (3.21)

and for  $r \in \mathbf{R}$ , put

$$\zeta_n(r) = \int_0^r \eta_n(s) ds. \tag{3.22}$$

Then for any  $x \in \mathbf{R}^d$  and  $t \in (0, \varepsilon^{-1})$ ,

$$0 \le \zeta_n(u_1(t,x) - u_2(t,x) - \varepsilon t)$$

$$= \int_0^t \eta_n(u_1(s,x) - u_2(s,x) - \varepsilon s)(u_1(ds,x) - u_2(ds,x) - \varepsilon ds).$$
(3.23)

(Notice that the function  $t \mapsto u_i(t, x)$  is nondecreasing for i = 1, 2 and  $x \in \mathbf{R}^d$ .) Hence by (1.4), for any r > 0

$$0 \leq \int_{|x| \leq r} \int_{0}^{\varepsilon^{-1}} \eta_{n}(u_{1}(s,x) - u_{2}(s,x) - \varepsilon s)$$

$$\times (u_{1}(ds,x) - u_{2}(ds,x) - \varepsilon ds) dx$$

$$= \int_{0}^{\varepsilon^{-1}} \int_{|x| \leq r} \eta_{n}(u_{1}(s,x) - u_{2}(s,x) - \varepsilon s)$$

$$\times (w(R, u_{1}(s, \cdot), dx) - w(R, u_{2}(s, \cdot), dx) - \varepsilon dx) ds$$

$$\rightarrow \int_{V_{\varepsilon}} w(R, u_{1}(t, \cdot), dx) dt - \int_{V_{\varepsilon}} w(R, u_{2}(t, \cdot), dx) dt - \varepsilon |V_{\varepsilon}|$$
(3.24)

as  $r \to \infty$  and then  $n \to \infty$ . This together with (3.20) implies that  $|V_{\varepsilon}| = 0$ , which is a contradiction.

Q. E. D.

**Lemma 8.** Suppose that  $u \in C((0,\infty) \times \mathbf{R}^d)$  and  $\psi \in C^2((0,\infty) \times \mathbf{R}^d)$ satisfy the following: for some  $(s, y) \in (0, \infty) \times \mathbf{R}^d$ ,

$$\chi^{-}(u, D\psi(s, y), s, y) = 1, \quad u(s, y) = \psi(s, y), \quad D^{2}\psi(s, y) > 0,$$

and there exists A > 0 such that for any  $\varepsilon > 0$ ,

$$U_{\varepsilon}^{-} \equiv \{(t,x) \in (0,\infty) \times \mathbf{R}^{d} | \psi(t,x) + \varepsilon > u(t,x)\} \subset U_{(\varepsilon/A)^{1/2}}((s,y)).$$
(3.25)

Then for sufficiently small  $\varepsilon > 0$  and any  $(\tau, \xi) \in U_{\varepsilon}^{-}$ ,  $D\psi(\tau, \xi) \in \partial u(\tau, z)$ for some z for which  $(\tau, z) \in U_{\varepsilon}^{-}$ .

(Proof). Take  $r \in (0, s)$  such that

$$D^2\psi(t,x) > 0$$
 for all  $(t,x) \in [s-r,s+r] \times U_r(y)$ . (3.26)

Since  $\chi^{-}(u, D\psi(s, y), s, y) = 1$ , by the continuity of u, there exists a constant  $\delta \in (0, r]$  such that for all  $p \in U_{\delta}(D\psi(s, y))$ , and  $(t, x) \in [s - \delta, s + \delta] \times (\mathbf{R}^{d} \setminus U_{r}(y))$ ,

$$u(t,x) \ge \psi(s,y) + \delta + \langle p, x - y \rangle \tag{3.27}$$

(see Definition 3). Take a constant  $\gamma \in (0, \delta]$  so that

$$\psi(s,y) + \delta \ge \psi(t,y) + \gamma \quad \text{for all } t \in [s - \gamma, s + \gamma],$$
(3.28)

$$D\psi(t,x) \in U_{\delta}(D\psi(s,y))$$
 for all  $(t,x) \in [s-\gamma,s+\gamma] \times U_{\gamma}(y)$ . (3.29)

Take  $\varepsilon \in (0, \gamma]$  sufficiently small so that  $U_{\varepsilon}^{-} \subset [s - \gamma, s + \gamma] \times U_{\gamma}(y)$ . Then for  $(\tau, \xi) \in U_{\varepsilon}^{-}$ , from (3.26), we see that

$$\psi(\tau, x) \ge \tilde{l}(x) := \psi(\tau, \xi) + \langle D\psi(\tau, \xi), x - \xi \rangle \quad \text{for all } x \in U_r(y).$$
(3.30)

In particular, we have from (3.28),

$$\psi(s,y) + \delta \ge \psi(\tau,y) + \varepsilon \ge \tilde{l}(y) + \varepsilon. \tag{3.31}$$

Hence for all  $x \in \mathbf{R}^d \setminus U_r(y)$ , by (3.27) and (3.29),

$$\tilde{l}(x) + \varepsilon = \tilde{l}(y) + \varepsilon + \langle D\psi(\tau,\xi), x - y \rangle$$

$$\leq \psi(s,y) + \delta + \langle D\psi(\tau,\xi), x - y \rangle \leq u(\tau,x).$$
(3.32)

We also have, by (3.30), for all  $x \in U_r(y)$  for which  $(\tau, x) \notin U_{\varepsilon}^-$ ,

$$\tilde{l}(x) + \varepsilon \le \psi(\tau, x) + \varepsilon \le u(\tau, x).$$
(3.33)

Since

$$u(\tau,\xi) - \tilde{l}(\xi) < \varepsilon,$$

the function  $x \mapsto u(\tau, x) - \tilde{l}(x)$  attains a minimum at z for which  $(\tau, z) \in U_{\varepsilon}^{-}$ , which means that  $D\psi(\tau, \xi) \in \partial u(\tau, z)$ .

Q. E. D.

The following two lemmas can be shown by the arguments in the proof of [15, Theorem 1], and we omit the proof.

**Lemma 9.** Suppose that (A.5)-(A.7) hold, and that  $u_1$  and  $u_2$  are continuous viscosity solutions, to (1.5) with  $u_1(0, \cdot) = u_2(0, \cdot)$ , for which

$$\sup\{|u_i(t,x) - u_i(0,x)| : i = 1, 2, (t,x) \in [0,T] \times \mathbf{R}^d\} < \infty \quad \text{for all } T > 0.$$
(3.34)

Then  $u_1 = u_2$ .

**Lemma 10.** Suppose that (A.1) and (A.5) hold, that v and u are a viscosity supersolution and a viscosity subsolution of (1.5) in  $(0,T) \times \mathbf{R}^d$ , respectively, where T > 0 is a constant, that v and u are lower semicontinuous and continuous on  $[0,T] \times \mathbf{R}^d$ , respectively, and that  $u(0,x) \leq v(0,x)$  for all  $x \in \mathbf{R}^d$  and

$$u(s,x) \le v(t,x) - \delta |x| + C_0$$
 for all  $s, t \in [0,T], x \in \mathbf{R}^d$  (3.35)

for some constants  $\delta > 0$  and  $C_0 > 0$ . Then  $u \leq v$  in  $[0, T) \times \mathbf{R}^d$ .

#### 4. Proof of Main Result

In this section we prove theorems given in Sect. ??.

(Proof of Theorem 1). First of all, we point out that one can show, from (A.4), that  $\partial v(t, \mathbf{R}^d) = \partial h(\mathbf{R}^d)$  for all  $t \ge 0$ , for any continuous solution v to (1.4) with  $v(0, \cdot) = h(\cdot)$ , by the argument of the first part of the proof of Lemma 6. In particular, a continuous solution v to (1.4) with  $v(0, \cdot) = h(\cdot)$  is unique by Lemma 7.

Suppose first that (A.2) does not hold. Then  $u(t, \cdot) \equiv h(\cdot)$  for all  $t \ge 0$ is a unique solution to (1.4) with  $u(0, \cdot) = h(\cdot)$ .

Suppose next that (A.2) holds. Take  $m_0 > 0$  such that

$$\sup_{|x| \le 2m_0 + 2} \{ 2|\hat{h}(x)| + |h(x)| \} > 0.$$

For  $t \ge 0$ , put

$$\Gamma_{t} \equiv \{\max(f,h)(x)|f: \mathbf{R}^{d} \mapsto \mathbf{R} \text{ is convex, and}$$

$$\sup_{|x| \le m} |f(x)| \le 2^{d+2} m |U_{m}(o)|^{-1} (\sup_{|x| \le 2m+2} \{2|\hat{h}(x)| + |h(x)|\} |U_{2m+2}(o)| + t ||R||_{L^{1}}) \text{ for all } m \ge m_{0} \}.$$
(4.1)

Then  $\Gamma_t$  is compact in  $C(\mathbf{R}^d)$  (see [2, section 3.3]).

By Lemma 3 and (3.11), the following holds: for any  $t \ge 0$ ,

$$\lim_{n \to \infty} P(X_n(t, \cdot) \in \Gamma_t) = 1.$$
(4.2)

This together with Lemma 4 implies the tightness of  $\{X_n(t,\cdot)\}_{0 \le t,n \ge 1}$  in  $D([0,\infty) : C(\mathbf{R}^d))$ , since for any  $\delta > 0$  and any t and s for which  $[t/\delta] = [s/\delta]$ ,

$$d_{C(\mathbf{R}^d)}(X_n(t,\cdot),X_n(s,\cdot)) \le d_{C(\mathbf{R}^d)}(X_n(([t/\delta]+1)\delta,\cdot),X_n([t/\delta]\delta,\cdot))$$

(see [7, p. 129, Corollary 7.4]).

One can also show, by Lemma 4, that any weak limit point of  $\{X_n(t, \cdot)\}_{t\geq 0}$ , as  $n \to \infty$ , belongs to the set  $C([0, \infty) : C(\mathbf{R}^d))$ , since for any t > 0 and  $\delta > 0$ ,

$$d_{C(\mathbf{R}^d)}(X_n(t,\cdot),X_n(t-,\cdot)) \le \max_{1\le i\le [t/\delta]+1} d_{C(\mathbf{R}^d)}(X_n(i\delta,\cdot),X_n((i-1)\delta,\cdot))$$

(see [7, p. 148, Theorem 10.2, (a)]).

Let  $\{X_{n_k}\}_{k\geq 1}$  be a weakly convergent subsequence of  $\{X_n\}_{n\geq 1}$ , and ube the weak limit of  $\{X_{n_k}\}_{k\geq 1}$ . Then by Skorohod's theorem (see [7, p. 102, Theorem 1.8]), taking a new probability space, one can assume that for any  $T \geq 0$ ,  $X_{n_k}(t, \cdot)$  converges, as  $k \to \infty$ , to  $u(t, \cdot)$  in  $C(\mathbf{R}^d)$  uniformly in  $t \in [0, T]$  a.s.. By (3.8), we have

$$E[\int_{\mathbf{R}^d} (u(t,x) - h(x))dx] \le t||R||_{L^1},$$
(4.3)

since

$$X_n(0, [nx]/n) = h([nx]/n) \to u(0, x) = h(x) \quad (\text{as } n \to \infty)$$
(4.4)

for  $x \in \mathbf{R}^d$ , and since for  $\varphi \in C_o(\mathbf{R}^d)$ 

$$|\sum_{z \in \mathbf{Z}^d/n_k} \varphi(z)(Z_{n_k}(t,z) - Z_{n_k}(0,z))n_k^{-d}$$

$$-\int_{\mathbf{R}^d} \varphi([n_k x]/n_k)(X_{n_k}(t,[n_k x]/n_k) - X_{n_k}(0,[n_k x]/n_k))dx|$$

$$\leq 2\varepsilon_{n_k} \int_{\mathbf{R}^d} |\varphi([n_k x]/n_k)|dx \to 0 \quad (\text{as } k \to \infty) \text{ a.s.}$$

$$(4.5)$$

by (3.3). Hence by (1.3), (4.3), Lemmas 5 and 6, for any  $\varphi \in C_o(\mathbf{R}^d)$  and any  $t \ge 0$ ,

$$\sum_{z \in \mathbf{Z}^d/n_k} \varphi(z) w(R, \hat{Z}_{n_k}(t, \cdot), \{z\}) = \int_{\mathbf{R}^d} \varphi(x) w(R, X_{n_k}(t, \cdot), dx) \quad (4.6)$$
$$\to \int_{\mathbf{R}^d} \varphi(x) w(R, u(t, \cdot), dx)$$

as  $k \to \infty$  a.s. (3.8) and (4.5)-(4.6) imply that u(t, x) is a unique continuous solution to (1.4) with u(0, x) = h(x). In particular, u is nonrandom and hence (2.6) holds.

Q. E. D.

(Proof of Theorem 2). The proof of (a) is standard and is omitted.

28

(b) Suppose that  $\hat{u}(t,x) < u(t,x)$ . Then there exists a constant r > 0 such that

$$\hat{u}(s,y) < u(s,y) \qquad \text{for all } (s,y) \in [t-r,t] \times U_r(x), \tag{4.7}$$

since the function  $s \mapsto \hat{u}(s, y)$  is non-decreasing in  $[0, \infty)$  for each  $y \in \mathbf{R}^d$ , and  $\hat{u}(s, \cdot) \in C(\mathbf{R}^d)$  for each  $s \in [0, \infty)$ , and  $u \in C([0, \infty) \times \mathbf{R}^d)$ . Therefore, we have

$$\int_{[t-r,t]\times U_r(x)} dsw(R, u(s, \cdot), dy) = 0.$$

$$(4.8)$$

This together with (1.4) implies the following:

$$u(s,y) = u(t,y) \qquad \text{for all } (s,y) \in [t-r,t] \times U_r(x). \tag{4.9}$$

Since the function  $s \mapsto u(s, y)$  is nondecreasing in  $[0, \infty)$  for each  $y \in \mathbf{R}^d$ , this implies that  $u(t, \cdot) = u(0, \cdot)$  in the set  $U_r(x)$ .

(c) If  $u(t,x) = \hat{u}(t,x)$ , then  $u(t,x) - \hat{u}(t,x) \le h(x) - \hat{h}(x)$  since  $h(x) \ge \hat{h}(x)$ . Suppose that  $\hat{u}(t,x) < u(t,x)$ . Then u(t,x) = h(x) by (b), from which we conclude that  $u(t,x) - \hat{u}(t,x) \le h(x) - \hat{h}(x)$  since  $\hat{u}(t,x) \ge \hat{h}(x)$ .

(d)  $\partial u(t, \mathbf{R}^d) = \partial h(\mathbf{R}^d)$  for all  $t \ge 0$  by the first argument of the proof of Theorem 1. Plugging functions  $\varphi_n \in C_o(\mathbf{R}^d)$   $(n \ge 1)$  into (1.4), where the sequence  $\{\varphi_n\}_{n\ge 1}$  is nondecreasing and approximates the constant function  $\varphi(x) \equiv 1$ , and sending  $n \to \infty$ , we get

$$\int_{\mathbf{R}^d} (u(t,x) - h(x)) dx = \int_0^t ds \int_{\mathbf{R}^d} w(R, u(s, \cdot), dx) = t \cdot w(R, h, \mathbf{R}^d).$$
(4.10)

## Hitoshi Ishii, Toshio Mikami

(e) We argue by contradiction. Suppose that  $\hat{u}(t,\cdot)=\overline{u}(t,\cdot)$  for some  $t\geq s.$  Then we have

$$\int_{\mathbf{R}^d} (u(t,x) - h(x)) dx < \int_{\mathbf{R}^d} (\hat{u}(t,x) - \hat{h}(x)) dx$$
(4.11)

since

$$u(t,\cdot) - \hat{u}(t,\cdot) \le u(s,\cdot) - \hat{u}(s,\cdot) \ne h - \hat{h},$$

in view of (c). Hence

$$\int_{\mathbf{R}^d} (u(t,x) - h(x)) dx < \int_{\mathbf{R}^d} (\overline{u}(t,x) - \hat{h}(x)) dx, \tag{4.12}$$

which is a contradiction in view of (d), since (A.4) with h replaced by  $\hat{h}$  holds (see Remark 2, (ii) in Sect. 2).

Q. E. D.

(Proof of Theorem 3). By (a) in Theorem 2, (1.4) is equivalent to

$$\int_{[0,\infty)\times\mathbf{R}^d}\varphi(t,x)[u(dt,x)dx - dtw(R,u(t,\cdot),dx)] = 0$$
(4.13)

for all  $\varphi \in C_o([0,\infty) \times \mathbf{R}^d)$ .

(Step I). We first show that u is a viscosity subsolution of (1.5).

Let  $\psi \in C^2((0,\infty) \times \mathbf{R}^d)$  and assume that  $u - \psi$  attains a maximum at  $(s,y) \in (0,\infty) \times \mathbf{R}^d$ . We may assume that  $u(s,y) = \psi(s,y)$ , so that  $u(t,x) < \psi(t,x)$  for all  $(t,x) \in (0,\infty) \times \mathbf{R}^d \setminus \{(s,y)\}$  (see [6]).

(i). Consider first the case when  $\hat{u}(s, y) = u(s, y)$ .

30

By adding to  $\psi$  the function  $(t, x) \mapsto A\{|t-s|^2+|x-y|^2\}$ , with a suitable A > 0, if necessary, we may assume that  $D^2\psi(s, y) > 0$  and that the following set

$$U_{\varepsilon}^{+} \equiv \{(t,x) \in (0,\infty) \times \mathbf{R}^{d} | \psi(t,x) - \varepsilon < u(t,x)\} \quad (\varepsilon > 0)$$
(4.14)

is contained in the set  $U_{(\varepsilon/A)^{1/2}}((s,y))$ .

In the same way as in (3.18)-(3.20), considering  $(u, \psi, -\varepsilon)$  instead of  $(u_1, u_2, \varepsilon t)$ , by the compactness of the closure of the set  $U_{\varepsilon}^+$ , one can show that if  $(\tau, \xi) \in U_{\varepsilon}^+$  and  $p \in \partial u(\tau, \xi)$ , then  $p \in \partial \psi(\tau, z) = \{D\psi(\tau, z)\}$  for some z for which  $(\tau, z) \in U_{\varepsilon}^+$  and that

$$\int_{U_{\varepsilon}^{+}} w(R, u(t, \cdot), dx) dt \leq \int_{U_{\varepsilon}^{+}} w(R, \psi(t, \cdot), dx) dt.$$
(4.15)

We argue by contradiction. Assume that the following holds:

$$\partial \psi(s, y) / \partial t > R(D\psi(s, y)) \det_{+} D^{2} \psi(s, y), \qquad (4.16)$$

since by the definition of  $\chi$ , we have

$$\chi(u, D\psi(s, y), s, y) = 1.$$

By reselecting  $\varepsilon > 0$  if necessary, we may assume that

$$\partial \psi(t,x)/\partial t > \varepsilon + R(D\psi(t,x)) \det_{+} D^{2}\psi(t,x), \quad \det_{+} D^{2}\psi(t,x) > 0 \quad (4.17)$$

for all  $(t,x) \in U_{\varepsilon/A}((s,y))$ . Hence in the same way as in (3.21)-(3.24), considering  $u - \psi + \varepsilon$  instead of  $u_1 - u_2 - \varepsilon t$ , we have

$$\varepsilon |U_{\varepsilon}^{+}| \leq \int_{U_{\varepsilon}^{+}} w(R, u(t, \cdot), dx) dt - \int_{U_{\varepsilon}^{+}} R(D\psi(t, x)) \det_{+} D^{2}\psi(t, x) dx dt \leq 0,$$
(4.18)

by (4.15), which is a contradiction.

(ii). Consider next the case when  $\hat{u}(s,y) < u(s,y).$ 

We have

$$\chi(u, D\psi(s, y), s, y) = 0,$$

from which we only have to show that

$$\partial \psi(s,y) / \partial t \le 0.$$
 (4.19)

(4.19) is true, since from (b) in Theorem 2, we have

$$u(t,y)=u(s,y) \quad \text{ for all } t\in (0,s),$$

from which we have

$$\psi(s, y) < \psi(t, y)$$
 for all  $t \in (0, s)$ .

(Step II). Next, we show that u is a viscosity supersolution of (1.5). Let  $\psi \in C^2((0,\infty) \times \mathbf{R}^d)$  and assume that  $u - \psi$  attains a minimum at  $(s,y) \in (0,\infty) \times \mathbf{R}^d$ . We may assume as well that  $u(s,y) = \psi(s,y)$ , so that  $u(t,x) > \psi(t,x)$  for all  $(t,x) \in (0,\infty) \times \mathbf{R}^d \setminus \{(s,y)\}$  (see [6]). By (a) in Theorem 2, we see that

$$\partial \psi(s, y) / \partial t \ge 0.$$

Hence we only have to consider the case when the following holds:

$$\chi^{-}(u, D\psi(s, y), s, y) = 1, \quad \det_{+} D^{2}\psi(s, y) > 0.$$

By subtracting from  $\psi$  the function  $(t, x) \mapsto A\{|t - s|^2 + |x - y|^2\}$ , with a sufficiently small A > 0, if necessary, we may assume that  $D^2\psi(s, y)$  is positive definite and that (3.25) holds. By Lemma 8, if  $\varepsilon > 0$  is sufficiently small, then for any  $(t, x) \in U_{\varepsilon}^-$ ,  $D\psi(t, x) \in \partial u(t, z)$  for some z for which  $(t, z) \in U_{\varepsilon}^-$ .

As in (Step I), we argue by contradiction. Suppose that

$$\partial \psi(s, y) / \partial t < R(D\psi(s, y)) \det_{+} D^{2} \psi(s, y).$$
(4.20)

Reselecting  $\varepsilon > 0$  sufficiently small, we may assume that

$$\partial \psi(t,x) / \partial t + \varepsilon < R(D\psi(t,x)) \det_{+} D^{2} \psi(t,x), \quad \det_{+} D^{2} \psi(t,x) > 0 \quad (4.21)$$

for all  $(t,x) \in U_{(\varepsilon/A)^{1/2}}((s,y))$ . Then in the same way as in (Step I), we get

$$\varepsilon |U_{\varepsilon}^{-}| \leq \int_{U_{\varepsilon}^{-}} [R(D\psi(t,x)) \det D^{2}\psi(t,x)dxdt - w(R,u(t,\cdot),dx)dt] \leq 0,$$
(4.22)

which is a contradiction.

Q. E. D.

(Proof of Theorem 4). By Lemma 9 and Theorems 1 and 3, we only have to show the following: for a solution u to (1.4) with  $u(0, \cdot) = h(\cdot)$  and any T > 0,

$$\sup\{|u(t,x) - h(x)| : (t,x) \in [0,T] \times \mathbf{R}^d\} < \infty.$$
(4.23)

This is true, since

$$u(t,x) - h(x) \tag{4.24}$$

$$\leq |U_1(o)|^{-1} \int_{|y| \leq 1} (u(t, x+y) - h(x+y) + u(t, x-y) - h(x-y)) dy/2 + |U_1(o)|^{-1} \int_{|y| \leq 1} (h(x+y) + h(x-y) - 2h(x)) dy/2 \leq t |U_1(o)|^{-1} \int_{\mathbf{R}^d} R(y) dy + C/2 < \infty,$$

by (A.1) and (A.8). Here we used the following. If u(t,x) - h(x) > 0, then by (b) in Theorem 2, for any  $y \in \mathbf{R}^d$ ,

$$u(t,x) = \hat{u}(t,x) \le (\hat{u}(t,x+y) + \hat{u}(t,x-y))/2 \le (u(t,x+y) + u(t,x-y))/2.$$

Q. E. D.

(Proof of Theorem 5). In  $\mathbf{R}^{d+1}$ , the moving ball with a fixed center and with radius given by

$$r(t) = (r_0 - (d+1)t)^{1/(d+1)} \qquad t \in [0, r_0/(d+1)), \tag{4.25}$$

with  $r_0 > 0$ , is a Gauss curvature flow. In particular, for fixed  $a \in \mathbf{R}^d$  and  $b \in \mathbf{R}$ , put

$$w(t, x; a, b, r_0) := \begin{cases} b - ((r_0 - (d+1)t)^{2/(d+1)} - |x-a|^2)^{1/2} \\ & \text{if } |x-a| \le (r_0 - (d+1)t)^{1/(d+1)}, \\ +\infty & \text{otherwise.} \end{cases}$$
(4.26)

Then  $w(\cdot; a, b, r_0)$  is a viscosity supersolution of (1.5) with  $R(p) = (1 + |p|^2)^{-(d+1)/2}$ . Applying Lemma 10 to u and  $w(\cdot; a, b, r)$ , where r > 0 and  $a \in \mathbf{R}^d$  are chosen arbitrarily, and  $b = h^r(a)$ , we find that

$$u(t,x) \le w(t,x;a,b,r)$$
 for all  $(t,x) \in [0,r/(d+1)) \times \mathbf{R}^d$ . (4.27)

In particular,

$$u(t,a) \le w(t,a;a,b,r) < h^r(a)$$
 for all  $(t,a) \in [0,r/(d+1)) \times \mathbf{R}^d$ . (4.28)

Let  $\theta > 1$ , which will be selected later, and put

$$\varepsilon := \sup_{x \in \mathbf{R}^d} (h(x) - h(\theta x)) (\ge h(o) - h(\theta o) = 0), \tag{4.29}$$

and define  $z: [0, \infty) \times \mathbf{R}^d \mapsto \mathbf{R}$  by

$$z(t,x) = \varepsilon + v(\theta^{2d}t, \theta x).$$
(4.30)

Then z is a viscosity supersolution of (1.5) with  $R(p) = (1 + |p|^2)^{-(d+1)/2}$ and satisfies

$$z(0,x) = \varepsilon + h(\theta x) \ge h(x) \qquad \text{for all } x \in \mathbf{R}^d.$$
(4.31)

Fix any r > 0. By (A.2)' we can choose a  $\theta > 1$  depending on r, which may be as close as required to 1, and  $\delta > 0$  and L > 0 depending on r and  $\theta$  such that

$$h(\theta x) > \delta |x| + h^r(x) \qquad \text{if } |x| \ge L. \tag{4.32}$$

Hence by (4.28), (4.31), and (4.32), if  $0 \le t < r/(d+1)$  and  $|x| \ge L$ , then

$$u(t,x) < h^{r}(x) < h(\theta x) - \delta |x| \le z(0,x) - \delta |x| \le z(t,x) - \theta |x|, \quad (4.33)$$

since for each  $x \in \mathbf{R}^d$  the function  $t \mapsto v(t, x)$  is non-decreasing in  $[0, \infty)$ .

In view of (4.31) and (4.33), again by Lemma 10, we have

$$u(t,x) \le z(t,x) \equiv \varepsilon + v(\theta^{2d}t,\theta x) \qquad \text{in } (0,r/(d+1)) \times \mathbf{R}^d.$$
(4.34)

Here we can choose  $\theta > 1$  arbitrarily close to 1 and, by (A.2)', the constant  $\varepsilon$ , as a function of  $\theta$ , converges to zero as  $\theta \downarrow 1$ . Hence, in view of the arbitrariness of r > 0, we have

$$u(t,x) \le v(t,x) \qquad \text{in } (0,\infty) \times \mathbf{R}^d. \tag{4.35}$$

Q. E. D.

## 5. Appendix

In this section we prove Remark 3.

**Proposition 1.** Let  $h \in C(\mathbf{R}^d : \mathbf{R})$ . Suppose that there exists a convex function  $h_0 : \mathbf{R}^d \mapsto \mathbf{R}$  such that  $h_0(x) \to \infty$  as  $|x| \to \infty$  and that (2.10) holds. Then for each  $\theta > 1$ , there exists  $\delta > 0$  such that the following holds:

$$\lim_{|x| \to \infty} [h(\theta x) - \delta |x| - h^r(x)] = \infty \quad \text{for all } r > 0,$$
(5.1)

$$\lim_{\theta \downarrow 1} \{ \sup_{x \in \mathbf{R}^d} [h(x) - h(\theta x)] \} = 0.$$
(5.2)

(Proof). Without loss of generality, we may assume that  $h_0 \ge 1$ .

For any  $\theta > 1$  and r > 0, by the convexity of  $h_0$ , the following holds: for any x and  $y \in \mathbf{R}^d$ , with  $|y| \leq r$ ,

$$h_0(x+y) = h_0(\theta^{-1}\theta x + (1-\theta^{-1})\theta(\theta-1)^{-1}y)$$
(5.3)  
$$\leq \theta^{-1}h_0(\theta x) + (1-\theta^{-1})h_0(\theta(\theta-1)^{-1}y) \leq \theta^{-1}h_0(\theta x) + C(r,\theta),$$

where  $C(r, \theta)$  is a constant, and also

$$h_0(x) \ge \varepsilon |x| \quad \text{for all } x \in \mathbf{R}^d,$$

where  $\varepsilon$  is a positive constant. Thus we have

$$\max_{|y| \le r} h_0(x+y) \le h_0(\theta x) - (1-\theta^{-1})\varepsilon \theta |x| + C(r,\theta).$$
(5.4)

## Hitoshi Ishii, Toshio Mikami

By (2.10) and the continuity of h and  $h_0$ , there exists a constant  $C_0 > 0$ such that

$$|h(x) - h_0(x)| \le C_0 \quad \text{for all } x \in \mathbf{R}^d.$$
(5.5)

We also have

$$h^{r}(x) \leq \sup_{|z| \leq r} h(x+z) + r \quad \text{for all } x \in \mathbf{R}^{d}$$
(5.6)

since  $U_r((x,y)) \subset \operatorname{epi}(h)$  for all  $y \ge \sup_{|z| \le r} h(x+z) + r$ .

Combining (5.4), (5.5), and (5.6), we get

$$h^{r}(x) - C_{0} - r \le h(\theta x) - (1 - \theta^{-1})\varepsilon\theta |x| + C_{0} + C(r, \theta) \quad \text{for all } x \in \mathbf{R}^{d},$$
 (5.7)

from which we see that

$$\lim_{|x|\to\infty} [h(\theta x) - \frac{1}{2}\varepsilon\theta(1-\theta^{-1})|x| - h^r(x)] = \infty,$$
(5.8)

and conclude that (5.1) holds with  $\delta = \frac{1}{2}\varepsilon\theta(1-\theta^{-1})$ .

We also have, by (5.3) with y = o, for any  $\theta > 1$ ,

$$h_0(x) - h_0(\theta x) \le \theta h_0(x) - h_0(\theta x) \le (\theta - 1)h_0(0).$$
 (5.9)

This together with (2.10) and (4.29) implies (5.2).

Q. E. D.

Acknowledgement: We would like to thank Prof. K. Ishii for informing us that Theorem 5 is similar to [3, Theorem 4.1] where they considered the

38

mean curvature flow with a convex coercive initial function by a different approach from ours.

#### References

- B. Andrews, Gauss curvature flow: the fate of the rolling stones, Invent. Math. 138, (1999) 151–161.
- I. J. Bakelman, Convex Analysis and Nonlinear Geometric Elliptic Equations (Springer-Verlag, 1994).
- G. Barles, S. Biton and O. Ley, Quelque résultats d'unicité pour l'equation du mouvement par courbure moyenne dans R<sup>N</sup>, preprint.
- D. Chopp, L. C. Evans, and H. Ishii, Waiting time effects for Gauss curvature flows, Indiana Univ. Math. J. 48, (1999) 311–334.
- B. Chow, Deforming convex hypersurfaces by the nth root of the Gaussian curvature, J. Differential Geom. 22, (1985) 117–138.
- M. G. Crandall, H. Ishii, and P.-L. Lions, User's guide to viscosity solutions of second order partial differential equations, Bull. Amer. Math. Soc. (N.S.) 27, (1992) 1–67.
- S. N. Ethier and T. G. Kurtz, Markov processes: characterization and convergence (Jone Wiley & Sons Inc., 1986).
- 8. W. J. Firey, Shapes of worn stones, Mathematika 21, (1974) 1-11.
- M. -H. Giga and Y. Giga, Crystalline and level set flow-convergence of a crystalline algorithm for a general anisotropic curvature flow in the plain, preprint.
- P. M. Girão, Convergence of a crystalline algorithm for the motion of a simple closed convex curve by weighted curvature, SIAM J. Numer. Anal. 32, (1995) 886-899.

- P. M. Girão and R. V. Kohn, Convergence of a crystalline algorithm for the heat equation in one dimension and for the motion of a graph by weighted curvature, Numer. Math. 67, (1994) 41-70.
- 12. C. E. Gutiérrez, The Monge-Ampère equation (Birkhäuser, 2001).
- N. Ikeda and S. Watanabe, Stochastic differential equations and diffusion processes (North-Holland/Kodansha, 1981).
- H. Ishii and T. Mikami, A two dimensional random crystalline algorithm for Gauss curvature flow, Adv. Appl. Prob. 34, (2002) 491-504.
- H. Ishii and T. Mikami, A mathematical model of the wearing process of a non-convex stone, SIAM J. Math. Anal. 33, (2001) 860-876.
- R. T. Rockafellar and R. J-B. Wets, Variational Analysis (Springer-Verlag, 1998).
- K. Tso, Deforming a hypersurface by its Gauss-Kronecker curvature, Comm. Pure Appl. Math. 38, (1985) 867–882.