



Title	UDP-D-galactose-dependent organization of the trans-Golgi network & Mechanisms for polar localization of boric acid channels in <i>Arabidopsis thaliana</i> [an abstract of dissertation and a summary of dissertation review]
Author(s)	汪, 社亮
Citation	北海道大学. 博士(農学) 甲第11997号
Issue Date	2015-09-25
Doc URL	http://hdl.handle.net/2115/60189
Rights(URL)	http://creativecommons.org/licenses/by-nc-sa/2.1/jp/
Type	theses (doctoral - abstract and summary of review)
Additional Information	There are other files related to this item in HUSCAP. Check the above URL.
File Information	Sheliang_Wang_abstract.pdf (論文内容の要旨)



[Instructions for use](#)

学位論文内容の要旨

博士の専攻分野名称：博士（農学）

氏名：Sheliang Wang

学位論文題名

UDP-D-galactose-dependent organization of the *trans*-Golgi network &
Mechanisms for polar localization of boric acid channels in *Arabidopsis thaliana*

(シロイヌナズナにおけるUDP-D-ガラクトースに依存したトランスゴルジ網の
構造維持およびホウ酸チャネルの細胞膜内偏在機構)

Boron (B) is essential for plant development but is toxic when presents in excess. In *Arabidopsis thaliana*, a boric acid channel NIP5;1 is required for B uptake under B deficiency. NIP5;1 is expressed in outermost cell layers of roots and shows polar localization toward the soil-side in the plasma membrane (PM). The polar localization is assumed to be important for the efficient uptake of boric acid from soil solution. However, the molecular mechanisms of PM targeting and polar localization of NIP5;1 and their physiological importance were not established.

In this study, the nature of a mutant developing intracellular aggregates containing GFP-tagged NIP5;1 was investigated. In addition, the mechanism of polar localization of NIP5;1 and its physiological importance on plant growth were elucidated.

1. UDP-D-galactose synthesis by UDP-glucose 4-epimerase 4 is required for organization of the *trans*-Golgi network/early endosome in *Arabidopsis thaliana* root epidermal cells

Previously, a fluorescence-imaging based screening isolated an *Arabidopsis* mutant developing intracellular aggregates containing GFP-NIP5;1 in epidermal cells of the root elongation zone. This mutant was identified as an allele of *UDP-glucose 4-epimerase 4 (uge4)* mutant which has defects in UDP-D-galactose synthesis and channeling into the Golgi.

Here, I further investigated the nature of intracellular aggregates in the *uge4* mutants. Live-cell imaging indicated that the markers of multi-vesicular bodies/late endosomes (MVB/LEs) and PM were strongly accumulated in the intracellular aggregates in the *uge4* mutant. Furthermore, electron microscopy analysis revealed accumulation of

abnormal high-electron-density vesicles in elongating epidermal cells. The abnormal vesicles were often associated or interconnected with *trans*-Golgi network/early endosomes (TGN/EEs) and contained ADP-ribosylation factor 1, which is usually localized to the Golgi and the TGN/EEs. Together, our data indicate the importance of UDP-D-galactose synthesis by UGE4 for the organization and function of endomembranes, especially TGN/EEs, which are a sorting station of the secretory and vacuolar pathways.

2. Polar localization of a boric acid channel is mediated by phosphorylation and plays a crucial role in growth of *Arabidopsis thaliana* under low boron conditions

To elucidate the mechanism of the polar localization of NIP5;1, I performed structure-localization analysis of NIP5;1 by expression of GFP-used homologs and chimera proteins in *Arabidopsis thaliana*. The N-terminal region of NIP5;1 was identified to be required for the polar localization and have the ability to target non-polar aquaporin proteins to the soliside PM domain. Further analysis revealed that the conserved threonine residues in the N-terminal region of NIP5;1 were phosphorylated and required for the polar localization.

I then investigated the physiological importance of the polar localization using transgenic *Arabidopsis* lines expressing a non-polar NIP5;1 variant. Although the non-polar variant showed comparable boric acid transport activity in *Xenopus* oocytes, the transgenic lines with the variant developed shorter roots than those with the wild-type NIP5;1 under low B conditions.

These results revealed that the polar localization of NIP5;1 is mediated by phosphorylation, and plays a crucial role in growth of *Arabidopsis thaliana* under low boron conditions probably by improving the efficiency of boric acid uptake in the roots.