



Title	Trilobodrilus itoi sp nov., with a Re-Description of T. nipponicus (Annelida: Dinophilidae) and a MolecularPhylogeny of the Genus
Author(s)	Kajihara, Hiroshi; Ikoma, Maho; Yamasaki, Hiroshi; Hiruta, Shimpei F.
Citation	Zoological science, 32(4), 405-417 <a href="https://doi.org/10.2108/zs140251">https://doi.org/10.2108/zs140251</a>
Issue Date	2015-08
Doc URL	<a href="http://hdl.handle.net/2115/62716">http://hdl.handle.net/2115/62716</a>
Type	article
File Information	zs_32_p405-.pdf



[Instructions for use](#)

# *Trilobodrilus itoi* sp. nov., with a Re-Description of *T. nipponicus* (Annelida: Dinophilidae) and a Molecular Phylogeny of the Genus

Hiroshi Kajihara<sup>1\*</sup>, Maho Ikoma<sup>1</sup>, Hiroshi Yamasaki<sup>2</sup>, and Shimpei F. Hiruta<sup>1</sup>

<sup>1</sup>Faculty of Science, Hokkaido University, Sapporo 060-0810, Japan

<sup>2</sup>Faculty of Science, University of the Ryukyus, Senbaru 1, Nishihara, Nakagami, Okinawa 903-0213, Japan

The marine interstitial annelid *Trilobodrilus itoi* sp. nov., the sixth member of the genus, is described on the basis of specimens collected intertidally at Ishikari Beach, Hokkaido, Japan; this is the second species in the genus described from the Pacific Rim. In addition, *T. nipponicus* Uchida and Okuda, 1943 is re-described based on fresh topotypic material from Akkeshi, Hokkaido, Japan. From both species, we determined sequences of the nuclear 18S and 28S rRNA genes, and the mitochondrial cytochrome c oxidase subunit I (COI) gene. Molecular phylogenetic trees based on concatenated sequences of the three genes showed that *T. itoi* and *T. nipponicus* form a clade, which was the sister group to a clade containing the two European congeners *T. axi* Westheide, 1967 and *T. heideri* Remane, 1925. The Kimura two-parameter distance for COI was 22.5–22.7% between *T. itoi* and *T. nipponicus*, comparable with interspecific values in other polychaete genera. We assessed the taxonomic utility of epidermal inclusions and found that the known six species can be classified into three groups.

**Key words:** North Pacific, marine invertebrates, archiannelid, mesopsammon, meiofauna, interstitial

## INTRODUCTION

The dinophilid annelid genus *Trilobodrilus* Remane, 1925 comprises marine interstitial ‘archiannelids’ that reach approximately 2 mm in body length and lack typical polychaete features such as distinct body segmentation, parapodia, and chitinous chaetae. It currently contains five named species: *T. heideri* Remane, 1925 and *T. axi* Westheide, 1967 in European waters (e.g., Remane, 1925; Boaden, 1963; Westheide, 1967; Ax, 1968; Scharnoffske, 1986); *T. indicus* Chandrasekhara Rao, 1973 from India (Chandrasekhara Rao, 1973); *T. nipponicus* Uchida and Okuda, 1943 from Japan (Uchida and Okuda, 1943), India (Chandrasekhara Rao and Ganapati, 1968), and the Pacific coast of the USA (Wieser, 1957); and *T. hermaphroditus* Riser, 1999 from the Atlantic coast of the USA (Riser, 1999). The hypothesis that these and other dinophilids have evolved from an ancestor most closely related to dorvilleid polychaetes in Eunicida (e.g., Westheide, 1987; Eibye-Jacobsen and Kristensen, 1994) was not supported by molecular phylogenetic analyses (Struck et al., 2002, 2005), and the sister taxon to Dinophilidae has not clearly been established (e.g., Rousset et al., 2007; Struck et al., 2008; Zrzavý et al., 2009). The family as currently diagnosed (Westheide, 1984) contains 16 species in three genera (Read, 2014). Dinophilids and other interstitial an-

nelid families, such as Diurodrilidae (Worsaae and Rouse, 2008; Golombek et al., 2013), Nerillidae (Worsaae et al., 2005), Protodrilidae (Bailey-Brock et al., 2010; Di Domenico et al., 2013; Martínez et al., 2013, 2014), and Saccocirridae (Di Domenico et al., 2014a, b, c) are important for understanding annelid evolution overall (Worsaae and Kristensen, 2005; Struck, 2006; Westheide, 2008) and the biogeography (Curini-Galletti et al., 2012) and macroevolutionary patterns (Rundell and Leander, 2010) of microscopic animals.

The Pacific species *Trilobodrilus nipponicus* needs re-description, as some of the morphological characters considered to be useful in distinguishing among congeners, e.g., the shape of ‘epidermal inclusions’ (see below), were not known for this species, nor were DNA sequence data available. Westheide (1967) pointed out that *T. nipponicus* is morphologically superficially similar to the European species *T. heideri*, and that a more detailed taxonomic description of the former is necessary to differentiate between them. While *T. nipponicus* has been reported from several localities distant from the type locality subsequent to the original description, insufficient original descriptions and lack of DNA data call into question previous species identifications of *Trilobodrilus* specimens worldwide (e.g., Rieger and Rieger, 1975).

Under light microscopy, specimens of *Trilobodrilus* show within the epidermis two types of small bodies up to 15 µm long: (1) a mosaic or aggregation of droplets, and (2) an elongated, darker body with granular contents. The terminology for these small bodies has been inconsistent in the literature. Westheide (1967) originally used the term “Epidermiseinschlüsse” [epidermis inclusions] collectively to

\* Corresponding author. Tel. : +81-11-706-2755;  
Fax : +81-11-706-4851;  
E-mail: kazi@mail.sci.hokudai.ac.jp

Supplemental material for this article is available online.  
doi:10.2108/zs140251

refer to both types, but called the aggregations of droplets “polygonele Felder” [polygonal fields] and the other, granular type “spindelförmige Drüsen” [spindle-shaped glands]. Chandrasekhara Rao (1973) did not use a collective term to refer to the two types, but called the former type “epidermal glands” and the latter “bacillary glands”. Riser (1999) likewise did not use a collective term, but called the former type “epidermal inclusions” and the latter “spindle glands”. Westheide (1967), who initially discovered epidermal inclusions in *Trilobodrilus*, noted differences in the inclusions between *T. axi* and *T. heideri*, which suggested the small bodies might provide useful taxonomic characters. Although the ‘epidermis inclusions’ *sensu* Westheide (1967) have been succinctly described in *T. indicus* (Chandrasekhara Rao, 1973) and *T. hermaphroditus* (Riser, 1999), their structure has not been explored in detail, nor has their taxonomic significance been rigorously assessed with statistical tests.

While Yamanishi (1983) listed 16 species of marine interstitial annelids reported from Japanese waters, representing 11 genera, these tiny worms remain understudied in this region. During the course of a faunal survey of marine interstitial animals around Hokkaido, northern Japan, we recognized two species of *Trilobodrilus*, one of which was referable to *T. nipponicus*, whereas the other turned out to be new to science. In this paper, we describe and illustrate these two species, and report the results of molecular phylogenetic analyses to infer their systematic position, making use of dinophilid sequences available in public databases. We also report intra- and interspecific variation in epidermis inclusions of the mosaic type (the “polygonele Felder” of Westheide [1967]), which we henceforth refer to as ‘epidermal inclusions’ *sensu* Riser (1999), and discuss the taxonomic utility of variation in these structures.

## MATERIALS AND METHODS

### Sampling and observation

Sampling was conducted at two localities, Ishikari Beach and Akkeshi Bay, Hokkaido, northern Japan (Fig. 1). Specimens were extracted from intertidal sediment samples by freshwater shock followed by stirring and decantation: sediments were brought back to the laboratory and agitated briefly but intensively in tap water in a bucket; the suspension was passed through a Gwen’s mermaid bra (Nybakken and Higgins, 2007) with a 32- $\mu\text{m}$ -mesh net; and the residue was immediately transferred to seawater to avoid rupture of the soft-bodied animals. Living worms were collected with a Pasteur pipette under a dissecting microscope and photographed with a Nikon D5200 digital camera attached to a Nikon SMZ 1500 microscope by adapters (NYPIXS2-3166, NY1S-FA, and NY1S-1501750, Micronet), with a pair of external strobe lights (Hikaru Komachi Di, Morris, Japan). For light microscopy, specimens were observed under an Olympus BX51 compound microscope after being anaesthetized in a  $\text{MgCl}_2$  solution isotonic to seawater and photographed with the same digital camera. For scanning electron microscopy (SEM), specimens were anaesthetized as just described and then fixed in formalin–seawater, dehydrated in an ethanol series, dried in a Hitachi HCP-2  $\text{CO}_2$  critical-point drier, mounted on an aluminium stub, coated with gold in a JEOL JFC-1100 ion sputter coater, and observed with a Hitachi S-3000N scanning electron microscope at 15-kV accelerating voltage. Measurements were taken from scaled digital images using ImageJ ver. 1.48 (Rasband, 1997–2014). Type and voucher material has been deposited in the Hokkaido University Museum (ZIHU), Sapporo, Japan.




Fig. 1. Map of Hokkaido, northern Japan, showing the sampling sites and the surrounding ocean currents.




Fig. 2. Schematic drawing of an epidermal inclusion, defining length and width, and the terminology for the components.

### Analyses of epidermal inclusions

Terminology and standard measurements pertaining to the mosaic-type epidermal inclusions treated here are given in Fig. 2. Each epidermal inclusion consists of an ‘envelope’ containing a number of ‘spherules’. The envelope is embedded in the epidermis, but may connect with the exterior via a small opening. The spherule situated closest to the opening is usually (but not always) the largest; we call the largest spherule the ‘major spherule’, corresponding to the “cap-like unit” of Riser (1999). Within the envelope there may be a ‘vacuolar region’ that lacks spherules, usually at the end opposite the major spherule. Because the envelope is so thin and the spherules are packed so tightly within it, the envelope itself cannot be observed by light microscopy; its outline distinct from the spherules is evident only in the vacuolar region. In the drawings of Westheide (1967, fig. 5) and Chandrasekhara Rao (1973, figs. 5–6), the envelope is not illustrated at all; in the photographs of Westheide (1967, fig. 6) and Riser (1999, fig. 1), the outline of the envelope is unclear. In this paper, to compare the measurements from *T. itoi* and *T. nipponicus* with those from other species, the length of the epidermal inclusion is defined as the long axis of the aggregation of spherules, rather than of the envelope itself, and the

width of the epidermal inclusion as the short axis of the spherule aggregation (Fig. 2).

To assess intraspecific variation in epidermal inclusions, the length and width of 11–26 epidermal inclusions were measured for each of five specimens of *T. itoi* sp. nov. Measurements were made from digital images, and differences in the variance and mean of the length-to-width ratio among the five specimens were examined by Bartlett's test and Tukey's HSD test, respectively. Interspecific variation in the epidermal inclusions was assessed in the same manner. For previously described species, lengths and widths were measured from figures in the published literature: Westheide (1967, figs. 5, 6) for *T. axi* ( $n = 13$ ) and *T. heideri* ( $n = 15$ ), Chandrasekhara Rao (1973, fig. 4) for *T. indicus* ( $n = 2$ ), and Riser (1999, fig. 1) for *T. hermaphroditus* ( $n = 15$ ). For *T. nipponicus*, 21 epidermal inclusions were measured from one specimen. Statistical tests were performed with R ver. 3.1.0 (R Core Team, 2014).

#### DNA extraction, PCR amplification, and sequencing

Total DNA was extracted by using a DNeasy Tissue Kit (Qiagen, USA). Fragments of the nuclear 18S and 28S rRNA genes (18S and 28S, respectively), and the mitochondrial cytochrome *c* oxidase subunit I gene (COI), were amplified by polymerase chain reaction (PCR) with the primer pairs listed in Table 1. PCR conditions were 95°C for 1 min; 35 cycles of 95°C for 30 sec, 45°C for 90 sec, and 72°C for 3 min (90 sec for COI); and 72°C for 7 min. Sequences were determined with the primers listed in Table 1 by using a Big-Dye Terminator Kit ver. 3.1 (Life Technologies, USA) and a 3730 DNA analyzer (Life Technologies, USA).

For the new species, we also obtained longer sequences by next-generation sequencing (NGS). DNA concentration was measured with a NanoDrop 8000 Spectrophotometer (Thermo Fisher Scientific, USA). DNA (100 ng) was fragmented using an Ion Xpress™ Plus Fragment gDNA Library Kit (Ion Torrent, USA). Fragments were ligated to barcode adapters using an Ion Xpress™ Barcode Adapters 33–48 Kit (Ion Torrent, USA). Using 2% E-Gel®

SizeSelect™ Agarose Gels (Invitrogen, USA), the resultant barcode library was subjected to size selection, targeting fragments 480 nt long. After eight cycles of PCR amplification, the size-selected library was purified twice with AMPureXP (Beckman Coulter, USA), and the quantity of the library was determined with a High Sensitivity DNA Kit (Agilent Technologies, USA). Based on this library, emulsion-PCR-based beads were prepared by using an Ion PGM™ Template OT2 400 Kit (Ion Torrent, USA). Sequencing was carried out with an Ion Torrent PGM™ sequencer, using an Ion 314™ Chip ver. 2 (Ion Torrent, USA)/Ion PGM™ Sequencing 400 Kit (Ion Torrent, USA). Base calling was done with an Ion Torrent Server, with the output for each barcode in the form of binary sequence alignment/map (BAM)-format files. The adapter sequences in the BAM files were eliminated by using CLC Genomic Workbench ver. 7 (CLC Bio, USA); in addition, *de novo* assembly was performed with CLC Genomic Workbench ver. 7, yielding 3155 contigs; these were exported in FASTA format and subjected to similarity searches using BLAST+ ver. 2.2.29.

#### Molecular phylogenetic analyses

To infer phylogenetic relationships among species in *Trilobodrilus*, a maximum-likelihood (ML) analysis and Bayesian inference (BI) were carried out that included sequences from our two species and four species of dinophilids for which sequences were available in the public databases (Table 2). Because the sister taxon to Dinophilidae is uncertain (e.g., Rousset et al., 2007), outgroups were chosen to include taxa in seven polychaete families, with special emphasis on Dorvilleidae (cf. Eibye-Jacobsen and Kristensen, 1994). The sipunculan *Sipunculus nudus* Linnaeus, 1766 was used to root the trees (Kvist and Siddall, 2013; Weigert et al., 2014). Sequences were aligned gene by gene by using MUSCLE (Edgar, 2004) implemented in MEGA ver. 5.2 (Tamura et al., 2011) with the following settings: Gap Open = –400; Gap Extend = 0; Max Iterations = 8; Clustering Method (Iteration 1, 2) = neighbor joining; Clustering Method (Other Iterations) = neighbor joining; Min Diag Length

**Table 1.** List of primers used for amplification and sequencing, with original references.

Gene	Primer name	Reaction*	Primer sequence (in 5'–3' direction)	Direction	Source
18S rRNA	K18SF	A/S	GTCATATGCTTGTCTTAAAGATTAAGC	Forward	Present study
	K18SR	A/S	GGAAACCTTGTTACGACTTTTACTTCA	Reverse	Present study
	F2	S	CCTGAGAAACGGCTRCCACAT	Forward	Yamaguchi and Endo (2003)
	F3	S	GYGRTCAGATACCRCCSTAGTT	Forward	Yamaguchi and Endo (2003)
	F4	S	GGTCTGTGATGCCCTYAGATGT	Forward	Yamaguchi and Endo (2003)
	R6	S	TYTCTCRKGCTBCCTCTCC	Reverse	Yamaguchi and Endo (2003)
	R7	S	GYYARAACCTAGGGCGGTATCTG	Reverse	Yamaguchi and Endo (2003)
	R8	S	ACATCTRAGGGCATCACAGACC	Reverse	Yamaguchi and Endo (2003)
28S rRNA	D1F	A/S	GGGACTACCCCTGAATTTAAGCAT	Forward	Park and Ó Foighil (2000)
	28Sb	A/S	TCGGAAGGAACAGCTAC	Reverse	Whiting et al. (1997)
	28S-01	A/S	GACTACCCCTGAATTTAAGCAT	Forward	Kim et al. (2000)
	28Sr	A/S	ACACACTCCTTAGCGGA	Reverse	Luan et al. (2005)
	28Sf	A/S	TGGGACCCGAAAGATGGTG	Forward	Luan et al. (2005)
	28S-3KR	A/S	CCAATCCTTTTCCCGAAGTT	Reverse	Yamasaki et al. (2013)
	28S-2KF	A/S	TTGGAATCCGCTAAGGAGTG	Forward	Yamasaki et al. (2013)
	28jj-3'	A/S	AGTAGGGTAAAACCTAACCT	Reverse	Palumbi (1996)
	28S-n05R	S	CTCACGGTACTTGTTCGCTAT	Reverse	Yamasaki et al. (2013)
	28SR-01	S	GACTCCTTGGTCCGTGTTTCAAG	Reverse	Kim et al. (2000)
	28S-15R	S	CGATTAGTCTTTTCGCCCTA	Reverse	Yamasaki et al. (2013)
	28S-3KF	S	AGGTGAACAGCCTCTAGTCG	Forward	Yamasaki et al. (2013)
	28v-5'	S	AAGGTAGCCAAATGCCTCATC	Forward	Palumbi (1996)
	28S-42F	S	GAGTTTGACTGGGGCGGTA	Forward	Yamasaki et al. (2013)
	COI	LCO1490	A/S	GGTCAACAAATCATAAAGATATTGG	Forward
HCO2198		A/S	TAAACTTCAGGGTGACCAAAAAATCA	Reverse	Folmer et al. (1994)

\*A, PCR amplification; S, cycle sequencing

**Table 2.** List of species included in the phylogenetic analysis, with public database accession numbers.

Taxa	Markers			Sources
	18S	28S	COI	
<b>Ingroup (Dinophilidae)</b>				
<i>Dinophilus gyrocolliatus</i> Schmidt, 1857	AF412805	—	—	Struck et al. (2002)
<i>Dinophilus</i> sp.	FJ200245	FJ200246	—	Worsaae and Rouse (2008)
<i>Trilobodrilus axi</i> Westheide, 1967	AF412806	AY894292	—	Struck et al. (2002, 2005)
<i>Trilobodrilus heideri</i> Remane, 1925	AF412807	AY732231	—	Struck et al. (2002, 2005)
<i>Trilobodrilus itoi</i> sp. nov.	DRA001682	DRA001682	AB924371	Present study
<i>Trilobodrilus nipponicus</i> Uchida and Okuda, 1943	LC009446	LC009447	LC009445	Present study
<b>Outgroup (Dorvilleidae)</b>				
<i>Dorvillea erucaeformis</i> (Malmgren, 1865)	AY176285	AY838859	AY838868	Worsaae et al. (2005); Struck et al. (2006)
<i>Exallopus jumarsi</i> Blake, 1985	—	—	JQ310755	Wiklund et al. (unpubl.)
<i>Iphitime hartmanae</i> Kirkegaard, 1977	—	—	GQ415472	Wiklund et al. (2009)
<i>Microdorvillea</i> sp.	AY527051	—	—	Struck (unpubl.)
<i>Ophryotrocha labronica</i> La Greca and Bacci, 1962	AY838855	DQ790047	AY838874	Struck et al. (2006, 2007)
<i>Parougia eliasoni</i> (Oug, 1978)	—	DQ790053	GQ415489	Struck et al. (2007); Wiklund et al. (2009)
<i>Pettiboneia urciensis</i> Campoy and San Martin, 1980	AF412801	—	—	Struck et al. (2002)
<i>Protodorvillea kefersteini</i> (McIntosh, 1869)	AF412799	AY732230	KF808171	Struck et al. (2002, 2005); Aylagas et al. (2014)
<i>Schistomeringos longicornis</i> (Ehlers, 1901)	—	HM473268	HM473662	Carr (unpubl.)
<b>Outgroup (others)</b>				
<i>Cirratulus spectabilis</i> (Kinberg, 1866)	AY708536	DQ790029	—	Burnette et al. (2005); Struck et al. (2007)
<i>Diurodrilus subterraneus</i> Remane, 1934	KC790349	KC790349	KC790350	Golombek et al. (2013)
<i>Eunice</i> sp.	AF412791	AY732229	—	Struck et al. (2002, 2005)
<i>Eurythoe complanata</i> (Pallas, 1766)	AY364851	AY364849	JN086548	Jördens et al. (2004); Borda et al. (2012)
<i>Nereis vexillosa</i> Grube, 1851	DQ790083	DQ790043	HM473512	Struck et al. (2007); Carr et al. (2011)
<i>Pectinaria gouldii</i> (Verrill, 1873)	DQ790091	DQ790054	FJ976040	Struck et al. (2007); Zhong et al. (unpubl.)
<i>Sipunculus nudus</i> Linnaeus, 1766	DQ300006	DQ300048	DQ300162	Schulze et al. (2007)

(lambda) = 24. Alignment-ambiguous regions were removed by using BMGE v. 1.1 (Criscuolo and Gribaldo, 2010); the 18S and 28S alignments were processed with the “-t DNA” option, and COI with the “-t CODON” option. The lengths of the resulting sequence alignments were 1535 nt (18S), 1735 nt (28S), and 645 nt (COI); the genes were concatenated for phylogenetic analyses by using MEGA ver. 5.2.

The ML analysis was conducted in RAxML ver. 8 (Stamatakis, 2014) under the GTR+G model; for 18S and 28S, the data were partitioned by gene; for COI by codon position. Nodal support values were estimated by bootstrapping with 1000 pseudoreplicates. RAxML was called as follows (except the options for input and output file names, as well as the partition file name): `raxmlHPC-PTHREADS-AVX -T 5 -f a -x12345 -p 12345 -#1000 -m GTR-GAMMA`.

Bayesian inference was conducted by using MrBayes ver. 3.2.3 (Ronquist and Huelsenbeck, 2003; Altekar et al., 2004), with two independent Metropolis-coupled analyses, each using four Markov chains of 1,000,000 generations. Trees were sampled every 100 generations. Run convergence was assessed by using Tracer ver. 1.6 (Rambaut et al., 2014). For BI, jModeltest2 (Darriba et al., 2012) was used to determine the most suitable substitution model for each gene partition under the Bayesian information criterion, with the following settings: number of substitution schemes = 3; including models with equal/unequal base frequencies (+F); including models with/without a proportion of invariable sites (+I); including models with/without rate variation among sites (+G) (nCat = 4); optimized free parameters (K) = substitution parameters + 29 branch lengths + topology; base tree for likelihood calculations = ML tree; tree topology search operation = NNI. The optimal models were K80+I+G for 18S, GTR+G for 28S, and GTR+I+G for COI.

Kimura (1980) two-parameter (K2P) genetic distances were calculated by using MEGA ver. 5.2.


## RESULTS

### Taxonomy

#### *Trilobodrilus itoi* sp. nov. (Fig. 3)

**Material examined.** Syntypes, ZIHU 4885, 17 specimens, Au-coated and mounted on SEM stub, collected by M. Ikoma and H. Kajihara, 15 January 2012, Ishikari Beach (43°15'26"N, 141°21'26"E; type locality), about 2 km south-southwest from the mouth of the Ishikari River, Hokkaido, northern Japan; from surface layer of sediment (medium sand) to about 10 cm depth at water's edge. Eleven specimens, same locality as for syntypes, collected by M. I., 30 September 2013, observed alive by light microscopy, destroyed after observation except for two non-type specimens: ZIHU 4886 (fixed in 10% formalin-seawater, mounted in glycerine on a glass slide, with the edge of the coverslip sealed with Canada balsam) and ZIHU 4887 (fixed in Bouin's fluid, mounted in the same manner as ZIHU 4886). More than 50 specimens collected sporadically at the type locality from 29 October 2012 to 30 September 2013, all destroyed after observation by light microscopy and DNA extraction.

**Sequences.** The following sequences were determined by standard sequencing from a single, non-type specimen collected on 17 December 2012, for which no morphological voucher remains: COI, AB924371 (658 nt, coding 219 aa); 18S rRNA, AB924372 (1714 nt); 28S rRNA, AB924373 (1120 nt). In addition, 75,340 reads (equivalent to 18.6 M bases) by new-generation sequencing from eight specimens



**Fig. 3.** *Trilobodrilus itoi* sp. nov. (A, F, G, J) Living, non-type specimens, destroyed by compression during observation; (B–E, H) SEM images, syntype, ZIHU 4885. (A) Anaesthetized whole specimen, photographed under dissecting microscope. (B) Head, front-lateral view. (C) Head, dorso-lateral view; dotted circle indicates wide gap in third ciliary band. (D) Head, lateral view. (E) Whole body, ventral view. (F) Photomicrograph of epidermal inclusions. (G) Magnification of epidermal inclusion. (H) Whole body, lateral view. (J) Photomicrograph of whole body.

in all, collected on 30 September 2013, have been deposited in the DDBJ Sequence Read Archive under accession number DRA001682. The complete 18S sequence with

parts of the ETS and ITS regions (3463 nt), as well as partial 28S sequences including part of the ITS region (4620 nt), both assembled from DRA001682, are available as Supple-

mentary Files S1 and S2, respectively.

**Diagnosis.** *Trilobodrilus* with second ciliary band incomplete dorsally, where single mid-dorsal ciliary tuft is present; third ciliary band dorsally discontinuous, with large gap.

**Description.** Body 0.9–1.3 mm long, 0.08–0.12 mm wide ( $n = 11$ ); colorless, translucent; stomach light yellow; salivary glands opaque white, sometimes tinged with green (Fig. 3A); oocytes cream white. Head with four apical ciliary tufts on anterior end, arranged dorsally and ventrally on both sides (Fig. 3B), as in congeners. Three ciliary bands present anteriorly (Fig. 3B–D); first band completely encircling head near tip (Fig. 3B, C); second band situated just anterior to mouth, dorsally discontinuous (Fig. 3C); third band post-oral on segment 1, dorsally discontinuous, with wide gap (Fig. 3C). Single, long, mid-dorsal ciliary tuft in gapped portion of second band, oriented slightly anteriorly (Fig. 3B, C). Pair of nuchal organs in segment 1, one on each side, just anterior to third ciliary band (Fig. 3D). Small, short ciliary tufts distributed sporadically on sides of body (Fig. 3E, H). In anaesthetized and fixed conditions, apparently 6–7 segments (except pygidium), although segmental boundaries posterior to segment 2 obscure; pygidium with unclear boundary (Fig. 3A, E, H). Pair of salivary glands (Fig. 3A, J) between segments 2 and 3. Width of central ciliary tract (Fig. 3E) about 40% (35–44%,  $n = 4$ ) of body diameter in middle of body. Epidermal inclusions highly variable in shape, mostly elliptical, up to  $7 \times 12 \mu\text{m}$  in size (Fig. 3F, G); spherules 3–22 (mostly 7–8) in number ( $n = 86$ , five specimens) (see below “Epidermal inclusions”). Spindle glands not found (Fig. 3F). No mature male found. Females contained up to two oocytes, each up to  $150 \mu\text{m}$  in diameter, arranged in tandem beneath intestine (Fig. 3J); “storage cells” or “coelenchyme” of Franssen (1980, fig. 4A), up to  $30 \mu\text{m}$  in diameter, packed within coelom in mature females (Fig. 3J). Mature individuals with oocytes found in September and October.

**Remarks.** *Trilobodrilus itoi* differs from congeners, except for *T. nipponicus*, in having a mid-dorsal ciliary tuft in the gapped portion of the second ciliary band on the head (Table 3). *Trilobodrilus itoi* differs from *T. nipponicus* in having the third ciliary band (on the first body segment) discontinuous dorsally (vs continuous in *T. nipponicus*), and the ventral ciliary tract is narrow, about 40% of body width (vs wide, about 45% in *T. nipponicus*).

**Ecology.** At Ishikari Beach, *T. itoi* appeared to be narrowly restricted to the swash zone; none were found land-

ward in subsurface water. Densities were locally variable; in calm sea conditions, up to 20 individuals were collected per 300 ml of sand. The salinity during the sampling period was generally 30; the water temperature was  $12^\circ\text{C}$  on 29 October 2012 and  $4^\circ\text{C}$  on 17 December 2012. Itô (1985) found the sediment grain size at the site ( $\phi$  scale; Krumbein, 1937) to be 1.2–1.7. Animals that co-occurred in the habitat included proseriate flatworms (one of which was *Archotoplana yamadai* Tajika, 1983); harpacticoid copepods (*Arenopontia ishikariana* Itô, 1968 and *Leptastacus japonicus* Itô, 1968); nematodes (genera *Axonolaimus*, *Bathylaimus*, *Enoploides*, *Enoplolaimus*, *Epacanthion*, *Epsilonema*, *Mesacanthion*, *Metoncholaimus*, *Neochromadora*, *Oncholaimus*, *Parasaveljevia*, *Rhynchonema*, *Theristus*, and *Trissonchulus*) (D. Shimada, pers. comm.); the non-interstitial gammarid amphipod *Anisogammarus pugettensis* (Dana, 1853) (cf. Hoshide, 1971); and the mysid *Archaeomysis articulata* Hanamura, 1997 (cf. Hanamura, 1999). With sporadic water changes, individuals of *T. itoi* were able to survive for more than a month without feeding in a Petri dish placed in an incubator at  $12^\circ\text{C}$ .

**Etymology.** The specific name, a noun in the genitive case, is in honor of Dr Tatsunori Itô (1945–1990), a Japanese meiobenthologist and crustacean systematist, whose handbook for the general public, *Organisms in Sand Interstices* (Itô, 1985), influenced several successive generations in Japan. Ishikari Beach, the type locality, was one of Itô’s primary research sites early in his career (e.g., Itô, 1970).

***Trilobodrilus nipponicus* Uchida and Okuda, 1943**  
(Fig. 4)

*Trilobodrilus nipponicus* Uchida and Okuda, 1943, p. 301, figs. 1, 2.

?*Trilobodrilus nipponicus*: Wieser (1957), p. 284, fig. 4; Chandrasekhara Rao and Ganapati (1968), p. 28, fig. 5.


**Material examined.** ZIHU 4946, 22 specimens, Au-coated and mounted on three SEM stubs, collected on a narrow beach about 50 m long at the base of a steep cliff near Cape Aikappu, Akkeshi, Hokkaido ( $43^\circ00'58''\text{N}$ ,  $144^\circ50'02''\text{E}$ ) by M. Ikoma, S. F. Hiruta, and H. Kajihara, 23 September 2014. In addition, one specimen examined alive with light microscopy was destroyed after observation, and four other specimens were used for DNA extraction.

**Sequences.** The following sequences were determined

**Table 3.** Morphological characters and their states in the six species in *Trilobodrilus*.

Species	Body length (mm)	Body width (mm)	Number of mid-dorsal tuft on 2nd ciliary band	3rd ciliary band dorsally continuous (+) or not (–)	Size of epidermal inclusions ( $\mu\text{m}$ )	Number of spherules per envelope	Mean ratio of length to width of epidermal inclusions	Spindle glands present (+) or absent (–)	Stomach color	Sex	Habitat	Source
<i>T. axi</i> Westheide, 1967	1	0.1	0	– <sup>a</sup>	$10 \times 15$	5–10	1.81	+	?	dioecious	intertidal	Westheide (1967)
<i>T. heideri</i> Remane, 1925	1.5–1.9	0.1–0.2	0	–	?	3–10	1.41	+	brown	dioecious	subtidal	Remane (1925); Westheide (1967)
<i>T. hermaphroditus</i> Riser, 1999	~1.7	~0.22	2	+ <sup>b</sup>	$6 \times 9$	3–7	1.42	+	greenish brown	hermaphroditic	intertidal/subtidal	Riser (1999)
<i>T. indicus</i> Chandrasekhara Rao, 1973	1.0–1.2	0.1–0.12	0	+	12	7–11	2.42	+	transparent?	dioecious	intertidal	Chandrasekhara Rao (1973)
<i>T. itoi</i> sp. nov.	0.9–1.3	0.08–0.12	1	–	$7 \times 9$	3–22	1.45	–	yellowish	dioecious	intertidal	Present study
<i>T. nipponicus</i> Uchida and Okuda, 1943	0.7–1.4	0.09–0.16	1	+	$5.5 \times 16$	9–13	2.28	–	pale green or yellowish brown	dioecious	intertidal	Uchida and Okuda (1943); present study

<sup>a</sup>Comprised of three rows, the 3rd row present only on ventral side; <sup>b</sup>horse-shoe shaped, not meeting ventral ciliary tract



**Fig. 4.** *Trilobodrilus nipponicus* Uchida and Okuda, 1943. **(A)** ZIHU 4946, photographed alive under dissecting microscope before preparation for SEM. **(B–E)** ZIHU 4946, SEM images. **(B)** Head, dorsal view; dotted circle indicates dorsally continuous third ciliary band. **(C)** Lateral view of whole body; arrowheads indicate lateral ciliary tufts. **(D)** Ventral view of whole body. **(E)** Middle portion of body, ventrolateral view; arrowheads showing lateral ciliary tufts. **(F)** Photomicrograph of epidermal inclusions.

by standard sequencing from four specimens, for which no morphological vouchers remain: COI, LC009442–LC009445 (658 nt, coding 219 aa); 18S rRNA, LC009446 (from one specimen, 1784 nt); 28S rRNA, LC009447 (from one specimen, 3596 nt).

**Diagnosis.** *Trilobodrilus* with second ciliary band dorsally incomplete, with single mid-dorsal tuft of cilia; third ciliary band nearly complete dorsally.

**Description.** Body 0.7–1.4 mm long, 0.09–0.16 mm wide ( $n = 23$ ); colorless, translucent (Fig. 4A); stomach orange yellow; oocytes opaque white. Ciliary pattern same as for *T. itoi*, except: 1) third ciliary band continuous dorsally ( $n = 14$ ), although sometimes sparse ( $n = 7$ ) (Fig. 4B), and 2) width of ventral ciliary tract about 52% (46–55%,  $n = 4$ )

of body width (Fig. 4D). Epidermal inclusions elongate, 11–16  $\mu\text{m}$  long, 3.5–5.5  $\mu\text{m}$  wide, with 9–13 spherules per envelope (Fig. 4F); size ratio of major spherule to other spherules in each envelope appearing to be greater than that in *T. itoi* (Figs. 3G, 4F). Spindle glands not found.


**Remarks.** Our topotypic material agrees completely with the original description of *T. nipponicus*, except that lateral ciliary tufts were sparse in our material (Fig. 4C, E), rather than regularly arranged on each segment as described and illustrated by Uchida and Okuda (1943, fig. 1). *Trilobodrilus nipponicus* has been distinguished morphologically from congeners in having a single, mid-dorsal ciliary tuft on the head. Chandrasekhara Rao and Ganapati (1968) identified specimens from India as *T. nipponicus*

because they had this type of mid-dorsal tuft. Because *T. itoi* also has a mid-dorsal tuft, the Indian form can no longer be positively identified as *T. nipponicus*. Rieger and Rieger (1975) also depicted a mid-dorsal tuft in an unidentified form collected from subtidal sand off Beaufort, North Carolina, Atlantic coast, USA; its identity should be investigated in future studies. A form from the Pacific coast near Seattle, USA, identified by Wieser (1957) as *T. nipponicus* lacks a mid-dorsal tuft; because the second ciliary band is continuous dorsally (Wieser, 1957, fig. 4), this form probably represents a different species.

**Ecology.** Although sediment-particle sizes were not measured, the sediment on the beach at Akkeshi appeared to be coarser than that at Ishikari Beach. *Trilobodrilus nipponicus* was extremely patchily distributed. We collected sediment at ~20 randomly chosen points along the beach in a zone a few meters landward or seaward from the water's edge, but detected *T. nipponicus* in only one sediment sample. We found 29 specimens in a 300 ml of sediment sample, in which unidentified acoels, harpacticoids, and nematodes predominated; fragments of cirratulid branchial filaments were also found. Uchida and Okuda (1943) reported that "the Archiannelid, *Saccocirrus major*, the Rhabdocoelid Turbellarian, *Thylacorhynchus* sp. and some nemerteans" co-occurred with *T. nipponicus*. While we did observe saccocirrids, they did not appear to utilize the same microhabitat as *T. nipponicus*. An unidentified species of *Proschizorhynchella* (Kalyptorhynchia: Schizorhynchia) co-occurred at low frequency (K. Tamura pers. comm.). Nemerteans in the genus *Cephalothrix* were found at the beach, but no species in *Ototyphlonemertes*, an interstitial ribbon-worm genus often found intertidally. We examined sediment on the northern and western coasts of Akkeshi Bay, and on two other beaches on the eastern coast (south of the type locality), but did not find *Trilobodrilus*. Sediments were gravely on the northern coast, silty on the western coast, and composed of finer sand on the two southeastern beaches.

### Epidermal inclusions


Spherules could be observed by light-microscopy when the animal was slightly squashed under a cover slip, but forceful squashing was necessary to count the number of spherules in a single envelope. After forceful squashing, however, the spherule(s) partly or entirely exuded from the envelope, especially near the edge of the body (Fig. 5). We



**Fig. 5.** *Trilobodrilus itoi* sp. nov., photomicrograph of epidermal inclusions near edge of body; arrowheads indicate major spherules appearing to be squeezed out of epidermis.

eventually gave up counting the exact number of spherules in each envelope, because 1) smaller spherules below 0.2  $\mu\text{m}$  in diameter (i.e., approaching the limit of the optical resolution for a 100  $\times$  oil-immersion lens) occurred between larger spherules, and 2) the spherules were not arranged at the same focal depth, with the deeper portion obscured.

The mean value in the ratio of length to width of the epidermal inclusions among the five specimens of *T. itoi* ranged from 1.35 to 1.55 (Fig. 6A), with no significant differences among the five in variation ( $P = 0.13$ : Bartlett's test) or mean



**Fig. 6.** Standard box-and-whisker plots of the length-width ratio of epidermal inclusions; the box encompasses the first and third quartiles; the thick horizontal bar within the box depicts the median; and whiskers depict the true minimum and maximum values, except where outliers (circles) greater than 1.5 times the interquartile range were identified. **(A)** Intraspecific variation in *Trilobodrilus itoi* sp. nov. among five specimens (specimen No. 1 was a mature, but tiny, female 0.9 mm in body length; Nos. 2–5 were immature specimens, 1.1–1.3 mm in body length). **(B)** Interspecific variation among six congeners; means with the same lower-case letter at the top of the graph are not significantly different based on Tukey's HSD test ( $\alpha = 0.05$ ). Data from the literature: *T. axi* and *T. heideri*, Westheide (1967); *T. indicus*, Chandrasekhara Rao (1973); *T. hermaphroditus*, Riser (1999).

( $P = 0.30\text{--}0.99$ ; Tukey's HSD test).

Significant differences in variation were observed among the six species of *Trilobodrilus* when two outliers (one each in *T. heideri* and *T. nipponicus*) were included ( $P = 0.0038$ ; Bartlett's test). Excluding these outliers resulted in no significance ( $P = 0.014$ ,  $\alpha = 0.01$ ); the greater  $P$  value was likely due to the small sample size in *T. indicus* ( $n = 2$ ). Excluding the outliers, three groups of mean length-width ratio values were evident: i) *T. axi* (1.81); ii) *T. heideri* (1.41), *T. hermaphroditus* (1.42), and *T. itoi* (1.45); and iii) *T. indicus* (2.42) and *T. nipponicus* (2.28) (Fig. 6B).

### Phylogeny and genetic distance

In the best ML tree ( $\ln L = -23774.622593$ ) (Fig. 7) and BI tree (Supplementary Fig. S1), Dinophilidae is monophyletic, with 97% bootstrap frequency (BF) and 1.00 posterior probability (PP). Within Dinophilidae, *T. itoi* and *T. nipponicus* form a clade (97% BF, 1.00 PP), as do the two European species, *T. axi* and *T. heideri* (96% BF, 1.00 PP). These two clades together comprise a monophyletic group (67% BF, 0.98 PP). *Dinophilus* sp. *sensu* Worsaae and Rouse (2008) is the sister group to *Trilobodrilus* spp. (97% BF, 1.00 PP), and *Dinophilus gyrociliatus* Schmidt, 1857, is the sister group to all other species in Dinophilidae (97% BF, 1.00 PP). Our results thus refute the suggestion (Westheide, 1967) that, based on morphological similarity, *T. heideri* might be more closely related to *T. nipponicus* than to *T. axi*.


Our study corroborated that of Struck et al. (2002, 2005) in failing to support the morphology-based hypothesis (Eibye-Jacobsen and Kristensen, 1994) of a close relationship between Dinophilidae and Dorvilleidae. All the dorvilleids included in our analyses (except for *Pettiboneia urciensis* Campoy and San Martin, 1980) formed a clade (91% BF, 1.00 PP) that was the sister group to *Eunice* sp. (64% BF, 0.99 PP).

The interspecific K2P genetic distance for COI based on one specimen of *T. itoi* and four of *T. nipponicus* was 22.5–22.7%. These values are comparable with interspecific K2P distances for COI in other genera of interstitial polychaetes, e.g., 23.6–27.0% among six species in *Protodrilus* (Di Domenico et al., 2013; Martines et al., 2013), and in genera of non-interstitial annelids, e.g., an average of 26.2% in *Hydroides* (Sun et al., 2012) and 19.2–26.2% among three species of *Glycera* (Schüller, 2011).

### DISCUSSION

We discovered two species of *Trilobodrilus* along the shores of Hokkaido Island, one at Ishikari Beach on the west coast, and the other in Akkeshi Bay on the east, with the two localities 1050 km (northward course) and 1230 km (southward course) apart in along-shore distances (Fig. 1). While we did not record precise physical/geological data for the habitat such as granulometry, the two beaches do differ in physical properties. In beach morphodynamics (Wright and Short, 1984), Ishikari Beach is the intermediate to dissipative type, with finer sand, while the beach in Akkeshi Bay is more or less reflective, with coarser sand. Our sampling site on Ishikari Beach, which is a stretch of sandy shore 25 km long, is 2 km from a river mouth, while the small, unnamed beach in Akkeshi Bay, roughly 50 m long, appeared to receive a continuous input of organic detritus from the nearby rocky and particulate shores that are rich in macroalgae and sea grasses. In addition to differences in the physical properties between the two beaches, they are under the influence of different ocean currents: the warm Tsushima Current flowing north off Ishikari Beach on the Sea of Japan, and the cold Oyashio Current flowing south outside Akkeshi Bay on the Pacific Ocean (Fig. 1).

Overall, the distribution of the two species appeared to be quite restricted, differing widely in scales ranging from microhabitat to the geographical and oceanographic levels. In this context, the putative wide distribution of *T. nipponicus* across the Indo-Pacific (Wieser, 1957; Chandrasekhara Rao and Ganapati, 1968) is likely the result



**Fig. 7.** Phylogenetic tree resulting from maximum-likelihood analysis based on combined COI, 18S, and 28S rRNA gene sequences ( $\ln L = -23774.622593$ ). Numbers near nodes indicate maximum-likelihood bootstrap support values in percent (> 60%) and posterior probability from Bayesian inference (> 0.90), respectively.

of misidentification (see Remarks above), as opponents of the “everything is everywhere” hypothesis (for review, De Wit and Bouvier, 2006; Fontaneto and Brodie, 2011; Williams, 2011) have put forward (e.g., Coleman, 2002; Foissner, 2006; Taylor et al., 2006). The species in *Trilobodrilus* so far investigated are all direct developers lacking a planktonic larval phase (Uchida and Okuda, 1943; Ax, 1968), which suggests relatively low dispersal ability, which in turn may be related to the restricted distributions observed in this study. The phylogeographic pattern in our phylogeny, in which the two Japanese species comprise the sister group to a clade containing the two European forms, may also be related to the poor dispersal potential in *Trilobodrilus*.

Species in *Trilobodrilus* were thought to be morphologically similar, distinguished by “the number of body segments, structure of head and pygidium, distribution of ciliary rings, lateral tufts of cilia, tactile bristles, structure of epidermal glands [sic], etc.” (Chandrasekhara Rao, 1973), until the fourth congener (*T. indicus*) was described. While the fifth species (*T. hermaphroditus*) differs from all of its congeners in being hermaphroditic, Riser (1999) pointed out that the body segmentation can vary during specimen preparation (pressure under the cover slip, anaesthetization, and fixation), and other characters, such as the four anterior tactile bristles (ciliary tufts) on the tip of the head, are present in all congeners and thus have little value in distinguishing between species. With regard to the epidermal inclusions, our observations indicated that 1) counting the number of spherules per envelope is impractical because there can be a number of extremely small (< 0.2 µm) spherules at different focus depths, but 2) the mean length–width ratio (L/W) may be of taxonomic utility, with three groups of species having been detected: *T. indicus* and *T. nipponicus* (elongate type, L/W > 2.0); *T. axi* (medium type, 1.5 ≤ L/W ≤ 2.0); and *T. heideri*, *T. hermaphroditus*, and *T. itoi* (oval type, L/W < 1.5).

The effectiveness of incorporating DNA data in studies of meiofaunal taxonomy has become apparent for many taxa, including Nemertodermatida (Meyer-Wachsmuth et al., 2014), Proseriata (Casu and Curini-Galletti, 2006; Scarpa et al., 2015), Gastrotricha (Kieneke et al., 2012), Nemertea (Tulchinsky et al., 2012; Leasi and Norenburg, 2014), Annelida (Di Domenico et al., 2014c; Martínez et al., 2014), and Rotifera (Fontaneto, 2014). While our SEM observations revealed some subtle differences in ciliation pattern between *T. itoi* and *T. nipponicus*, before comparing their DNA sequences we were uncertain whether these two represented different species. DNA taxonomy (e.g., Fontaneto et al., 2009), or “turbo-taxonomy”—an approach combining COI sequences, concise morphological descriptions, and high-resolution digital imaging (Butcher et al., 2012; Riedel et al., 2013)—will likely be essential in future studies to elucidate the diversity of these morphologically character-poor meiofauna, especially for material from areas previously not investigated.

#### ACKNOWLEDGMENTS

We thank Professor Matthew H. Dick for reviewing and editing the manuscript; two anonymous reviewers for constructive criticism on an earlier version of the manuscript; Professor/Director Masahiro Nakaoka and other staff at the Akkeshi Marine Station for providing

facilities; Naoya Takeda and Kohtarō Tamura for information on Kalyptorhynchia; Daisuke Shimada for information on nematodes; Shohei Yamauchi for help in collecting; Daisuke Shimada and Shinri Tomioka for assisting with SEM observation; and Keiichi Kakui for assisting with photography. This study was financially supported by JSPS KAKENHI Grant Numbers 23370038 and 2565013.

#### REFERENCES

- Altekar G, Dwarkadas S, Huelsenbeck JP, Ronquist F (2004) Parallel Metropolis-coupled Markov chain Monte Carlo for Bayesian phylogenetic inference. *Bioinformatics* 20: 407–415
- Ax P (1968) Das Fortpflanzungsverhalten von *Trilobodrilus* (Archiannelida, Dinophilidae). *Mar Biol* 1: 330–335
- Aylagas E, Borja Á, Rodríguez-Ezpeleta N (2014) Environmental status assessment using DNA metabarcoding: towards a genetics based marine biotic index (GAMBI). *PLoS ONE* 9: e90529 doi:10.1371/journal.pone.0090529
- Bailey-Brock JH, Jouin-Toulmond C, Brock RE (2010) Protodrilidae (Annelida: Polychaeta) from the Hawaiian Islands and comparison with specimens from French Polynesia. *Pac Sci* 64: 463–472
- Blake JA (1985) Polychaeta from the vicinity of deep-sea geothermal vents in the eastern Pacific. I. Euprosinidae, Phyllococidae, Hesionidae, Nereididae, Glyceridae, Dorvilleidae, Orbiniidae, and Maldanidae. *Bull Biol Soc Wash* 6: 67–101
- Boaden PJS (1963) Behaviour and distribution of the archiannelid *Trilobodrilus heideri*. *JMBA UK* 43: 239–250
- Borda E, Kudenov JD, Bienhold C, Rouse G (2012) Towards a revised Amphinomidae (Annelida, Amphinomida): description and affinities of a new genus and species from the Nile Deep-sea Fan, Mediterranean Sea. *Zool Scr* 41: 307–325
- Burnette AB, Struck TH, Halanych KM (2005) Holopelagic *Poeobius meseres* (“Poeobiidae,” Annelida) is derived from benthic flabelligerid worms. *Biol Bull* 208: 213–220
- Butcher BA, Smith MA, Sharkey MJ, Quicke DLJ (2012) A turbo-taxonomic study of Thai *Aleiodes* (*Aleiodes*) and *Aleiodes* (*Arcaleiodes*) (Hymenoptera: Braconidae: Rogadinae) based largely on COI barcoded specimens, with rapid descriptions of 179 new species. *Zootaxa* 3457: 1–232
- Campoy A, San Martín G (1980) *Pettiboneia urciensis* sp. n.: un nouveau Dorvilleidae (Polychètes: Errantes) de la Méditerranée. *Cah Biol Mar* 21: 201–207
- Carr CM, Hardy SM, Brown TM, Macdonald TA, Hebert PD (2011) A tri-oceanic perspective: DNA barcoding reveals geographic structure and cryptic diversity in Canadian polychaetes. *PLoS ONE* 6: e22232 doi:10.1371/journal.pone.0022232
- Casu M, Curini-Galletti M (2006) Genetic evidence for the existence of cryptic species in the mesopsammic flatworm *Pseudomonocelis ophiocephala* (Rhabditophora: Proseriata). *Biol J Linn Soc* 87: 553–576
- Chandrasekhara Rao G (1973) *Trilobodrilus indicus* n. sp. (Dinophilidae, Archiannelida) from Andhra coast. *Proc Indian Acad Sci Sect B* 77: 101–108
- Chandrasekhara Rao G, Ganapati PN (1968) On some archiannelids from the beach sands of Waltair Coast. *Proc Indian Acad Sci Sect B* 67: 24–30
- Criscuolo A, Gribaldo S (2010) BMGE (block mapping and gathering with entropy): selection of phylogenetic informative regions from multiple sequence alignments. *BMC Evol Biol* 10: 210
- Coleman AW (2002) Microbial eukaryote species. *Science* 297: 337
- Curini-Galletti M, Artois T, Delogu V, De Smet WH, Fontaneto D, Jondelius U, et al. (2012) Patterns of diversity in soft-bodied meiofauna: dispersal ability and body size matter. *PLoS ONE* 7: e33801 doi:10.1371/journal.pone.0033801
- Dana JD (1853) Crustacea. Part II. United States Exploring Expedition During the Years 1838, 1839, 1840, 1841, 1842 under the Command of Charles Wilkes, U.S.N., Vol 13. Sherman,

- Philadelphia, pp 689–1618
- Darriba D, Taboada GL, Doallo R, Posada D (2012) jModelTest2: more models, new heuristics and parallel computing. *Nat Methods* 9: 772
- De Wit R, Bouvier T (2006) ‘Everything is everywhere, but, the environment selects’; what did Baas Becking and Beijerinck really say? *Environ Microbiol* 8: 755–758
- Di Domenico M, Martínez A, Almeida TCM, Martins MO, Worsaae K, Lana PC (2014a) Response of the meiofaunal annelid *Saccocirrus pussicus* (Saccocirridae) to sandy beach morphodynamics. *Hydrobiologia* 734: 1–16
- Di Domenico M, Martínez A, Amaral ACZ, Lana P da C, Worsaae K (2014b) Saccocirridae (Annelida) from the southern and south-eastern Brazilian coasts. *Mar Biodiv* 44: 313–325
- Di Domenico M, Martínez A, Lana P da C, Worsaae K (2013) *Protodrilus* (Protodrilidae, Annelida) from the southern and southeastern Brazilian coasts. *Helgol Mar Res* 67: 733–748
- Di Domenico M, Martínez A, Lana P, Worsaae K (2014c) Molecular and morphological phylogeny of Saccocirridae (Annelida) reveals two cosmopolitan clades with specific habitat preferences. *Mol Phylogenet Evol* 75: 202–218
- Edgar RC (2004) MUSCLE: a multiple sequence alignment method with reduced time and space complexity. *BMC Bioinformatics* 5: 113
- Ehlers E (1901) Die Polychaeten des magellanischen und chilenischen Strandes. Ein faunistischer Versuch. Festschrift zur Feier des hundertfünfzigjährigen Bestehens des königlichen Gesellschaft der Wissenschaften zu Göttingen. Abteilungen der mathematisch-physikalischen Klasse. Weidmann, Berlin
- Eiby-Jacobsen D, Kristensen RM (1994) A new genus and species of Dorvilleidae (Annelida, Polychaeta) from Bermuda, with a phylogenetic analysis of Dorvilleidae, Iphitimidae and Dinophilidae. *Zool Scr* 23: 107–131
- Foissner W (2006) Biogeography and dispersal of microorganisms: a review emphasizing protists. *Acta Protozool* 45: 111–136
- Folmer O, Black M, Hoeh W, Lutz R, Vrijenhoek R (1994) DNA primers for amplification of mitochondrial cytochrome c oxidase subunit I from diverse metazoan invertebrates. *Mol Mar Biol Biotech* 3: 294–299
- Fontaneto D (2014) Molecular phylogenies as a tool to understand diversity in rotifers. *Int Rev Hydrobiol* 99: 178–187
- Fontaneto D, Brodie J (2011) Why biogeography of microorganisms? In “Biogeography of Microscopic Organisms: Is Everything Small Everywhere?” Ed by D Fontaneto, Cambridge University Press, Cambridge, pp 3–10
- Fontaneto D, Kaya M, Herniou EA, Barraclough TG (2009) Extreme levels of hidden diversity in microscopic animals (Rotifera) revealed by DNA taxonomy. *Mol Phylogenet Evol* 53: 182–189
- Fransen ME (1980) Ultrastructure of coelomic organization in annelids. I. Archannelids and other small polychaetes. *Zoomorphologie* 95: 235–249
- Golombek A, Tobergte S, Nesnidal MP, Purschke G, Struck TH (2013) Mitochondrial genomes to the rescue—Diurodrilidae in the myzostomid trap. *Mol Phylogenet Evol* 68: 312–326
- Grube AE (1851) Annulaten. In “Reise in den äussersten Norden und osten Sibiriens während der Jahre 1843 und 1844 mit allerhöchster Genehmigung auf Veranstaltung der kaiserlichen Akademie der Wissenschaften zu St. Petersburg ausgeführt und in Verbindung mit vielen Gelehrten herausgegeben, Vol 2, Part 1” Ed by A Th von Middendorff, Kaiserlichen Akademie der Wissenschaften, St. Petersburg, pp 1–24
- Hanamura Y (1997) Review of the taxonomy and biogeography of shallow-water mysids of the genus *Archaeomysis* (Crustacea: Mysidacea) in the North Pacific Ocean. *J Nat Hist* 31: 669–711
- Hanamura Y (1999) Seasonal abundance and life cycle of *Archaeomysis articulata* (Crustacea: Mysidacea) on a sandy beach of western Hokkaido, Japan. *J Nat Hist* 33: 1811–1830
- Hoshide K (1971) Studies on gregarines from Japan II. *Cephaloidophora anisogammari* n. sp. and *Cephaloidophora elongata* n. sp. from amphipoda. *J Fac Sci Hokkaido Univ Ser VI Zool* 18: 186–192
- Itô T (1968) Description and records of marine harpacticoid copepods from Hokkaido I. *J Fac Sci Hokkaido Univ Ser VI Zool* 16: 369–381
- Itô T (1970) The study of interstitial fauna on Ishikari coast, Hokkaido. *Benthos Res* 1: 21–23 [in Japanese]
- Itô T (1985) Organisms in Sand Interstices. Kaimeisha, Tokyo [in Japanese]
- Jördens J, Struck T, Purschke G (2004) Phylogenetic inference regarding Parergodrilidae and *Hrabeiella periglandulata* (‘Polychaeta’, Annelida) based on 18S rDNA, 28S rDNA and COI sequences. *J Zool Syst Evol Res* 42: 270–280
- Kieneke A, Martínez Arbizu PM, Fontaneto D (2012) Spatially structured populations with a low level of cryptic diversity in European marine Gastrotricha. *Mol Ecol* 21: 1239–1254
- Kim C-G, Zhou H-Z, Imura Y, Tominaga O, Su Z-H, Osawa S (2000) Pattern of morphological diversification in the *Leptocarabus* ground beetles (Coleoptera: Carabidae) as deduced from mitochondrial ND5 gene and nuclear 28S rRNA sequences. *Mol Biol Evol* 17: 137–145
- Kimura M (1980) A simple method for estimating evolutionary rates of base substitutions through comparative studies of nucleotide sequences. *J Mol Evol* 16: 111–120
- Kinberg JGH (1866) *Annulata nova*. *Ofv Ak Forh* 22: 239–258
- Kirkegaard JB (1977) A new species of *Iphitime* (Polychaeta: Iphitimidae) living under the tail of *Hyas* (Crustacea: Decapoda) in the Oslo Fjord. In “Essays on Polychaetous Annelids in Memory of Dr Olga Hartman” Ed by DJ Reich, K Fauchald, Allan Hancock Foundation, University of Southern California Press, Los Angeles, pp 199–209
- Krumbein WC (1937) The sediments of Barataria Bay. *J Sed Petrol* 7: 3–17
- Kvist S, Siddall ME (2013) Phylogenomics of Annelida revised: a cladistic approach using genome-wide expressed sequence tag data mining and examining the effects of missing data. *Cladistics* 29: 435–448
- La Greca M, Bacci G (1962) Una nuova specie di *Ophryotrocha* delle coste tirreniche (Annelida, Polychaeta). *Bollettin Zool* 29: 7–18
- Leasi F, Norenburg JL (2014) The necessity of DNA taxonomy to reveal cryptic diversity and spatial distribution of meiofauna, with a focus on Nemertea. *PLoS ONE* 9: e104385 doi:10.1371/journal.pone.0104385
- Linnaeus C (1766) *Systema naturae per regna tria naturae, secundum classes, ordines, genera, species, cum characteribus, differentiis, synonymis, locis, tomus I, editio duodecima, reformata*. Salvius, Stockholm
- Luan Y-X, Mallatt JM, Xie R-D, Yang Y-M, Yin W-Y (2005) The phylogenetic positions of three basal-hexapod group (Protura, Diplura, and Collembola) based on ribosomal RNA gene sequences. *Mol Biol Evol* 22: 1579–1592
- Malmgren AJ (1865) Nordiska hafs-annulater. *Ofv Ak Forh* 22: 181–192
- Martínez A, Di Domenico M, Jörgen K, Norenburg JL, Worsaae K (2013) Description of three new species of *Protodrilus* (Annelida, Protodrilidae) from Central America. *Mar Biol Res* 9: 676–691
- Martínez A, Di Domenico M, Rouse GW, Worsaae K (2014) Phylogeny and systematics of Protodrilidae (Annelida) inferred with total evidence analyses. *Cladistics* (early view) doi:10.1111/cla.12089
- McIntosh WC (1869) On the structure of British nemerteans, and some new British annelids. *Tr R Soc Edinb* 25: 305–433
- Meyer-Wachsmuth I, Curini Galletti M, Jondelius U (2014) Hyper-

- cryptic marine meiofauna : species complexes in Nemertodermitida. *PLoS ONE* 9(9): e107688. doi:10.1371/journal.pone.0107688
- Nybakken JW, Higgins RP (2007) Intertidal meiobenthos. In "The Light and Smith Manual: Intertidal Invertebrates from Central California to Oregon, Completely Revised and Expanded" Ed by JT Carlton, University of California Press, Oakland, CA, pp 18–23
- Oug E (1978) New and lesser known Dorvilleidae (Annelida, Polychaeta) from Scandinavian and northeast American waters. *Sarsia* 63: 285–303
- Pallas PS (1766) *Miscellanea zoologica quibus novae imprimis atque obscurae animalium species describuntur et observationibus iconibusque illustrantur*. Pieter van Cleef, The Hague
- Palumbi SR (1996) Nucleic acids II: The polymerase chain reaction. In "Molecular Systematics, 2nd ed" Ed by DM Hills, C Moritz, BK Mable, Sinauer Associates, Sunderland, Massachusetts, pp 205–247
- Park J-K, Ó Foighil D (2000) Sphaeriid and corbiculid clams represent separate heterodont bivalve radiations into freshwater environments. *Mol Phylogenet Evol* 14: 75–88
- R Core Team (2014) R: A Language and Environment for Statistical Computing. R Foundation for Statistical Computing, Vienna, available at <http://www.R-project.org>
- Rambaut A, Suchard MA, Xie W, Drummond AJ (2014) Tracer v 1.6, available at <http://beast.bio.ed.ac.uk/Tracer>
- Rasband WS (1997–2014) ImageJ, U. S. National Institutes of Health, Bethesda, Maryland, USA, <http://imagej.nih.gov/ij/>
- Read G (2014) Dinophilidae Remane, 1925. In "World Polychaeta Database" Ed by G Read, K Fauchald, World Register of Marine Species, <http://www.marinespecies.org/aphia.php?p=taxdetails&id=155089>, accessed on 10 December 2014
- Remane A (1925) Diagnosen neuer Archianneliden. *Zool Anz* 65: 15–17
- Remane A (1934) *Diurodrilus subterraneus* nov. spec., ein Archiannelide aus dem Küstengrundwasser. *Schr Ver Schleswig-Holst* 20: 479
- Riedel A, Sagata K, Suhardjono YR, Tänzler R, Balke M (2013) Integrative taxonomy on the fast track—towards more sustainability in biodiversity research. *Front Zool* 10: 15
- Rieger MM, Rieger GE (1975) Fine structure of the pharyngeal bulb in *Trilobodrillus* and its phylogenetic significance within Archiannelida. *Tissue Cell* 7: 267–279
- Riser NW (1999) Description of a new species of dinophilid polychaete, with observations on other dinophilids and interstitial polychaetes in New England. *Northeast Nat* 6: 211–220
- Ronquist F, Huelsenbeck JP (2003) MrBayes 3: Bayesian phylogenetic inference under mixed models. *Bioinformatics* 19: 1572–1574
- Rousset V, Pleijel F, Rouse GW, Erséus C, Siddall ME (2007) A molecular phylogeny of annelids. *Cladistics* 23: 41–63
- Rundell R, Leander BS (2010) Masters of miniaturization: convergent evolution among interstitial eukaryotes. *BioEssays* 32: 430–437
- Scarpa F, Cossu P, Sanna D, Lai T, Norenburg JL, Curini-Galletti M, et al. (2015) An 18S and 28S-based clock calibration for marine Proseriata (Platyhelminthes). *J Exp Mar Biol Ecol* 463: 22–31
- Scharnoffsk P (1986) Ultrastructure of sperm morphology of *Trilobodrillus axi* and *T. heideri* (Dinophilidae, Polychaeta). *Helgol Meeresunters* 40: 419–430
- Schmidt O (1857) Zur Kenntnis der Turbellaria Rhabdocoela und einiger anderer Würmer des Mittelmeeres. *SB Ak Wien* 23: 347–366
- Schüller M (2011) Evidence for a role of bathymetry and emergence in speciation in the genus *Glycera* (Glyceridae, Polychaeta) from the deep Eastern Weddell Sea. *Polar Biol* 34: 549–564
- Schulze A, Cutler EB, Giribet G (2007) Phylogeny of sipunculan worms: a combined analysis of four gene regions and morphology. *Mol Phylogenet Evol* 42: 171–192
- Stamatakis A (2014) RAxML version 8: a tool for phylogenetic analysis and post-analysis of large phylogenies. *Bioinformatics* 30: 1312–1313
- Struck TH (2006) Progenetic species in polychaetes (Annelida) and problems assessing their phylogenetic affiliation. *Integr Comp Biol* 46:558–568
- Struck TH, Halanych KM, Purschke G (2005) Dinophilidae (Annelida) is most likely not a progenetic Eunicida: evidence from 18S and 28S rDNA. *Mol Phylogenet Evol* 37: 619–623
- Struck TH, Nesnidal MP, Purschke G, Halanych KM (2008) Detecting possibly saturated positions in 18S and 28S sequences and their influence on phylogenetic reconstruction of Annelida (Lophotrochozoa). *Mol Phylogenet Evol* 48: 628–645
- Struck TH, Purschke G, Halanych KM (2006) Phylogeny of Eunicida (Annelida) and exploring data congruence using a partition addition bootstrap alteration (PABA) approach. *Syst Biol* 55: 1–20
- Struck TH, Schult N, Kusen T, Hickman E, Bleidorn CB, McHugh D, et al. (2007) Annelid phylogeny and the status of Sipuncula and Echiura. *BMC Evol Biol* 7: 57 doi:10.1186/1471-2148-7-57
- Struck TH, Westheide W, Purschke G (2002) Progenesis in Eunicida ("Polychaeta," Annelida)—separate evolutionary events? Evidence from molecular data. *Mol Phylogenet Evol* 25: 190–199
- Sun Y, Kupriyanova EK, Qiu JW (2012) COI barcoding of *Hydroides*: a road from impossible to difficult. *Invert Syst* 26: 539–547
- Tajika K-I (1983) Zwei neue interstitielle Turbellarien der Gattung *Archotoplana* (Proseriata, Otoplanidae) aus Hokkaido, Japan. *J Fac Sci Hokkaido Univ Ser VI Zool* 23: 179–194
- Tamura K, Perterson D, Peterson N, Stecher G, Nei M, Kumar S (2011) MEGA5: molecular evolutionary genetics analysis using maximum likelihood, evolutionary distance, and maximum parsimony methods. *Mol Biol Evol* 28: 2731–2739
- Taylor JW, Turner E, Townsend JP, Dettman JR, Jacobson D (2006) Eukaryotic microbes, species recognition and the geographic limits of species: examples from the kingdom Fungi. *Phil Trans Roy Soc B* 361: 1947–1963
- Tulchinsky AY, Norenburg JL, Turbeville JM (2012) Phylogeography of the marine interstitial nemertean *Ototyphlonemertes parmula* (Nemertea, Hoplonemertea) reveals cryptic diversity and high dispersal potential. *Mar Biol* 159: 661–674
- Uchida T, Okuda S (1943) A new species of Archiannelida, *Trilobodrillus nipponicus* n. sp. *J Fac Sci Hokkaido Imp Univ Ser VI Zool* 8: 301–305
- Verrill AE (1873) Report upon the invertebrate animals of Vineyard Sound and the adjacent waters, with an account of the physical characters of the region. In "Report on the Condition of the Sea Fisheries of the South Coast of New England in 1871 and 1872" Ed by SF Baird, Government Printing Office, Washington DC, pp 295–778
- Weigert A, Helm C, Meyer M, Nickel B, Arendt D, Hausdorf B, et al. (2014) Illuminating the base of the annelid tree using transcriptomics. *Mol Biol Evol* 31: 1391–1401
- Westheide W (1967) Die Gattung *Trilobodrillus* (Archiannelida, Polychaeta) von der deutschen Nordseeküste. *Helgoländ wiss Meer* 16: 207–215
- Westheide W (1984) The concept of reproduction in polychaetes with small body size: adaptation in interstitial species. In "Polychaete Reproduction, Fortschritte der Zoologie, Vol 29" Ed by A Fischer, H-D Pfannenstiel, Gustav Fischer, New York, pp 265–287
- Westheide W (1987) Progenesis as a principle in meiofauna evolution. *J Nat Hist* 21: 843–854
- Westheide W (2008) Polychaetes: Interstitial Families. Field Studies Council, Shrewsbury
- Whiting MF, Carpenter JM, Wheeler QD, Wheeler WC (1997) The

- Strepsiptera problem: phylogeny of the holometabolous insect orders inferred from 18S and 28S ribosomal DNA sequences and morphology. *Syst Biol* 46: 1–68
- Wieser W (1957) Archannelids from the intertidal of Puget Sound. *T Am Microsc Soc* 76: 275–285
- Wiklund H, Glover AG, Dahlgren TG (2009) Three new species of *Ophryotrocha* (Annelida: Dorvilleidae) from a whale-fall in the North-East Atlantic. *Zootaxa* 2228: 43–56
- Williams DM (2011) Historical biogeography, microbial endemism and the role of classification: everything is endemic. In “Biogeography of Microscopic Organisms: Is Everything Small Everywhere?” Ed by D Fontaneto, Cambridge University Press, Cambridge, pp 11–31
- Worsaae K, Nygren A, Rouse GW, Giribet G, Persson J, Sundberg P, et al. (2005) Phylogenetic position of Nerillidae and *Aberranta* (Polychaeta, Annelida), analysed by direct optimization of combined molecular and morphological data. *Zool Scr* 34(3): 313–328
- Worsaae K, Kristensen RK (2005) Evolution of interstitial Polychaeta (Annelida). *Hydrobiologia* 535–536: 319–340
- Worsaae K, Rouse GW (2008) Is *Diurodrilus* an annelid? *J Morphol* 269: 1426–1455
- Wright LD, Short AD (1984) Morphodynamic variability of surf zones and beaches: a synthesis. *Mar Geol* 56: 93–118
- Yamaguchi S, Endo K (2003) Molecular phylogeny of Ostracoda (Crustacea) inferred from 18S ribosomal RNA sequences: implication for its origin and diversification. *Mar Biol* 143: 23–38
- Yamanishi R (1983) Preliminary report on the fauna of interstitial polychaetes (Annelida) of Japan. *Benthos Res* 24: 41–48 (in Japanese)
- Yamasaki H, Hiruta SF, Kajihara H (2013) Molecular phylogeny of kinorhynchs. *Mol Phylogenet Evol* 67: 303–310
- Zrzavý J, Říha P, Piálek L, Janouškovec J (2009) Phylogeny of Annelida (Lophotrochozoa): total-evidence analysis of morphology and six genes. *BMC Evolutionary Biology* 9: 189 doi:10.1186/1471-2148-9-189

(Received November 6, 2014 / Accepted April 7, 2015)