Title	Changes in S1P(1) and SIP(2) expression during embryonal development and primitive endoderm differentiation of F9 cells
Author(s)	Hiraga, Yuki; Kihara, Akio; Sano, Takamitsu; Igarashi, Yasuyuki
Citation	Biochemical and biophysical research communications, 344(3), 852-858 https://doi.org/10.1016/j.bbrc.2006.04.002
Issue Date	2006-06-09
Doc URL	http://hdl.handle.net/2115/11391
Type	article (author version)
File Information	Hiraga et al.pdf

[Instructions for use](#)

Changes in S1P₁ and S1P₂ expression during embryonal development and primitive endoderm differentiation of F9 cells

Yuki Hiraga, Akio Kihara*, Takamitsu Sano, and Yasuyuki Igarashi

*Department of Biomembrane and Biofunctional Chemistry, Graduate School of
Pharmaceutical Sciences, Hokkaido University, Kita 12-jo, Nishi 6-choume, Kita-ku,
Sapporo 060-0812, Japan*

*Corresponding author

Address correspondence to:

Akio Kihara

Department of Biomembrane and Biofunctional Chemistry,
Graduate School of Pharmaceutical Sciences, Hokkaido University,
Kita 12-jo, Nishi 6-choume, Kita-ku, Sapporo 060-0812, Japan

Telephone: +81-11-706-3971

Fax: +81-11-706-4986

E-mail: kihara@pharm.hokudai.ac.jp

Abstract

Sphingosine 1-phosphate (S1P) is a ligand for S1P family receptors (S1P₁- S1P₅). Of these receptors, S1P₁, S1P₂, and S1P₃ are ubiquitously expressed in adult mice, while S1P₄ and S1P₅ are tissue-specific. However, little is known of their expression during embryonal development. We performed Northern blot analyses in mouse embryonal tissue and found that such expression is developmentally regulated. We also examined the expression of these receptors during primitive endoderm (PrE) differentiation of mouse F9 embryonal carcinoma (EC) cells, a well-known *in vitro* endoderm differentiation system. *SIP*₂ mRNA was abundantly expressed in F9 EC cells, but little *SIP*₁ and no *SIP*₃, *SIP*₄, or *SIP*₅ mRNA was detectable. However, *SIP*₁ mRNA expression was induced during EC-to-PrE differentiation. Studies using small interference RNA of *SIP*₁ indicated that increased S1P₁ expression is required for PrE differentiation. Thus, S1P₁ may play an important function in PrE differentiation that is not substituted for by S1P₂.

Keywords: Sphingolipid; Sphingosine 1-phosphate; S1P receptor; S1P₁; Embryonal development; Differentiation; Endoderm; F9 cells

Introduction

As an extracellular signaling molecule, sphingosine 1-phosphate (S1P) affects several cellular processes including proliferation, migration, differentiation, actin cytoskeleton reorganization, and adherens junction assembly [1]. Through these cellular responses, S1P plays important functions in the vascular and immune systems. S1P is a ligand for S1P/Edg family receptors, which are seven-span membrane proteins coupled to heterotrimeric G proteins. To date, five members of this family have been identified in mammals, S1P₁/Edg1, S1P₂/Edg5/H218, S1P₃/Edg3, S1P₄/Edg6, and S1P₅/Edg8 [2-5]. Of the five, S1P₁ is arguably the most important, since only disruption of the *S1P₁* gene causes embryonic lethality, which is due to severe hemorrhage resulting from a defect in vascular stabilization [6]. In contrast, *S1P₂* and *S1P₃* knockout mice are viable and exhibit only mild phenotypes [7-9]. In adult mice, S1P₁, S1P₂, and S1P₃ are known to be ubiquitously expressed, whereas S1P₄ and S1P₅ have been detected only in specific tissues [4,7]. However, the expression of the receptors has not been well studied during embryonal stages.

Each S1P receptor specifically couples to certain G proteins. S1P₁ couples only with G_i proteins [10,11], whereas S1P₂ and S1P₃ couple with G_i, G_q and G_{12/13} [11]. Although S1P₂ and S1P₃ couple with the same set of G proteins, their affinities differ. S1P₂ exhibits the highest affinity toward G_{12/13}, whereas S1P₃ appears to primarily couple with G_q, then G_i [9,12]. Reflecting their distinct preferences and affinities for specific G-proteins, S1P₁, S1P₂, and S1P₃ can induce different, sometimes opposite, cellular responses. For example, S1P₁ and S1P₃ stimulation by S1P induces cell

migration, but S1P₂ stimulation inhibits it [12-14]. Thus, regulating the expression of S1P receptors to provide a cellular balance may determine the effect of S1P on cellular processes.

Differentiation during embryogenesis is accompanied by dynamic changes in protein expression that promote cell and tissue-specific development. To date, information regarding the expression of S1P receptors during embryogenesis is limited. Changes in S1P₁ expression during mouse embryonic development have been observed in the embryo proper and in two extra-embryonic tissues (the allantois and the yolk sac) [15]. S1P₁ expression patterns during mice embryogenesis have also been examined [6]. In the present study we performed Northern blot analyses in mouse embryonal tissues and found that *S1P₁*, *S1P₂*, and *S1P₃* mRNA expression is developmentally regulated. In addition, we examined the expression and regulation of these receptors using an F9 *in vitro* differentiation model. F9 embryonal carcinoma (EC) cells are a well-known system for extra-endoderm differentiation in early mouse embryogenesis. In the presence of retinoic acid (RA), F9 EC cells differentiate toward primitive endoderm (PrE) [16], and if dibutyl cyclic AMP (db-cAMP) is added together with the RA, the F9 EC cells differentiate to parietal endoderm (PE) via PrE [17]. We examined the expression profile for each S1P receptor throughout this differentiation. Little S1P₁ is expressed in F9 EC cells, but expression is induced during PrE differentiation. This expression is in direct contrast to S1P₂ expression, which is down-regulated during the EC-to-PE (via PrE) differentiation [18]. The other S1P receptors are not detectable at either stage. Thus, the expression profile of the S1P receptors is altered during

differentiation. We also report here that *SIP₁* small interference RNA (siRNA) inhibited PrE differentiation, suggesting that the induction of *SIP₁* is required for this differentiation.

Materials and methods

Cell culture and transfection. Mouse F9 cells were grown on 0.1% gelatin-coated dishes in Dulbecco's modified Eagle's medium (Sigma, St. Louis, MO; D6429) containing 10% fetal bovine serum and supplemented with 100 units/ml penicillin and 100 µg/ml streptomycin. For differentiation experiments, 1 µM all-*trans*-RA (Sigma) with or without 250 µM bt_2cAMP (Sigma) was added to the medium. Transfections were performed using Lipofectamine™ 2000 reagent (Invitrogen, Carlsbad, CA).

Plasmids. The pCE-puro plasmid, a mammalian expression vector, contains a puromycin-resistant gene used for selecting stable transformants [19]. The pCE-puro 3xFLAG-4 plasmid is a derivative of the pCE-puro plasmid and was designed to produce a C-terminal triple FLAG (3xFLAG)-tagged protein. The pCE-puro SIP_1 -3xFLAG and pCE-puro SIP_2 -3xFLAG plasmids encoding SIP_1 -3xFLAG and SIP_2 -3xFLAG, respectively, were constructed by cloning SIP_1 or SIP_2 into the pCE-puro 3xFLAG-4 plasmid.

Preparing stable transformants of F9 cells. F9 derivatives that stably express SIP_1 -3xFLAG (F9-18 and F9-19) or SIP_2 -3xFLAG (F9-14 and F9-15) were prepared by transfection of the pCE-puro SIP_1 -3xFLAG or pCE-puro SIP_2 -3xFLAG plasmid, respectively, into F9 cells and by subsequent puromycin selection at 0.5 µg/ml for 1 week. F9-17 cells are control cells that were generated by transfection of pCE-puro 3xFLAG-4 vector and subsequent puromycin-selection.

Northern blot analysis and reverse transcription (RT)-PCR. To examine the embryonal development-specific expression of *SIP*₁, *SIP*₂, and *SIP*₃ mRNA, blots containing 20 µg total RNA from mouse embryos at several stages of development were purchased from Seegene (Seoul, Korea). To likewise examine mRNA expression in F9 cells, blots were prepared using total RNA isolated from F9 EC, PrE, and PE cells with Trizol Reagent (Invitrogen) as instructed by the manufacturer. Fixed amounts of total RNA (20 µg) were separated on a 1% agarose/2.2 M formaldehyde gel and transferred to positively charged nylon membranes (Roche Diagnostics, Indianapolis, IN).

³²P-labeled probes were prepared from cDNA for each SIP receptor, which had been amplified using specific primers (for SIP₁,

5'-ATGGTGTCCACTAGCATCCCGG-3' and

5'-TTAGGAAGAAGAATTGACGTTTCCAGACG-3'; for SIP₂,

5'-CCACCATGGGCGGCTTATACTCAGAGTACC-3' and

5'-TCAGACCACTGTGTTACCCTCC-3'; for SIP₃,

5'-AAGCCATGGCAACCACGCATGCGCAGG-3' and

5'-TCACTTGCAGAGGACCCCGTTCTG-3'; for SIP₄,

5'-GGCCATGAACATCAGTACCTGG-3' and

5'-CTAGGTGCTGCGGACGCTGGAAATGC-3'; and for SIP₅,

5'-GTGCTGGTCCGCCACCCTCGCTTCC-3' and

5'-TCAGTCTGTAGCAGTAGGCACCAGG-3') then labeled with [³²P] using the

TAKARA Bio (Shiga, Japan) random primer DNA labeling kit, version 2.

Hybridization of the probes to the mRNA blots was carried out in ExpressHyb Hybridization Solution (BD Biosciences Clontech, Palo Alto, CA) at 68°C for 2 h. Radioactivities were quantified using a Bio-Imaging Analyzer BAS2500 (Fuji Photo Film, Tokyo, Japan).

RT-PCR was performed using total RNAs prepared from untreated or differentiated F9 cells as templates and SuperScript™ One-Step RT-PCR System with Platinum Taq (Invitrogen). The primers used included for *SIP₁*, 5'-ATGGTGTCCACTAGCATCCCGG-3' and 5'-TTAGGAAGAAGAATTGACGTTTCCAGACG-3'; for *SIP₂*, 5'-CCACCATGGGCGGCTTATACTCAGAGTACC-3' and 5'-TCAGACCACTGTGTTACCCTCC-3'; for *actin*, 5'-ATGGATGACGATATCGCTGCGCTGG-3' and 5'-CTAGAAGCACTTGCGGTGCACGATG-3'; for *GATA-4*, 5'-ATGTACCAAAGCCTGGCCATGGCCGCCAACC-3' and 5'-TTACGCGGTGATTATGTCCCCATGACTGTC-3'; and for *GATA-6*, 5'-ATGTACCAGACCCTCGCCGCCCTGTCCAGC-3' and 5'-TCAGACCAGGGCCAGAGCACACCAAGAATCC-3'.

Phalloidin staining. Phalloidin staining of F-actin was performed on F9 EC, PrE, and PE cells as described previously using a fluorescence microscopy AxioSkop 2 plus (Carl Zeiss, Oberkochen, Germany) [19].

Immunoblotting. Cells were washed twice with PBS, suspended in buffer A (62.5 mM Tris-HCl (pH 6.8), 2% SDS, and 10% glycerol), and sonicated. After removal of cell debris by centrifugation, protein concentrations of the resulting lysates were quantified using BCA protein assay kit (Pierce Biotechnology, Rockford, IL). Samples were mixed with 1X SDS sample buffer (buffer A containing a trace amount of bromophenol blue and 10% 2-mercaptoethanol) and boiled at 100°C for 3 min. Proteins were separated by SDS-PAGE and transferred to Immobilon™ polyvinylidene difluoride membrane (Millipore, Billerica, MA). After it was blocked with 5% skim milk in TBST (Tris-buffered-saline with 0.05% Tween 20), the membrane was incubated for 1 h with an anti-Dab2 (BD Biosciences Clontech) or anti-actin (Sigma) antibody, each diluted 1/1000. The membranes were washed three times with TBST, incubated for 1 h with an HRP-conjugated anti-mouse IgG F(ab')₂ fragment (1/7500 dilution; Amersham Biosciences, Piscataway, NJ), then washed three times with TBST three times and once with TBS. Labeling was detected by the ECL detection method (Amersham Biosciences).

RNA interference. The siRNAs were purchased from Greiner Bio-One (Kremsmünster, Austria). The nucleotide sequences of the respective RNAs targeted by the siRNAs are as follows: *SIP₁* siRNA, 5'-AACTGACTTCAGTGGTGTTC-3' (nucleotides 137-157 of mouse *SIP₁* ORF); *SIP₂* siRNA, 5'-AAGGTCAAGCTCTACGGCAGT-3' (nucleotides 406-426 of mouse *SIP₂* ORF); and scrambled *SIP₁* siRNA, 5'-AATTGATGATCGTTCCGGACT-3'. F9 cells were

transfected with the appropriate siRNA together with the pCE-puro vector, which carries a puromycin resistant gene. Sixteen h after transfection on day 0, 1 μ M RA and 0.5 μ g/ml puromycin were added to the culture medium to initiate differentiation and to kill the untransfected cells, respectively. Medium was changed every 2 days, and puromycin was used for 48 h. Cells were recovered on days 0 through 4, and total cell lysates were prepared. Dab2 and actin were detected in the cell lysates by immunoblotting. Total RNA was prepared from cells on day 1 (40 h after transfection), and subjected to RT-PCR as described above.

Results

Regulation of SIP_1 , SIP_2 , and SIP_3 expression during embryonal development

SIP_1 , SIP_2 , and SIP_3 are expressed ubiquitously among tissues in adult mice [7]. However, their developmental stage-specific expression patterns have not been studied in depth. Therefore, we performed Northern blot analyses using a mouse embryo full stage blot. As shown in Fig. 1, SIP_1 mRNA is abundantly expressed throughout mouse embryogenesis with a transient peak around embryonic day (E) 8.5. The expression of SIP_2 mRNA is high in earlier developmental stages but low in later stages. The highest SIP_2 mRNA expression was observed at E7.5; the expression gradually decreased toward E18.5. Conversely, the SIP_3 mRNA is low in earlier developmental stages (E4.5 to E9.5) but rapidly increase at E10.5, reaching a maximal level at E11.5. The expression then gradually decreases. These results indicate that SIP_1 , SIP_2 , and SIP_3 each exhibits a characteristic, developmental stage-specific expression pattern.

Induction of SIP_1 during PrE differentiation of F9 cells

Having determined that the expression of SIP receptors changes during development, we set out to explore the molecular mechanisms responsible. However, a whole animal model is not suitable for such analysis. Therefore, we used F9 cells, a well-known model system for extra-endoderm differentiation in early mouse embryogenesis. Reportedly, SIP_2 mRNA is gradually down-regulated during the EC-to-PE (via PrE) differentiation of F9 cells [18]. However, in that study expression of the other SIP receptors was not examined. Therefore, we investigated the expression

levels of all S1P receptors in F9 cells at each differentiation stage. Treatment with 1 μ M RA for 3-4 days induced F9 EC cells to differentiate into PrE cells, so that the cells became enlarged, and numerous, F-actin-dense projections appeared (Fig. 2). Combined treatment with 1 μ M RA and 250 μ M bt₂cAMP induced further differentiation of PrE cells into PE cells. PE cells were readily distinguished from PrE cells by their morphology, which is characterized as round shapes with long cell processes, usually two (Fig. 2). Most of the EC cells had differentiated to PE cells by 6 days post-treatment.

We prepared total RNA from F9 cells treated with RA alone or with RA/bt₂cAMP for 0, 3, or 6 days, and performed Northern blot analyses using specific probes for each S1P receptor. Untreated, F9 EC cells expressed a significant amount of *SIP*₂ mRNA (Fig. 3). As reported [18], the *SIP*₂ mRNA levels decreased by day 3 of treatment with RA/bt₂cAMP, and were further decreased at day 6 (Fig. 3). Treatment with RA alone did not cause a reduction in *SIP*₂ at day 3, but the levels were slightly decreased by day 6. In contrast, *SIP*₁ mRNA was barely detected in F9 EC cells, however its expression was induced by a 3 day treatment with RA or with RA/bt₂cAMP (Fig. 3). Overall, the *SIP*₁ levels were higher in the presence of bt₂cAMP than in its absence, but the expression decreased in cells treated for 6 days with either treatment (Fig. 3). These results indicate that *SIP*₁ mRNA is induced during EC-to-PrE differentiation. No *SIP*₃, *SIP*₄, or *SIP*₅ mRNA was detected at any differentiation stage (Fig. 3).

We further investigated the *SIP*₁ mRNA induction over time. When F9 EC cells were treated with RA alone, the *SIP*₁ mRNA levels gradually increased, reaching

maximum at day 3 (Fig. 4A). Maximum expression was maintained through day 5, but the levels slightly decreased at day 6. Thus, SIP_1 was induced during the EC-to-PrE differentiation. Treatment with RA/bt₂cAMP induced more SIP_1 mRNA expression than treatment with RA alone did, however, after reaching maximum level at day 3, the SIP_1 levels gradually decreased (Fig. 4A). These results suggest that in the course of successive differentiation from EC-to-PrE-to-PE, SIP_1 mRNA transiently peaks at the PrE stage.

Consistent with the previous report [18], during treatment with RA/bt₂cAMP SIP_2 mRNA levels slightly increased at day 1 then gradually decreased, reaching minimal levels at day 4 (Fig. 4B). In contrast, RA treatment caused a large increase at day 1, followed by only a slight decrease. At days 3 and 4, around which PrE differentiation was complete, the SIP_2 levels were equivalent to those observed in the untreated EC cells. These results indicate that SIP_2 levels are high at the EC and PrE stages but are nearly absent at the PE stage. Thus, S1P may function only through S1P₁ at the PE stage in contrast to the EC stage, at which S1P₂ is predominant.

Requirement of SIP_1 in the PrE differentiation

To investigate whether S1P₁ or S1P₂ is involved in the differentiation of EC cells into PrE cells, we prepared siRNAs specific to each of their sequences, as well as SIP_1 scrambled siRNA. RT-PCR using total RNAs prepared from F9 cells transfected with the SIP_1 , SIP_2 , or the scrambled siRNA confirmed a reduction in the SIP_1 and SIP_2 mRNAs by their specific siRNAs (Fig. 5A). F9 cells were transfected with each of these

siRNAs and treated with RA to induce PrE differentiation. Differentiation to PrE was monitored by investigating the expression of the PrE marker Dab2 [20]. As shown in Fig. 5B, F9 cells transfected with the scrambled siRNA expressed two spliced isoforms of Dab2, p96 and p67, around days 3 and 4 of treatment, much like the expression in untransfected F9 cells (data not shown). Similar bands were also observed in cells transfected with the *SIP*₂ siRNA, indicating that it had no effect. However, the *SIP*₁ siRNA significantly inhibited the induction of Dab2 expression. Further RT-PCR analysis revealed that similar induction of two other PrE markers, the transcription factors GATA-4 [21] and GATA-6 [22], was also reduced by the *SIP*₁ siRNA but not by the *SIP*₂ siRNA (Fig. 5C). These results suggest that the induction of *SIP*₁ is required for the progression of PrE differentiation.

Overexpression of SIP₁ has no effect on PrE differentiation of F9 cells

We next examined whether overexpression of *SIP*₁ in F9 EC cells could stimulate PrE differentiation. Two independent F9 cells stably expressing *SIP*₁-3xFLAG or *SIP*₂-3xFLAG were incubated with RA/bt₂cAMP, and the expression of Dab2 was examined. As shown in Fig. 6, neither overexpression of *SIP*₁-3xFLAG nor that of *SIP*₂-3xFLAG stimulated PrE differentiation. Thus, *SIP*₁ expression is necessary but not sufficient for the PrE differentiation.

Discussion

In the present study, we have demonstrated that *SIP₁*, *SIP₂*, and *SIP₃* mRNA each exhibit a characteristic, developmental stage-specific expression pattern. *SIP₁* mRNA transiently increases around E8.5, soon after the beginning of vasculogenesis.

Vasculogenesis is initiated by the birth of angioblasts (endothelial cell precursors), followed by angioblast aggregation in extraembryonic regions. High levels *SIP₁* expression have been observed in dorsal aorta, intersomitic arteries, and capillaries at E9.5 [6]. Thus, the high expression of *SIP₁* at E8.5 and E9.5 may be related to the active vasculogenesis.

Although *SIP₁* is the most important *SIP* receptor in the development of a stable and mature vascular system, *SIP₂* and *SIP₃* also function coordinately. *SIP₁*-null mice appeared normal up to E11.5, but bleeding was evident at E12.5 [6]. Wide-spread hemorrhage and severe edema were observed at E13.5, and the mice could not survive beyond E14.5 [6]. Although both *SIP₂*- and *SIP₃*-null mice developed normally vascular systems, *SIP₁ SIP₂* double knockout mice and *SIP₁ SIP₂ SIP₃* triple knockout mice exhibited more severe vascular defects than those of the *SIP₁* single knockout mice [23]. About half of the triple null mice exhibited bleeding at E10.5, and most of those died at E11.5. This timing coincides with a rapid induction of *SIP₃* at E10.5 (Fig. 1), suggesting that *SIP₃* has some function in the development of vascular system around these embryonic stages.

We have also demonstrated here that the expression patterns of *SIP₁* and *SIP₂* mRNAs change during the differentiation of F9 cells, with EC expressing little *SIP₁* and

high levels of *SIP*₂; PrE, high levels of *SIP*₁ and high levels of *SIP*₂; and PE, intermediate levels of *SIP*₁ and little *SIP*₂ (Figs 3 and 4). In addition, analysis using siRNA revealed that *SIP*₁ expression is required for PrE differentiation (Fig. 5). Considering that PrE cells contain a high amount of *SIP*₂, *SIP*₁ might have an important function in PrE differentiation that is not substituted for by *SIP*₂. Such differences in function are well established in regards to cell migration [12-14].

Differentiation is a complicated process, in which many transcriptional factors, and thus many genes, are temporally regulated. During F9 differentiation, the most immediate gene response is likely mediated by RA receptors bound to RA. Two classes of such receptors are known, retinoic acid receptors (RARs) and retinoid X receptors (RXRs), each of which has three isotypes (α , β , and γ). These receptors form homodimers or heterodimers and activate the transcription of genes containing the retinoic acid response element (RARE). The RXR α /RAR γ heterodimer has been shown to be important for the PrE and PE differentiation of F9 cells [24].

Genes exhibiting changes in their levels during RA-mediated PrE differentiation are classified mainly into two groups, those whose expression changes within the first 24 h and those peaking around day 3 [25,26]. The first group includes genes directly activated by RA, but also those activated indirectly. Several transcriptional factors, including AP-1, GATA-4, and GATA-6, are induced around day 3 [21,27-29] and may be involved in the expression of genes in the second group. We found that *SIP*₁ mRNA expression reaches maximum at day 3, whereas *SIP*₂ mRNA transiently increases on day 1. These results suggest that *SIP*₁ and *SIP*₂ belong to the second and first groups,

respectively, although the transcription factors involved in their increases are presently unknown. Using the transcription factor database TRANSFAC and accompanying programs (<http://transfac.gbf.de/TRANSFAC>), we found two putative AP-1 binding sites in the *S1P₁* promoter region. Thus, it is possible that AP-1 is responsible for the induction of *S1P₁* during EC-to-PrE differentiation.

S1P is abundant in blood [30] and, therefore, in extracellular spaces. However, *S1P* also exists intracellularly in most cells, although the levels are quite low. Intracellular *S1P* is proposed to function as a second messenger of various stimuli such as growth factors and cytokines [31]. We recently reported that sphingosine kinase and *S1P* lyase, both of which are involved in *S1P* metabolism, were up-regulated during F9 PrE differentiation [19]. Moreover, *S1P* accumulation, resulting from the disruption of the *S1P* lyase gene or the overproduction of sphingosine kinase, resulted in accelerated of the PrE differentiation [19], suggesting that intracellular *S1P* is also involved in this process. Thus, it appears that extracellular and intracellular *S1P* cooperatively regulate PrE differentiation.

Expression of *S1P₁* is known to be altered during several types of development and differentiation stages. For example, in human endothelial cells the expression of *S1P₁* was rapidly increased by treatment with phorbol 12-myristate 13-acetate, which induces differentiation of these cells into capillary-like, tubular structures [32]. Changes in *S1P₁* expression during embryonic development has also been shown in the embryo proper and in two extra-embryonic tissues (the allantois and the yolk sac) [15].

Treatment of endothelial cells with vascular endothelial growth factor (VEGF) rapidly

induces expression of S1P₁, which is attenuated by the tyrosine kinase inhibitor genistein and by the protein kinase C inhibitor calphostin C [33]. Moreover, S1P₁ levels change during T cell maturation and activation. In the course of T cell maturation in the thymus, S1P₁ is strongly upregulated [34,35]. Later, when mature naïve T cells encounter their specific antigens in the lymph node, S1P₁ is downregulated to prevent S1P gradient-induced exit from the lymph node [34]. After activation and clonal expansion, T cells restore their S1P₁ levels and exit [34]. The molecular mechanism regulating such S1P₁ expression is unclear, however, and further analysis is needed. For this purpose, the well-examined F9 system will be useful.

Acknowledgments

This work was supported by a Grant-in-Aid for Scientific Research on Priority Areas (B) (12140201) from the Ministry of Education, Culture, Sports, Science and Technology of Japan.

References

- [1] T.A. Taha, K.M. Argraves, L.M. Obeid, Sphingosine-1-phosphate receptors: receptor specificity versus functional redundancy, *Biochim. Biophys. Acta.* 1682 (2004) 48-55.
- [2] M.J. Lee, J.R. Van Brocklyn, S. Thangada, C.H. Liu, A.R. Hand, R. Menzeleev, S. Spiegel, T. Hla, Sphingosine-1-phosphate as a ligand for the G protein-coupled receptor EDG-1, *Science* 279 (1998) 1552-1555.
- [3] S. An, T. Bleu, W. Huang, O.G. Hallmark, S.R. Coughlin, E.J. Goetzl, Identification of cDNAs encoding two G protein-coupled receptors for lysosphingolipids, *FEBS Lett.* 417 (1997) 279-282.
- [4] M.H. Gräler, G. Bernhardt, M. Lipp, EDG6, a novel G-protein-coupled receptor related to receptors for bioactive lysophospholipids, is specifically expressed in lymphoid tissue, *Genomics* 53 (1998) 164-169.
- [5] R.L. Malek, R.E. Toman, L.C. Edsall, S. Wong, J. Chiu, C.A. Letterle, J.R. Van Brocklyn, S. Milstien, S. Spiegel, N.H. Lee, Nrg-1 belongs to the endothelial differentiation gene family of G protein-coupled sphingosine-1-phosphate receptors, *J. Biol. Chem.* 276 (2001) 5692-5699.
- [6] Y. Liu, R. Wada, T. Yamashita, Y. Mi, C.X. Deng, J.P. Hobson, H.M. Rosenfeldt, V.E. Nava, S.S. Chae, M.J. Lee, C.H. Liu, T. Hla, S. Spiegel, R.L. Proia, Edg-1, the G protein-coupled receptor for sphingosine-1-phosphate, is essential for vascular maturation, *J. Clin. Invest.* 106 (2000) 951-961.
- [7] I. Ishii, B. Friedman, X. Ye, S. Kawamura, C. McGiffert, J.J. Contos, M.A.

- Kingsbury, G. Zhang, J.H. Brown, J. Chun, Selective loss of sphingosine 1-phosphate signaling with no obvious phenotypic abnormality in mice lacking its G protein-coupled receptor, LP_{B3}/EDG-3, J. Biol. Chem. 276 (2001) 33697-33704.
- [8] A.J. MacLennan, P.R. Carney, W.J. Zhu, A.H. Chaves, J. Garcia, J.R. Grimes, K.J. Anderson, S.N. Roper, N. Lee, An essential role for the H218/AGR16/Edg-5/LP_{B2} sphingosine 1-phosphate receptor in neuronal excitability, Eur. J. Neurosci. 14 (2001) 203-209.
- [9] I. Ishii, X. Ye, B. Friedman, S. Kawamura, J.J. Contos, M.A. Kingsbury, A.H. Yang, G. Zhang, J.H. Brown, J. Chun, Marked perinatal lethality and cellular signaling deficits in mice null for the two sphingosine 1-phosphate (S1P) receptors, S1P₂/LP_{B2}/EDG-5 and S1P₃/LP_{B3}/EDG-3, J. Biol. Chem. 277 (2002) 25152-25159.
- [10] M.J. Lee, M. Evans, T. Hla, The inducible G protein-coupled receptor *edg-1* signals via the G_i/mitogen-activated protein kinase pathway, J. Biol. Chem. 271 (1996) 11272-11279.
- [11] R.T. Windh, M.J. Lee, T. Hla, S. An, A.J. Barr, D.R. Manning, Differential coupling of the sphingosine 1-phosphate receptors Edg-1, Edg-3, and H218/Edg-5 to the G_i, G_q, and G₁₂ families of heterotrimeric G proteins, J. Biol. Chem. 274 (1999) 27351-27358.
- [12] H. Okamoto, N. Takuwa, T. Yokomizo, N. Sugimoto, S. Sakurada, H. Shigematsu, Y. Takuwa, Inhibitory regulation of Rac activation, membrane ruffling, and cell migration by the G protein-coupled sphingosine-1-phosphate receptor EDG5 but not EDG1 or EDG3, Mol. Cell. Biol. 20 (2000) 9247-9261.

- [13] F. Wang, J.R. Van Brocklyn, J.P. Hobson, S. Movafagh, Z. Zukowska-Grojec, S. Milstien, S. Spiegel, Sphingosine 1-phosphate stimulates cell migration through a G_i -coupled cell surface receptor. Potential involvement in angiogenesis, *J. Biol. Chem.* 274 (1999) 35343-35350.
- [14] J. Kon, K. Sato, T. Watanabe, H. Tomura, A. Kuwabara, T. Kimura, K. Tamama, T. Ishizuka, N. Murata, T. Kanda, I. Kobayashi, H. Ohta, M. Ui, F. Okajima, Comparison of intrinsic activities of the putative sphingosine 1-phosphate receptor subtypes to regulate several signaling pathways in their cDNA-transfected Chinese hamster ovary cells, *J. Biol. Chem.* 274 (1999) 23940-23947.
- [15] K.M. Argraves, B.A. Wilkerson, W.S. Argraves, P.A. Fleming, L.M. Obeid, C.J. Drake, Sphingosine-1-phosphate signaling promotes critical migratory events in vasculogenesis, *J. Biol. Chem.* 279 (2004) 50580-50590.
- [16] S. Strickland, V. Mahdavi, The induction of differentiation in teratocarcinoma stem cells by retinoic acid, *Cell* 15 (1978) 393-403.
- [17] S. Strickland, K.K. Smith, K.R. Marotti, Hormonal induction of differentiation in teratocarcinoma stem cells: generation of parietal endoderm by retinoic acid and dibutyryl cAMP, *Cell* 21 (1980) 347-355.
- [18] Y. Li, A.J. MacLennan, M.B. Rogers, A putative G-protein-coupled receptor, H218, is down-regulated during the retinoic acid-induced differentiation of F9 embryonal carcinoma cells, *Exp. Cell Res.* 239 (1998) 320-325.
- [19] A. Kihara, M. Ikeda, Y. Kariya, E.Y. Lee, Y.M. Lee, Y. Igarashi, Sphingosine-1-phosphate lyase is involved in the differentiation of F9 embryonal

- carcinoma cells to primitive endoderm, *J. Biol. Chem.* 278 (2003) 14578-14585.
- [20] E.R. Smith, C.D. Capo-chichi, J. He, J.L. Smedberg, D.H. Yang, A.H. Prowse, A.K. Godwin, T.C. Hamilton, X.X. Xu, Disabled-2 mediates c-Fos suppression and the cell growth regulatory activity of retinoic acid in embryonic carcinoma cells, *J. Biol. Chem.* 276 (2001) 47303-47310.
- [21] R.J. Arceci, A.A. King, M.C. Simon, S.H. Orkin, D.B. Wilson, Mouse GATA-4: a retinoic acid-inducible GATA-binding transcription factor expressed in endodermally derived tissues and heart, *Mol. Cell. Biol.* 13 (1993) 2235-2246.
- [22] S. Futaki, Y. Hayashi, T. Emoto, C.N. Weber, K. Sekiguchi, Sox7 plays crucial roles in parietal endoderm differentiation in F9 embryonal carcinoma cells through regulating Gata-4 and Gata-6 expression, *Mol. Cell. Biol.* 24 (2004) 10492-10503.
- [23] M. Kono, Y. Mi, Y. Liu, T. Sasaki, M.L. Allende, Y.P. Wu, T. Yamashita, R.L. Proia, The sphingosine-1-phosphate receptors S1P₁, S1P₂, and S1P₃ function coordinately during embryonic angiogenesis, *J. Biol. Chem.* 279 (2004) 29367-29373.
- [24] H. Chiba, J. Clifford, D. Metzger, P. Chambon, Specific and redundant functions of retinoid X Receptor/Retinoic acid receptor heterodimers in differentiation, proliferation, and apoptosis of F9 embryonal carcinoma cells, *J. Cell Biol.* 139 (1997) 735-747.
- [25] M.J. Sleight, Differentiation and proliferation in mouse embryonal carcinoma cells, *Bioessays* 14 (1992) 769-775.
- [26] T.M. Harris, G. Childs, Global gene expression patterns during differentiation of

F9 embryonal carcinoma cells into parietal endoderm, *Funct. Integr. Genomics* 2 (2002) 105-119.

- [27] J.L. Bjersing, A. Brorsson, O. Heby, Increased expression of *c-jun*, but not retinoic acid receptor β , is associated with F9 teratocarcinoma stem cell differentiation induced by polyamine depletion, *J. Cell. Biochem.* 67 (1997) 378-385.
- [28] S. Futaki, Y. Hayashi, M. Yamashita, K. Yagi, H. Bono, Y. Hayashizaki, Y. Okazaki, K. Sekiguchi, Molecular basis of constitutive production of basement membrane components. Gene expression profiles of Engelbreth-Holm-Swarm tumor and F9 embryonal carcinoma cells, *J. Biol. Chem.* 278 (2003) 50691-50701.
- [29] A. Murakami, J. Thurlow, C. Dickson, Retinoic acid-regulated expression of fibroblast growth factor 3 requires the interaction between a novel transcription factor and GATA-4, *J. Biol. Chem.* 274 (1999) 17242-17248.
- [30] Y. Yatomi, Y. Igarashi, L. Yang, N. Hisano, R. Qi, N. Asazuma, K. Satoh, Y. Ozaki, S. Kume, Sphingosine 1-phosphate, a bioactive sphingolipid abundantly stored in platelets, is a normal constituent of human plasma and serum, *J. Biochem.* 121 (1997) 969-973.
- [31] S. Pyne, N.J. Pyne, Sphingosine 1-phosphate signalling in mammalian cells, *Biochem. J.* 349 (2000) 385-402.
- [32] T. Hla, T. Maciag, An abundant transcript induced in differentiating human endothelial cells encodes a polypeptide with structural similarities to G-protein-coupled receptors, *J. Biol. Chem.* 265 (1990) 9308-9313.

- [33] J. Igarashi, P.A. Erwin, A.P. Dantas, H. Chen, T. Michel, VEGF induces S1P₁ receptors in endothelial cells: Implications for cross-talk between sphingolipid and growth factor receptors, *Proc. Natl. Acad. Sci. USA* 100 (2003) 10664-10669.
- [34] M. Matloubian, C.G. Lo, G. Cinamon, M.J. Lesneski, Y. Xu, V. Brinkmann, M.L. Allende, R.L. Proia, J.G. Cyster, Lymphocyte egress from thymus and peripheral lymphoid organs is dependent on S1P receptor 1, *Nature* 427 (2004) 355-360.
- [35] M.L. Allende, J.L. Dreier, S. Mandala, R.L. Proia, Expression of the sphingosine 1-phosphate receptor, S1P₁, on T-cells controls thymic emigration, *J. Biol. Chem.* 279 (2004) 15396-15401.

Figure legends

Fig. 1. Expression of *SIP*₁, *SIP*₂, and *SIP*₃ mRNA during mouse embryonal development. (A) A ³²P-labeled *SIP*₁, *SIP*₂, or *SIP*₃ probe was hybridized to 20 µg of total RNA from each stage of a mouse embryo (upper three panels). To demonstrate uniform RNA loading, 28S and 18S ribosomal RNAs were stained with ethidium bromide (bottom panel). Embryo samples from E4.5 to E6.5 included extra-embryonic tissues and maternal uterus, while the samples from E7.5 to E9.5 were conceptuses, including embryo and extraembryonic tissues. The samples from E10.5 to E18.5 were solely embryos. (B) Radioactivity associated with each SIP receptor in (A) was quantified using a BAS-2500 bioimaging analyzer and expressed as a percentage relative to the value at E4.5.

Fig. 2. Differentiation of F9 EC cells to PrE and PE cells. F9 EC cells are differentiated to PrE cells by treatment with 1 µM RA over 3 days. Co-incubation with 1 µM RA and 250 µM bt₂cAMP causes differentiation of F9 EC cells to PrE around day 3 and further differentiation to PE at days 5 to 6. All cells were fixed, permeabilized, stained with phalloidin to visualize F-actin, and observed under a fluorescence microscope.

Fig. 3. Only *SIP*₁ and *SIP*₂ are expressed and regulated during F9 differentiation. Total RNA was prepared from F9 cells incubated with 1 µM RA or with 1 µM RA/250 µM bt₂cAMP for the indicated times. Fixed amounts of RNA (20 µg) were separated by agarose gel electrophoresis and subjected to Northern blotting using a ³²P-labeled probe

specific for the indicated *SIP* receptor or for *actin* to demonstrate uniform RNA loading.

Fig. 4. Kinetics of *SIP*₁ and *SIP*₂ expression during F9 differentiation. F9 cells were incubated with 1 μ M RA or with 1 μ M RA/250 μ M bt₂cAMP for the indicated times. Total RNA was prepared from each culture and equivalent amounts (20 μ g) were subjected to Northern blotting using a ³²P-labeled probe for *SIP*₁ (A) or *SIP*₂ (B). Radioactivities associated with the *SIP*₁ and *SIP*₂ mRNAs were quantified using a Bio-Imaging Analyzer BAS2500. Values are illustrated relative to the amount of the *SIP*₁ mRNA at day 1 or the *SIP*₂ mRNA at day 0, and represent the mean $\pm$ SD from three independent experiments.

Fig. 5. Involvement of *SIP*₁ in PrE differentiation. F9 cells were transfected with *SIP*₁ siRNA, *SIP*₂ siRNA, or *SIP*₁ scrambled (sc) siRNA and incubated for 16 h. PrE differentiation was then initiated by the addition of 1 μ M RA. (A) Total RNA was prepared from F9 cells at day 1 following RA treatment and subjected to RT-PCR analysis using primers specific for *SIP*₁, *SIP*₂, or *actin*. (B) At the indicated times following RA treatment, total lysates were prepared. Proteins (10 μ g) were separated by SDS-PAGE, followed by immunoblotting with an anti-Dab2 antibody, or, to demonstrate uniform protein loading, an anti-actin antibody. (C) Total RNA was prepared at the indicated times following RA treatment and was subjected to RT-PCR analysis using primers specific for *GATA-4*, *GATA-6*, or *actin*.

Fig. 6. Forced expression of S1P₁ or S1P₂ has no effect on PrE differentiation. F9 cells stably expressing vector only (F9-17), S1P₁-3xFLAG (F9-18 and F9-19), or S1P₂-3xFLAG (F9-14 and F9-15) were incubated with 1 μ M RA and 250 μ M bt₂cAMP for the indicated times. Total proteins (15 μ g) were separated by SDS-PAGE, followed by immunoblotting with an anti-Dab2 or anti-actin antibody.