



Title	Cytopathic Changes in Viroid-infected Leaf Tissues
Author(s)	KOJIMA, Makoto; MURAI, Madoka; SHIKATA, Eishiro
Citation	Journal of the Faculty of Agriculture, Hokkaido University, 61(2), 219-223
Issue Date	1983-03
Doc URL	http://hdl.handle.net/2115/12983
Type	bulletin (article)
File Information	61(2)_p219-223.pdf



[Instructions for use](#)

CYTOPATHIC CHANGES IN VIROID-INFECTED LEAF TISSUES¹⁾

Makoto KOJIMA

(Laboratory of Plant Pathology, Faculty of Agriculture,
Niigata University, Niigata, Japan)

Madoka MURAI and Eishiro SHIKATA

(Department of Botany, Faculty of Agriculture,
Hokkaido University, Sapporo, Japan)

Received November 11, 1982

Introduction

Eight different viroids have been described so far as a causal agent of plant diseases since potato spindle tuber disease was confirmed to be caused by a viroid, a novel pathogen¹⁾. Recently, the ultrastructural changes caused by viroid infections have been studied by several workers^{2,3,6,7,8,11,13,14)} but their results are not always coincident at present. SEMANCIK and VANDERWOUDE¹⁰⁾ first pointed out that citrus exocortis viroid(CEV) induced abundant production of paramural bodies (plasmalemmasomes), cytopathic effects at the plasma membrane in *Gynura aurantiaca*. In contrast to this finding, using the same system, WAHN *et al.*^{13,14)} reported that the actual cytopathic changes were aberrations of the cell walls and of the plasmalemmasomes themselves. HARI⁹⁾ also reported on ultrastructural changes of tomato leaf tissues infected with potato spindle tuber viroid(PSTV) showing production of paramural bodies and aberrations of the thylakoid membrane systems of the chloroplasts.

To determine which cytopathic effects occurred in each host-viroid combination, we conducted some comparative experiments using cucumber plants infected with hop stunt viroid(HSV) and cucumber pale fruit viroid(CPFV) and tomato plants infected with PSTV, CEV, HSV and CPFV, respectively.

[J. Fac. Agr. Hokkaido Univ., Vol. 61, Pt. 2, 1983]

1) This work was supported in part by a Grant-in-Aid from Ministry of Education, Science and Culture, Japan (No. 56360004).

Materials and Methods

Viroids and plants

Viroids used in this experiment were HSV¹⁰, CPFV* (from Dr. SÄNGER, Max-Planck-Institut, W. GERMANY), PSTV* (from Dr. SÄNGER) and CEV* (from Dr. SÄNGER). They were propagated in tomato (*Lycopersicon esculentum* cv. Rentita) and cucumber (*Cucumis sativus* cv. Suyu) plants. Inoculated plants were grown in a greenhouse at 28 C¹⁰.

Electron microscopy

Apical leaves of each plant were cut off 5 weeks after inoculation. Fixation and embedding procedures were the same as those described in a previous report¹². Ultrathin sections were cut by an ultramicrotome (Reihelt-Jung OmU4) equipped with a diamond knife. Specimens were viewed by an electron microscope (JEM 100 B, 80 kV) after double staining by uranyl acetate and lead citrate.

Results and Discussion

Cytopathic changes in cucumber plants infected with HSV and CPFV

Symptoms caused by HSV were quite similar to those caused by CPFV in cucumber plants. Each cucumber plant infected with HSV and CPFV showed very severe symptoms such as yellowing and shortening of internodes⁹. Table 1 summarized the cytopathic changes caused by both viroids in the infected cucumber plants and indicated that their internal symptoms were almost identical. One of the typical alterations is aberration of cell

TABLE 1. Cytopathic changes in cucumber and tomato plants infected by viroids

Host	Cucumber (Suyo)			Tomato (Rentita)					
	Viroid	HSV	CPFV	control†	HSV	CPFV	PSTV	CEV	control†
External symptom		+	+	-	-	-	+	+	-
Papamural body		+	+	+	+	+	+	+	+
Aberration of cell wall		+	+	-	-	-	+	+	-
Chloroplast disintegration		+	+	-	+	+	+	+	-
Disappearance of tonoplast		+	+	-	+	+	+	+	-

† healthy plants

* CPFV, PSTV, CEV, were imported with Quarantine permission No. 55 Yokoshoku Dai 1035 gou.

walls in all tissues, such as epidermis, palisade parenchyma, spongy parenchyma and vascular bundles. As shown in Plate I, the cell walls of HSV and CPFV-infected cucumber leaves are extremely distorted. These areas were characterized by cell walls with corrugated profile and irregular thickness. These aberrations appear to lead to a structural deformation of the cells themselves, resulting in external symptoms. Another alteration in the infected leaf cells was disintegration of the chloroplasts (Plate II). Furthermore, these cells having disintegrated chloroplasts usually accompanied with breakdown of tonoplast and production of granules within the cytoplasm. These structural changes are almost the same as those of tomato plants infected with PSTV and CEV (Plate III and IV).

Cytopathic changes in tomato plants infected with HSV and CPFV

Tomato plants showed no symptoms when infected with HSV or CPFV, but was proved to be capable of carrying these agents⁹. Therefore, it was rather difficult to find out distinctive alterations of leaf cells in the infected tomato plants. Disintegration of chloroplasts, however, sometimes (not often) occurred in these systems.

Cytopathic changes in tomato plants infected with PSTV and CEV

Symptoms caused by CEV were similar to those caused by PSTV in tomato plants, and were very severe in both cases. Cytopathic changes caused by both viroid infections were also almost the same (Table 1). Aberrations of cell walls such as corrugated profile and irregular thickness in tomato plants infected by PSTV and CEV were frequently observed. In addition, characteristic alterations of the chloroplasts within mesophyll cells were also frequently found (Plate III and IV).

Sequential events of chloroplast disintegration by viroid infections

Disintegration process of the chloroplasts in tomato palisade and spongy parenchymatous mesophyll cells infected by PSTV and CEV has some similarities in cucumber plants as mentioned above. They are as follows; ① swelling of the chloroplasts, ② aberration of their thylakoid membranes system, ③ distortion of their grana, ④ irregular arrangement of their stroma lamellar, ⑤ breakdown of their en-




Fig. 1. Schematic presentation of chloroplast disintegration procedure in association with viroid infections.

velope, and ⑥ release of their inner contents into cytoplasm (Fig. 1). However, no differences were observed in the size and number of starch grains and plastoglobules (osmiophilic globules) in the infected and healthy plants (Plate II, III and IV).

Since no aberration of membrane systems of the chloroplasts was found in healthy plants (Plate II A and B), and the mesophyll cells adjacent to the diseased cells showing chloroplast aberrations showed no cytopathic changes having intact chloroplasts (Plate III, left), we presumed that such aberrations of membrane systems of the chloroplasts were one of the cytopathic effects in viroid infections. There are some similarities between these cytopathic changes and those caused by some plant virus infections^{1,6}.

Other cytopathic changes in viroid-infected plants

In healthy mature mesophyll cells, cytoplasm containing cell organelles is usually located at the periphery of cell walls, and vacuoles limited by tonoplast are located in the middle of the cells (Plate II A and B). On the other hand, in the viroid-infected plants, other cytopathic changes such as disappearance of central vacuole and plasmolysis were also frequently found (Plate III and IV). These denatured cells were vertically distributed from palisade to spongy parenchymatous cells to form a denatured cell zone (Fig. 2).

On the basis of our observations, cytopathic changes in viroid-infected plants, such as aberrations of cell walls and distortions of chloroplast membranes, in association with their external symptoms, were frequently found in both cucumber-HSV and-CPFV, and tomato-PSTV and-CEV systems, but indistinctive in tomato-HSV and-CPFV symptomless system.

It is unlikely that paramural bodies^{8,9,11} are the primary cytopathic effects in viroid-infected plants because these structures are also present in healthy plants. It is also rather difficult to conclude from our investigation that "the viroid specific cytopathic changes are alterations of the paramural bodies themselves" as mentioned by WAHN *et al.*^{13,14}




Fig. 2. Schematic presentation of a denatured cell zone in a viroid-infected plant, in which some denatured cells (oblique lined) were vertically distributed from palisade to spongy parenchymatous cells.

Acknowledgement

The authors wish to express appreciation to Dr. H. L. SÄNGER, Max-

Planck-Institut für Biochemie, Martinsried bei München, West Germany, for his kind supply of CPFV, CEV and PSTV, and tomato (cv. Rentita) seeds.

Literature Cited

1. APPIANO, A., D'AGOSTINO, G., REDOLFI, P. and PENNAZIO, S.: Sequence of cyto-logical events during the process of local lesion formation in the tomato bushy stunt virus- *Gomphrena globosa* hypersensitive system, *J. Ultrastr. Res.* **76**: 173-180. 1981
2. DAGRACA, J. V. and MARTIN, M. M.: Ultrastructural changes in avocado leaf tissue infected with avocado sunblotch, *Phytopath. Z.* **102**: 185-194. 1981
3. DESJARDINS, P. R. and DRAKE, R. J.: Ultrastructural cytology of avocado infected with the sunblotch viroid, *Abst. 5th Internat. Cong. Virology, Strasbourg* p 28. 1981
4. DIENER, T. O.: Viroids and Viroid Diseases, Wiley- Interscience Publication John Wiley and Sons, Inc. 1979
5. EHARA, Y. and MISAWA, T.: Occurrence of abnormal chloroplasts in tobacco leaves infected systemically with the ordinary strain of cucumber mosaic virus, *Phytopath. Z.* **84**: 233-252. 1975
6. HARI, V.: Ultrastructure of potato spindle tuber viroid infected tomato leaf tissue, *Phytopathology* **70**: 385-387. 1980
7. MONMA, T. and TAKAHASHI, T.: Pathogenicity of plant viroid VI. Ultrastructure of hop stunt viroid infected cells, *Ann. Phytopath. Soc. Japan* **47**: 417 (abstr.). 1981
8. PALIWAL, Y. C. and SINH, R. P.: Cytopathological changes induced by potato spindle tuber viroid in *Scopolia sinensis*, *Can. J. Bot.* **59**: 677-682. 1981
9. SANO, T., SASAKI, M. and SHIKATA, E.: Comparative studies on hop stunt viroid, cucumber pale fruit viroid and potato spindle tuber viroid, *Ann. Phytopath. Soc. Japan* **47**: 599-605. 1981
10. SASAKI, M. and SHIKATA, E.: Studies on the host range of hop stunt disease in Japan, *Proc. Japan Acad.* **53**, Ser. B: 103-108. 1977
11. SEMANCIK, J. S. and VANDERWOUDE, W. J.: Exocortis viroid: Cytopathic effects at the plasma membrane in association with pathogenic RNA, *Virology* **69**: 719-726. 1976
12. SHIKATA, E. and KITAGAWA, Y.: Rice blace-streaked dwarf virus. Its properties, morphology and intracellular localization, *Virology* **77**: 826-842. 1977
13. WAHN, K., GOMEZ, F. R. und SÄNGER, H. L.: Cytopathologie von viroid-infizierten bei *Gynura aurantiaca* nach Infektion mit dem Viroid der Citrus-Exocortis-Krankheit (CEV), *Phytopath. Z.* **98**: 1-18. 1980
14. WAHN, K., GOMEZ, F. R. and SÄNGER, H. L.: Cytopathic changes in leaf tissue of *Gynura aurantiaca* infected with the viroid of exocortis disease, *J. gen. Virol.* **49**: 355-365. 1980

Explanation of Plate

Plate I

Aberrations (arrows) of cell walls of cucumber plants infected with CPFV (A and B) and HSV (C).

Photo A. upper epidermis. $\times 6,400$

Photo B. vascular bundles. $\times 6,400$

Photo C. lower epidermis. $\times 6,400$






Plate II

Disintegration of chloroplasts in plants associated with viroid infections.

Photos A, B. Ultrathin sections of palisade (A) and spongy (B) parenchymatous mesophyl cells of healthy cucumber plants. $\times 6,000$

N: nucleus; C: chloroplast

Photo C. A palisade parenchymatous cell of a cucumber plant infected by HSV. Note disappearance of the envelopes and aberrations of thylakoid membrane systems of chloroplasts. $\times 6,000$

Photo D. A spongy parenchymatous cell of a tomato plant infected by CEV. Note various stages of chloroplast disintegration. $\times 6,000$

double arrow: double layers; single arrow: single layer; star: no layers; white arrow: plastoglobule






Plate III

Cytopathic changes in the palisade parenchymatous cells of a tomato plant infected by CEV. Note a cell (left) in which cell organelles are normally arranged and two neighbouring affected cells showed disappearance of tonoplast and breakdown of chloroplast envelope. $\times 6,400$




Plate IV

Cytopathic changes in the palisade parenchymatous cells of a tomato plant infected by PSTV. Note disintegration of chloroplast membrane systems, disappearance of tonoplasts and plasmolysis (arrow). $\times 6,400$

