



Title	SEROLOGICAL PROPERTIES OF THE MOSQUITOCIDAL PROTEIN OF BACILLUS THURINGIENSIS AND THE MORPHOLOGY OF ITS PARASPORAL CRYSTAL
Author(s)	IIZUKA, Toshihiko; YAMAMOTO, Takashi
Citation	Journal of the Faculty of Agriculture, Hokkaido University, 62(1), 98-114
Issue Date	1984-11
Doc URL	http://hdl.handle.net/2115/13015
Type	bulletin (article)
File Information	62(1)_p98-114.pdf



[Instructions for use](#)

**SEROLOGICAL
PROPERTIES OF THE MOSQUITOCIDAL PROTEIN
OF *BACILLUS THURINGIENSIS* AND
THE MORPHOLOGY OF ITS
PARASPORAL CRYSTAL**

Toshihiko IZUKA

(Laboratory of Sericology, Faculty of Agriculture,
Hokkaido University, Sapporo 060, Japan)

Takashi YAMAMOTO

(USDA, ARS, Cotton Insects Research Unit,
Brownsville, Texas 78520, USA)

Received May 15, 1984

Introduction

Bacillus thuringiensis has been known as an insect pathogen and numerous strains have been isolated from diseased insects in many countries. They are classified into some 20 subspecies based on the flagellar antigen (H-antigen) and biochemical tests^{4,5}. Most subspecies normally produce a spore and a bipyramidal crystal in the cell at the end of the logarithmic phase of growth. Several subspecies, which have mosquitocidal activity, produce irregular and/or cuboidal crystals^{6,10}. The crystals have been purified by isopycnic centrifugation in a NaBr density gradient^{1,18}.

YAMAMOTO and McLAUGHLIN¹⁸ have isolated two proteins of Mr 135,000 and Mr 65,000 from the crystals of mosquitocidal HD-1 strain (subsp. *kurstaki*) after an exposure of its crystals to 2-mercaptoethanol at pH 10. The 135 Kdal protein, which was the major component of the crystal and was toxic to several lepidopterous species, showed no toxicity to mosquito larvae. The 65 Kdal protein, which differed entirely from the 135 Kdal protein on the basis of biochemical tests, showed a toxicity to mosquito larvae as high as its toxicity to larvae of lepidopterous species. Furthermore, IZUKA and YAMAMOTO¹¹ suggested that the mosquitocidal 65 Kdal protein crystalize in a cuboidal form.

The irregularly shaped crystals of *B. thuringiensis* subsp. *israelensis* (ONR-60 A), which was isolated by GOLDBERG and MARGALIT⁷ from soil

samples taken at a mosquito larval breeding site, has showed very high toxicity to mosquito larvae¹⁷. Mosquitocidal toxin of subsp. *israelensis* have been identified as 28 Kdal protein¹⁷.

In the present study, we tested the crystal proteins from 42 strains of different *B. thuringiensis* serotypes by immunoelectrophoresis, and their crystal shapes were examined with a scanning electron microscope (SEM).

Materials and Methods

Bacterial strains

Strains and sources of *B. thuringiensis* used in this study were listed in the Table 1.

Immunoelectrophoresis

We used rocket-immunoelectrophoresis which has been described by LAURELL¹². P-1 (135 Kdal protein) and P-2 (65 Kdal protein) were isolated from the lyophilized crystal of subsp. *kurstaki* HD-1 according to the method of YAMAMOTO and McLAUGHLIN¹⁸ and used as antigens. Three antisera directed against HD-1, against P-2 protein, and against *israelensis* crystal were made in rabbit. The antigens were dialyzed in 5 mM Tris-HCl (pH 8) at 4°C and inoculated into rabbit's shoulder with Freund's adjuvants. Details of the immunization procedures were as described by CLARK and FREEMAN² and by YAMAMOTO and TANADA¹⁹.

For immunoelectrophoresis, a loopful of spores and crystals which were incubated at 30°C, for 72 hr, was suspended in 30 μ l of 2% 2-mercaptoethanol whose pH was adjusted to 10 with NaOH. After 30 min incubation on ice, 10 μ l of Tris-HCl (pH 7) were added. The sample (4 μ l) was electrophoresed on 1.2% agarose gel plate (10 \times 10 \times 0.1 cm) containing 125-250 μ l antiserum. After electrophoresis at 100 V for 3 hr, the gel was washed in 0.8% NaCl-10 mM Tris-HCl (pH 8) for 16 hr at 37°C and stained with 0.25% Coomassie blue R-250 in 7% acetic acid and 30% methanol.

SEM

Bacteria were cultured on nutrient agar (Difco) at 30°C until almost all cells lysed (about 72 hr). The crystals and spores (about 100 mg wet weight) were washed in 10 ml of 50 mM Tris-HCl (pH 8) followed by water by repeated centrifugation at 10,000 rpm for 10 min. The final precipitate was resuspended in 1 ml of water, and 20 μ l of the suspension were air-dried on an aluminum disk. After the sample was coated with carbon and gold, it was observed and photographed with a SEM (JEOL, JSM-S1).

TABLE 1. Sources of the strains in *Bacillus thuringiensis*

Serotype H	Subspecies epithet	Source
1	<i>thuringiensis</i> Berliner	H. DE BARJAC, France
1	<i>thuringiensis</i> (BA-068)	E. REEVES, USA
2	<i>finitimus</i>	A. M. HEIMPEL, USA
3a	<i>alesti</i>	H. DE BARJAC, France
3a, 3b	<i>kurstaki</i> (HD-1)	H. DULMAGE, USA
3a, 3b	<i>kurstaki</i> (MC)	S. T. AMONKAR, India
4a, 4b	<i>sotto</i>	Insect Pathology Lab., Beltsville, USA
4a, 4b	<i>dendrolimus</i>	(id.)
4a, 4c	<i>kenyae</i>	H. DE BARJAC, France
5a, 5b	<i>galleriae</i>	(id.)
5a, 5c	<i>canadensis</i>	(id.)
6	<i>subtoxicus</i>	(id.)
6	<i>entomocidus</i>	Insect Pathology Lab., Beltsville, USA
7	<i>aizawai</i>	K. AIZAWA, Japan
7	<i>aizawai</i> (<i>juroi</i>)	ATCC (#21281) Rockville, USA
7	<i>aizawai</i> (HU)	Induced by T. IIZUKA from <i>aizawai</i> .
7	<i>aizawai</i> (B106)	Shionogi Co. Ltd., Japan
8a, 8b	<i>morrisoni</i>	H. DE BARJAC, France
8a, 8c	<i>ostrinae</i> (HD-501)	H. DULMAGE, USA
9	<i>tolworthi</i>	H. DE BARJAC, France
10	<i>darmstadiensis</i>	(id.)
10	<i>darmstadiensis</i> (73-E-10-2)	K. AIZAWA, Japan
10	<i>darmstadiensis</i> (73-E-10-16)	(id.)
10	<i>darmstadiensis</i> (73-E-37-14)	(id.)
11a, 11b	<i>toumanoffi</i>	H. DE BARJAC, France
11a, 11c	<i>kyushuensis</i>	K. AIZAWA, Japan
12	<i>thompsoni</i>	H. DE BARJAC, France
13	<i>pakistani</i> (HD-395)	H. DULMAGE, USA
14	<i>israelensis</i> (ONR-60A)	L. GOLDBERG, USA
14	<i>israelensis</i> (922903)	L. A. BULLA
14	<i>israelensis</i> (922906)	(id.)
14	<i>israelensis</i> (922910)	(id.)
14	<i>israelensis</i> (922917)	(id.)
14	<i>israelensis</i> (922918)	(id.)

Table 1. Continued

Serotype H	Subspecies epithet	Source
14	<i>israelensis</i> (922920)	(id.)
15	<i>dakota</i> (HD-511)	H. DULMAGE, USA
16	<i>indiana</i> (HD-516)	(id.)
17	<i>tohokuensis</i>	K. AIZAWA, Japan
18	<i>kumamotoensis</i>	(id.)
19	<i>tochigiensis</i>	(id.)
	<i>yunnanensis</i>	(id.)
	<i>wuhanensis</i>	H. DULMAGE, USA

Results and Discussion

The crystals of *B. thuringiensis* strains listed in Table 1, were examined by immunoelectrophoresis using anti-*kurstaki*-serum and the results were shown in Fig. 1 and Fig. 2. The HD-1 crystal showed two peaks, one with 135 Kdal protein and the other 65 Kdal protein¹⁰. The peak running from the left (negative electrode) to the right (positive electrode) on the gel was of 135 Kdal protein, and the peak running to the opposite direction was of 65 Kdal protein. It was recognized that four strains, subspp. *kurstaki* HD-1, *kurstaki* MC, *kenyae* and *tolworthi* produced the peak on the negative electrode side. This indicates that crystals of subspp. *kenyae* and *tolworthi* contain a protein similar to, if not the same as, the 65 Kdal mosquitocidal toxin of HD-1.

Anti-P-2-serum was also used to check crystals of the strains listed in Table 1 (Fig. 3 and Fig. 4). It confirmed the observation with anti-*kurstaki*-serum that subspp. *kurstaki* HD-1, *kurstaki* MC, *kenyae* and *tolworthi* contains the 65 Kdal P-2 protein.

Anti-*israelensis*-serum precipitated proteins of subspp. *israelensis* and *aizawai* HU when the crystal proteins from the strains listed in Table 1 were examined by immunoelectrophoresis (Fig. 5 and Fig. 6).

Previously, crystals of many strains in *B. thuringiensis* have already been photographed by SEM¹⁰. In the present study, we observed morphological details of the crystals from strains which are reportedly mosquitocidal. They are subspp. *kurstaki* HD-1, *kenyae*, *galleriae*, *entomocidus*, *aizawai* and *tolworthi*⁸⁾; subspp. *israelensis* ONR-60 A³⁾; subspp. *kyushuensis*¹⁴⁾; and subspp. *darmstadiensis* 73-E-10-2 and 73-E-10-16¹⁶⁾. (Fig. 7, Fig. 8 and Fig. 9). In addition to these mosquitocidal strains, the crystal of subspp. *morrisoni* was photographed under SEM and the electron micrograph re-

vealed two crystal forms.

The shape of crystals of subspp. *kurstaki* HD-1, *kenyae*, and *tolworthi*, which were shown to produce P-2 toxin by immunoelectrophoresis, appeared to have a cuboidal body some of which were partly embedded in the bipyramidal crystal matrix as previously reported by SHARPE and BAKER¹⁶. The crystals of subspp. *galleriae* also contained cuboidal ones, but 65 Kdal protein did not appear on the agarose gel (Fig. 1, 2, and 3). Therefore, we re-examined subspp. *galleriae* by immunoelectrophoresis using a high concentration of antiserum (Fig. 10). As a result, the crystal protein of subspp. *galleriae* was faintly recognized by an increased volume of anti-P-2-serum, but it appeared to be different from the reaction on the proteins of subspp. *kurstaki* HD-1, *kurstaki* MC, *kenyae* and *tolworthi*. It was suggested that a crystal protein of subspp. *galleriae* had some limited antigenic sites similar to the P-2 protein of HD-1.

There were no differences in the shape of crystals among subspp. *israelensis* ONR-60 A, *israelensis* 922910 and *aizawai* HU whose proteins were precipitated with anti-*israelensis*-serum. However, crystals of subspp. *darmstadiensis* 73-E-10-2 and 73-E-10-16 which have been reported to have irregular shape by IIZUKA *et al.*⁹, did not react with anti-*israelensis*-serum.

Shapes of the crystals produced by the mosquitocidal strains of *B. thuringiensis*^{3,8,14,15}, include cuboidal form as seen in *kurstaki* HD-1 type crystals and irregular form in subspp. *israelensis* type crystals.

We stock four strains of subspp. *aizawai*, the original *aizawai* from Dr. K. AIZAWA, *aizawai* (*juroi*) from ATCC, USA, *aizawai* HU selected by T. IIZUKA, and *aizawai* B106 (sporeless) from Shionogi Co., Ltd.. Electron micrography showed significant differences in crystal morphology among these strains. We are very concerned about the differences and propose further studies on the plasmid patterns and the insecticidal activity of these strains.

Summary

In the present study, we have demonstrated that there are three serologically groups in a number of mosquitocidal *B. thuringiensis* δ -endotoxins. They are: group 1 including those produced by subspp. *kurstaki* HD-1, *kenyae* and *tolworthi* which were precipitated with anti-*kurstaki* HD-1-serum; group 2, subspp. *israelensis* and *aizawai* HU which reacted with anti-*israelensis*-serum; group 3, subspp. *galleriae*, *entomocidus* and *darmstadiensis* which showed no crossreaction with these two antisera. Morphology, of the crystals appeared to be different among these groups when they were examined by electron microscopy.

References

1. ANG, B. J. and NICKERSON, K. W.: Purification of the protein crystal from *Bacillus thuringiensis* by zonal gradient centrifugation. *Appl. Environ. Microbiol.* **36**: 625-626. 1978
2. CLARKE, H. G. M. and FREEMAN, T.: A quantitative immuno-electrophoresis method (Laurell electrophoresis). *Protides Biol. Fluids. Proc. Colloq.* **14**: 503-509. 1966
3. DE BARJAC, H.: Toxicite de *Bacillus thuringiensis* var. *israelensis* pour les larves d'*Aedes aegypti* et d'*Anopheles stephensi*. *Compt. Rend. Acad. Sci. (Paris)* **286**: 1175-1178. 1978
4. DE BARJAC, H. and BONNEFOI, A.: Essai de classification biochimique et serologique de 24 souches de *Bacillus* du type *Bacillus thuringiensis*. *Entomophaga* **7**: 5-31. 1962
5. DE BARJAC, H. and BONNEFOI, A.: Mise au point sur la classification des *Bacillus thuringiensis*. *Entomophaga* **18**: 5-17, 1973
6. FAUST, R. M., ADAMS, J. R., ABE, K., IIZUKA, T. and BULLA, L. A.: Comparative morphology and size distribution of the parasporal crystals from various subspecies of *Bacillus thuringiensis*. *J. Sericult. Sci. (Japan)* **51**: 316-324. 1982
7. GOLDBERG, L. J. and MARGALIT, J.: A bacterial spore demonstrating rapid laticidal activity against *Anopheles sergentii*, *Uranotaenia unguiculata*, *Culex univittatus*, *Aedes aegypti* and *Culex pipiens*. *Mosquito News* **37**: 355-358. 1977
8. HALL, I. M., ARAKAWA, K. Y., DULMAGE, H. D. and CORREA, J. A.: The pathogenicity of strains of *Bacillus thuringiensis* to larvae of *Aedes* and *Culex* mosquitoes. *Mosquito News* **37**: 246-251. 1977
9. IIZUKA, T., FAUST, R. M. and OHBA, M.: Comparative profile of plasmid DNA and morphology of parasporal crystals in four strains of *Bacillus thuringiensis* subsp. *darmsladiensis*. *App. Ent. Zool.* **18**: 486-494. 1983
10. IIZUKA, T., ISHINO, M. and NAKASHIMA, T.: Comparative morphology of parasporal crystals and characterizations of plasmid DNA from various subspecies of entomopathogenic bacteria, *Bacillus thuringiensis*. *Mem. Fac. Agric. Hokkaido Univ.* **13**: 423-431. 1982
11. IIZUKA, T. and YAMAMOTO, T.: Possible location of the mosquitocidal protein in the crystal preparation of *Bacillus thuringiensis* subsp. *kurstaki*. *FEMS Microbiol. Lett.* **19**: 187-192. 1983
12. LAURELL, C. B.: Electroimmunoassay. *Scand. J. Clin. Lab. Invest.* **29** (Suppl. 124): 21-37. 1972
13. NISHITSUTSUJI-UWO, J., WAKISAKA, Y. and EDA, M.: Sporeless mutant of *Bacillus thuringiensis*. *J. Invertebr. Pathol.* **25**: 355-361. 1975
14. OHBA, M. and AIZAWA, K.: A new subspecies of *Bacillus thuringiensis* possessing 11a: 11c flagellar antigenetic structure: *Bacillus thuringiensis* subsp. *kyushuensis*. *J. Invertebr. Pathol.* **33**: 387-388. 1979
15. PADUA, L. E., OHBA, M. and AIZAWA, K.: The isolates of *Bacillus thuringiensis*

- serotype 10 with a highly preferential toxicity to mosquito larva. *J. Invertebr. Pathol.* **36**: 180-186. 1980
16. SHARPE, E. S. and BAKER, F. L.: Ultrastructure of the unusual crystal of the HD-1 isolate of *Bacillus thuringiensis* var. *kurstaki*. *J. Invertebr. Pathol.* **34**: 320-322. 1979
 17. YAMAMOTO, T., IIZUKA, T. and ARONSON, J. N.: Mosquitocidal protein of *Bacillus thuringiensis* subsp. *israelensis*: Identification and partial isolation of the protein. *Curr. Microbiol* **9**: 279-284. 1983
 18. YAMAMOTO, T. and MCLAUGHLIN, R. E.: Isolation of a protein from the parasporal crystal of *Bacillus thuringiensis* var. *kurstaki* toxic to the mosquito larva, *Aedes taeniorhynchus*. *Biochem. Biophys. Res. Commun.* **103**: 414-421. 1981
 19. YAMAMOTO, T. and TANADA, Y.: Comparative analysis of the enveloped virions of two granulosis viruses of the armyworm, *Pseudaletia unipuncta*. *Virology* **94**: 71-81. 1979





Fig. 1. Crossed immunoelectrophoresis of dissociated crystals of a variety of *B. thuringiensis* strains against anti-*kurstaki* HD-1 serum.

The sample (4 μ l each) was electrophoresed on an agarose gel plate at 100 V for 3 hr at 4°C. The gel, which was made from 10 ml of 1.2% low electroendosmos agarose and contained 250 μ l antiserum and 100 mM Tris-acetate (pH 8.2), was cast on a 10 \times 10 cm glass plate. After the electrophoresis, the gel plate was washed in 0.8% NaCl buffered with 10 mM Tris-HCl (pH 8) for 16 hr at 37°C to remove unreacted serum component, dried in air, and stained with 0.25% Coomassie blue R-250. The positive electrode was on the right.



- | | |
|-------------------------------------|-------------------------------------|
| 1. <i>thuringiensis</i> Berliner | 16. <i>tolworthi</i> |
| 2. <i>fnitimus</i> | 17. <i>darmstadiensis</i> |
| 3. <i>alesti</i> | 18. <i>darmstadiensis</i> 73-E-10-2 |
| 4. <i>kurstaki</i> HD-1 | 19. <i>toumanoffi</i> |
| 5. <i>kurstaki</i> MC | 20. <i>kyushuensis</i> |
| 6. <i>sotto</i> | 21. <i>thompsoni</i> |
| 7. <i>dendrolimus</i> | 22. <i>pakistani</i> |
| 8. <i>kenyae</i> | 23. <i>israelensis</i> ONR-60A |
| 9. <i>galleriae</i> | 24. <i>indiana</i> |
| 10. <i>canadensis</i> | 25. <i>dakota</i> |
| 11. <i>subtoxicus</i> | 26. <i>tohokuensis</i> |
| 12. <i>entomocidus</i> | 27. <i>kumamotoensis</i> |
| 13. <i>aizawai</i> (<i>juroi</i>) | 28. <i>tochigiensis</i> |
| 14. <i>morrisoni</i> | 29. <i>yunnanensis</i> |
| 15. <i>ostrinae</i> | 30. <i>wuhanensis</i> |

Fig. 2. Crossed immunoelectrophoresis of dissociated crystals against anti-*kurstaki* HD-1-serum in the additional strains to Fig. 1.





Fig. 3. Crossed immunoelectrophoresis of dissociated crystals against anti-P-2-serum in the strains of *B. thuringiensis*.



- | | |
|--------------------------------------|-------------------------------------|
| 1. <i>thuringiensis</i> Berliner | 16. <i>tolworthi</i> |
| 2. <i>finitimus</i> | 17. <i>darmstadiensis</i> |
| 3. <i>alesti</i> | 18. <i>darmstadiensis</i> 73-E-10-2 |
| 4. <i>kurstaki</i> HD-1 | 19. <i>toumanoffi</i> |
| 5. <i>kurstaki</i> MC | 20. <i>kyushuensis</i> |
| 6. <i>sotto</i> | 21. <i>thompsoni</i> |
| 7. <i>dendrolimus</i> | 22. <i>pakistani</i> |
| 8. <i>kenyae</i> | 23. <i>israelensis</i> ONR-60A |
| 9. <i>galleriae</i> | 24. <i>indiana</i> |
| 10. <i>canadensis</i> | 25. <i>dakota</i> |
| 11. <i>subtoxicus</i> | 26. <i>tohokuensis</i> |
| 12. <i>entomocidus</i> | 27. <i>kumamotoensis</i> |
| 13. <i>aizawaai</i> (<i>juroi</i>) | 28. <i>tochigiensis</i> |
| 14. <i>morrisoni</i> | 29. <i>yunnanensis</i> |
| 15. <i>ostrinae</i> | 30. <i>wuhanensis</i> |

Fig. 4. Crossed immunoelectrophoresis of dissociated crystals against anti-P-2-serum in the additional strains to Fig. 3.





Fig. 5. Crossed immunoelectrophoresis of dissociated crystals against anti-*israelensis*-serum in the strains of *B. thuringiensis*.



- | | |
|-------------------------------------|-------------------------------------|
| 1. <i>thuringiensis</i> Berliner | 16. <i>tolworthi</i> |
| 2. <i>finitimus</i> | 17. <i>darmstadiensis</i> |
| 3. <i>alesti</i> | 18. <i>darmstadiensis</i> 73-E-10-2 |
| 4. <i>kurstaki</i> HD-1 | 19. <i>toumanoffi</i> |
| 5. <i>kurstaki</i> MC | 20. <i>kyushuensis</i> |
| 6. <i>sotto</i> | 21. <i>thompsoni</i> |
| 7. <i>dendrolimus</i> | 22. <i>pakistani</i> |
| 8. <i>kenyae</i> | 23. <i>israelensis</i> ONR-60A |
| 9. <i>galleriae</i> | 24. <i>indiana</i> |
| 10. <i>canadensis</i> | 25. <i>dakota</i> |
| 11. <i>subtoxicus</i> | 26. <i>tohokuensis</i> |
| 12. <i>entomocidus</i> | 27. <i>kumamotoensis</i> |
| 13. <i>aizawai</i> (<i>juroi</i>) | 28. <i>tochigiensis</i> |
| 14. <i>morrisoni</i> | 29. <i>yunnanensis</i> |
| 15. <i>ostrinae</i> | 30. <i>wuhanensis</i> |

Fig. 6. Crossed immunoelectrophoresis of dissociated crystals against anti-*israelensis*-serum in the additional strains to Fig. 5.




Fig. 7. Electron micrographs of crystals of *B. thuringiensis* subspecies associating which were reported to have mosquitocidal activity by HALL *et al.*⁹⁾.




Fig. 8. Electron micrographs of crystals of *B. thuringiensis* subsp. *aizawai*.




Fig. 9. Electron micrographs of crystals of *B. thuringiensis* subspecies whose mosquitocidal activity were reported by HALL *et al.*⁸⁾, by DE BARJAC³⁾, and by OHBA and AIZAWA¹⁴⁾. The micrographs of subsp. *morrisoni* crystals appeared to be similar to that of subsp. *pakistani* crystals.




Fig. 10. Crossed immunoelectrophoresis of dissociated crystals against anti-P-2-serum in the subsp. *galleriae*. An increased volume of anti-serum ($450 \mu\text{l}$) was added to the agarose gel.