Title	Harlequin ichthyosis and other autosomal recessive congenital ichthyoses : The underlying genetic defects and pathomechanisms
Author(s)	Akiyama, Masashi
Citation	Journal of Dermatological Science, 42(2), 83-89 https://doi.org/10.1016/j.jdermsci.2006.01.003
Issue Date	2006-05
Doc URL	http://hdl.handle.net/2115/13466
Type	article (author version)
File Information	main.pdf

[Instructions for use](#)

Journal of Dermatological Science

Manuscript No. JDS-05-332 Revised Version

REVIEW ARTICLE

Harlequin ichthyosis and other autosomal recessive congenital ichthyoses: the underlying genetic defects and pathomechanisms

Masashi Akiyama

Department of Dermatology, Hokkaido University Graduate School of Medicine,
Sapporo 060-8638, Japan

Word count (text only): 2137 words

Correspondence and reprint requests to:

Masashi Akiyama, MD, PhD
Department of Dermatology
Hokkaido University Graduate School of Medicine
North 15 West 7, Sapporo 060-8638, Japan
Telephone: +81-11-716-1161, ext. 5962
Fax: +81-11-706-7820
e-mail <akiyama@med.hokudai.ac.jp>

Summary

Autosomal recessive congenital ichthyoses (ARCI) include several severe subtypes including harlequin ichthyosis (HI), lamellar ichthyosis and non-bullous congenital ichthyosiform erythroderma. Patients with these severe types of ichthyoses frequently show severe hyperkeratosis and scales over a large part of the body surface from birth and their quality of life is often severely affected. Recently, research into the pathomechanisms of these severe congenital ichthyoses have advanced dramatically and led to the identification of several causative genes and molecules underlying the genetic defects. To date, seven loci have been identified that are associated with ARCI and, among them, five causative genes and molecules have been detected. The five genes are transglutaminase 1 gene (*TGMI*), *ABCA12*, two lipoxygenase genes, *ALOXE3* and *ALOX12B*, and *ichthyin*. One of these components, *ABCA12*, has recently been shown to be a keratinocyte lipid transporter associated with lipid transport in lamellar granules and loss of *ABCA12* function leads to a defective lipid barrier in the stratum corneum, resulting in the HI phenotype. Transglutaminase 1 deficiency was reported to cause a malformed cornified cell envelope leading to a defect in the intercellular lipid layers in the stratum corneum and defective stratum corneum barrier function resulting in an ichthyosis phenotype. Thus, defective intercellular lipid layers are major findings in autosomal recessive congenital ichthyoses. Information concerning ARCI genetic defects and disease pathomechanisms are beneficial for providing better treatments and genetic counseling including prenatal diagnosis for families affected by ichthyoses.

Keywords: *ABCA12*; harlequin ichthyosis; lamellar granules; lamellar ichthyosis; non-bullous congenital ichthyosiform erythroderma; prenatal diagnosis

1. Introduction

Severe autosomal recessive congenital ichthyoses (ARCI) can be devastating to a patients' quality of life in those seriously affected, even though other organs are uninvolved. Harlequin ichthyosis (HI; OMIM #242500), especially, is often fatal in affected newborns [1]. For a long time, the pathomechanisms and underlying genetic defects were unknown, although significant progress has recently been made in the understanding of the molecular basis of human epidermal keratinization processes. Since transglutaminase 1 gene (*TGMI*) mutations were identified as the cause in lamellar ichthyosis (LI) in 1995 [2, 3], mutations in several other genes have also been identified in LI and non-bullous congenital ichthyosiform erythroderma (NBCIE). Mutations in any of the known causative genes can lead to either NBCIE or LI and candidate genes specific to either NBCIE or LI alone have not yet been identified. Based on these facts, it is wise to consider NBCIE and LI as a general blanket diagnosis of covering and range of different disorders with similar and overlapping clinical features [4].

In 2003, *ABCA12* was first reported as the causative gene in type 2 LI (OMIM #601277) [5]. Recently, we have determined *ABCA12* function and clarified the pathomechanisms of *ABCA12* mutations leading to the HI phenotype [6]. *ABCA12* is a keratinocyte transmembrane lipid transporter protein associated with lipid transport in lamellar granules to the surface of granular layer keratinocytes [6]. In addition, several newly recognized molecules have been identified as causes of ARCI. In this article, recent research advances into the pathomechanisms of ARCI including HI and the feasibility of performing DNA-based prenatal diagnoses for ARCI are further discussed.

2. Defective intercellular lipid is the main idea behind the pathomechanisms of autosomal recessive congenital ichthyosis (ARCI)

The underlying genetic defects in ARCI patients, to date, a total of seven loci including candidate genes have been reported as shown in Table 1 [5-13]. The genes responsible in 14q11.2, 2q34, 17p13.1 and 5q33 were identified as *TGMI* [2, 3], *ABCA12* [5, 14], two lipoxygenase genes, *ALOXE3* and *ALOX12B* [15], and *ichthyin* [12], respectively (see Table 1).

Formation of the intercellular lipid layers is essential for epidermal barrier function and defective formation of the lipid layers is thought to result in serious loss of the epidermal barrier function, and to abnormal hyperkeratosis (Fig. 2) [16, 17]. Mutations in the lipid transporter protein, *ABCA12* cause defective lipid secretion into lamellar granules which then become expelled from the apical surface of keratinocytes [6] as mentioned above and lead to either LI [5] and HI [6, 18] phenotypes. Lipoxygenase-3 and 12(R)-lipoxygenase are non-heme iron-containing dioxygenases expressed in the epidermis, and their exact functions are unknown [19, 20]. They may be associated with lipid metabolism in the lamellar granule contents and/or intercellular lipid layers in the epidermis. In addition to HI and LI with defective lipid layers in the stratum corneum as their main pathogenetic mechanism, ichthyosis syndromes are also thought to share similar pathomechanisms. For example, Dorfman-Chanarin syndrome (neutral lipid storage disease) showed malformation of lamellar granules and defective lipid production in lamellar granules caused by a deficiency in the CGI-58 protein that is thought to be involved in the pathogenesis of this form of ichthyosis [21]. In Sjgren-Larsson syndrome harboring fatty aldehyde dehydrogenase (FALDH) gene

(ALDH3A2) mutations, defective lamellar granule formation was reported as one sign of a putative pathogenetic mechanism producing ichthyotic lesions in the patients' skin [22].

Similarly, transglutaminase 1 is an enzyme that was reported to crosslink hydroxyceramide to the cornified cell envelope [23, 24]. Cornified cell envelope formation is essential as the scaffold upon which normal intercellular lipid layer formation in the stratum corneum can then take place [25]. Mutations in the transglutaminase 1 gene (*TGMI*) cause defects in the intercellular lipid layers in the stratum corneum leading to defective barrier function of the stratum corneum resulting in the ichthyosis phenotype seen in LI patients [25] and in transglutaminase 1 knockout mice [26]. It remains to be clarified to what extent a defective cornified cell envelope alone contributes to the barrier abnormality in these conditions. In this context, it is of no doubt that defective formation of the intercellular lipid layers in the stratum corneum due to abnormal keratinocyte lipid metabolism, transport, and/or secretion is a major pathogenetic mechanism in this group of congenital ichthyoses, even if known or as yet undiscovered molecules may also play a significant part in the disease process.

It is certain that transglutaminase 1 is one of the major causative molecules of ARCI [4, 27]. Majority of the cases with *TGMI* mutations show LI phenotype. Most *TGMI* mutations reported in LI families are located in the core domain or upstream of it [4, 27]. *TGMI* mutations were also reported to lead to an NBCIE phenotype in a small number of the families [28-30] and defective transglutaminase 1 is not specific for the classic LI phenotype. We can hypothesize that serious loss of transglutaminase 1

activity might lead to the classic LI phenotype [27] and a partial loss of transglutaminase 1 activity to a mild form of LI [31] or a phenotype of NBCIE [29]. Further accumulation of the ARCI patients with *TGMI* mutations is needed to confirm the correlation between genotype and phenotype in this subgroup of ARCI patients.

Ichthyin is a protein with several transmembrane domains, which belongs to a new family of proteins with an unknown function. Ichthyin-like proteins are localized in the plasma membrane, and share with homologies to both transporters and G-protein coupled receptors [12]. Ichthyin was suggested to be a membrane receptor for certain ligands (trioxilins A3 and B3) from the hepoxilin pathway [12]. However, the exact mechanisms of how *ichthyin* mutations cause an ichthyosis phenotype remain to be clarified.

A large outstanding number of patients with unknown genetic defects still suffer from LI and NBCIE forms. Thus, we cannot exclude the possibility that distinct pathomechanisms not involving abnormal stratum corneum lipid barrier formation underlie such groups of patients. For example, an enhanced serine protease activity was demonstrated as a pathomechanism in the mouse model of Netherton syndrome, one of the major ichthyosis syndromes [32].

From the studies of molecular genetic defects underlying each type of autosomal recessive congenital ichthyosis subtype, the majority of missense mutations in ABCA12 were reported to cause LI phenotype, although one NBCIE case was also reported to harbor ABCA12 missense mutations [5]. Mutations in transglutaminase 1 gene result in phenotypes with both LI [2, 3] and NBCIE [28-30].

Mutations in lipoxygenase-3 and 12(R)-lipoxygenase genes have been reported to lead to both LI and NBCIE phenotypes [15]. Mutations in *ichthyin*, in the majority of patients caused the NBCIE phenotype, however an LI phenotype has been reported in one case [12]. We have clarified that even HI shares the same causative gene, *ABCA12* with type 2 LI [6]. These facts clearly indicate that mutations in any of the known causative genes are not specific to one disease phenotype alone. Thus, based on a comprehensive understanding of the pathophysiology of these diseases, HI, LI and NBCIE can be considered to be a group of distinct but related disorders with overlapping phenotypes.

Until now, no clear genotype/phenotype correlation has been defined for mutations in any of the causative genes leading to the ARCI patient phenotype, except for HI and type 2 LI. Thus, it seems that for *ABCA12* related diseases truncation and/or deletion mutations, at least in one allele, cause an HI phenotype and that a homozygote or a compound heterozygote harboring missense mutations results in LI [5, 6, 18].

3. *ABCA12* deficiency in harlequin ichthyosis and lamellar ichthyosis type 2

Among the severe ARCI, harlequin ichthyosis is the most devastating congenital ichthyosis and affected newborns show large, thick, plate-like scales over the whole body and severe ectropion, eclabium and flattened ears [17]. Type 2 LI is a subtype of LI which links to 2q33-35. Clinically, no distinct characteristic features are known for this type of LI [5, 7, 8]. Mutations in *ABCA12* underlie HI and type 2 LI.

ABCA12 is a member of the large superfamily of the ATP-binding cassette (ABC)

transporters, which bind and hydrolyze ATP to transport various molecules across a limiting membrane or into a vesicle [33]. The ABCA subfamily members are thought to be lipid transporters [34].

Ultrastructurally, lamellar granule abnormalities are apparent in HI patient epidermis [35-38]. The cornified cell envelope appears to be normal in HI and major cornified cell envelope precursor proteins (involucrin, small proline-rich proteins 1 and 2, and loricrin) are normally distributed in the HI epidermis [39, 40]. From these findings, HI and type 2 LI diseases are thought to be caused by a different pathomechanisms from malformations in cornified cell envelope. Several morphologic abnormalities, for example the abnormal lamellar granules in the granular layer keratinocytes and a lack of extracellular lipid lamellae in the stratum corneum, reflect the defective lipid transport via lamellar granules and the malformation of intercellular lipid layers in the stratum corneum in HI [35-39]. In type 2 LI, no distinct ultrastructural features have yet been reported. Abnormal lamellar granules in the granular layer keratinocytes and accumulation of lipid droplets in the stratum corneum, similar findings to that seen in HI, were reported in LI cases without transglutaminase 1 deficiency and some in these LI cases might belong to type 2 LI [41, 42].

The HI patients' epidermis was shown to have defective lipid transport using lamellar granules [6]. In addition, cultured epidermal patient keratinocytes carrying *ABCA12* mutations demonstrated defective glucosylceramide transport but that this phenotype was recoverable by ABCA12 corrective gene transfer [6]. From these findings, we have shed light on the pathomechanisms of HI with *ABCA12* mutations that cause a loss of ABCA12 function. Lack of ABCA12 function subsequently leads to disruption

of lamellar granule lipid transport in the upper keratinizing epidermal cells resulting in malformation of the intercellular lipid layers of the stratum corneum [6]. The fact that ABCA3 (a member of the same protein superfamily as ABCA12) works in pulmonary surfactant lipid secretion again using lamellar granules in lung alveolar type II cells [43] further supports our concept.

Genotype/phenotype correlations with five distinct ABCA12 mutations were reported in nine families of type 2 LI and all five mutations were missense mutations resulting in only one amino acid alteration [5]. In contrast, most mutations in HI are truncation or deletion mutations which lead to severe loss of function of ABCA12 peptide affecting important nucleotide-binding fold domains and/or transmembrane domains. In HI, at least one mutation on each allele must be a truncation or deletion mutation in the conserved region which seriously affects ABCA12 function [6]. Further accumulation of data on ABCA12 mutation protein effects and specific sites is needed to elucidate the genotype/phenotype correlations to aid in predicting HI patients' prognosis once novel combinations of ABCA12 mutations have been identified.

4. DNA-based prenatal diagnosis of severe congenital ichthyoses

ARCI are a group of the most severe genodermatoses and often the patients' quality of life is seriously affected. Thus, the parents' request for prenatal diagnosis is not to be ignored easily. Before the causative genes were identified, prenatal diagnosis had been performed by fetal skin biopsy and electron microscopic observation during the later stages of pregnancies at 19-23 weeks estimated gestational age mainly for HI for more than 20 years [44, 45]. Prenatal diagnosis of LI had been possible by ultrastructural observation of fetal skin samples, although prenatal diagnosis of LI was

the most high risk among the keratinization diseases because LI shows regional, individual, and familial variability in its expression. In the last decade, causative genes for ARCI have been clarified one by one and, as previously described, in 2005, we have identified the underlying gene causing HI [6]. Due to the recent advance in understanding of causative genetic defects for ARCI, it has now become possible to make DNA-based prenatal diagnosis for several ARCI diseases by chorionic villus or amniotic fluid sampling procedures in the earlier stages of pregnancy with a lower risk to fetal health and with a reduced burden on the mothers, as is the case with other severe genetic disorders [46]. In LI families with transglutaminase 1 gene mutations, successful prenatal diagnosis and prenatal exclusion of LI by transglutaminase 1 gene mutation analysis has already been reported [47, 48]. Theoretically, prenatal diagnosis by mutation analysis in lipoyxygenase-3, 12(R)-lipoyxygenase and ABCA12 is available in LI families with previously identified mutations on a case by case basis depending on gene mutations. In the near future, even earlier prenatal detection of ARCI by DNA analysis using fetal cells in maternal peripheral blood circulation [49], and, furthermore, preimplantation genetic diagnosis [50] should be available for ARCI.

Acknowledgements

I would like to thank Professor Hiroshi Shimizu for his critical reading of this manuscript and Professor. James R. McMillan for his proofreading and comments in the preparation of the manuscript.

References

- [1] Akiyama M. The pathogenesis of severe congenital ichthyosis of the neonate. *J Dermatol Sci* 1999; 21: 96-104.
- [2] Huber M, Rettler I, Bernasconi K, et al. Mutations of keratinocyte transglutaminase in lamellar ichthyosis. *Science* 1995; 267: 525-8.
- [3] Russell LJ, DiGiovanna JJ, Rogers GR, et al. Mutations in the gene for transglutaminase 1 in autosomal recessive lamellar ichthyosis. *Nature Genet* 1995; 9: 279-83.
- [4] Akiyama M, Sawamura D, Shimizu H. The clinical spectrum of nonbullous congenital ichthyosiform erythroderma and lamellar ichthyosis. *Clin Exp Dermatol* 2003; 28: 235-40.
- [5] Lefvire C, Audebert S, Jobard F, et al. Mutations in the transporter ABCA12 are associated with lamellar ichthyosis type 2. *Hum Mol Genet* 2003; 12: 2369-78.
- [6] Akiyama M, Sugiyama-Nakagiri Y, Sakai K, et al. Mutations in ABCA12 in harlequin ichthyosis and functional rescue by corrective gene transfer. *J Clin Invest* 2005; 115: 1777-84.
- [7] Parmentier L, Lakhdar H, Blanchet-Bardon C, Marchand S, Dubertret L, Weissenbach J. Mapping of a second locus for lamellar ichthyosis to chromosome 2q33-35. *Hum Mol Genet* 1996; 5: 555-9.
- [8] Parmentier L, Clepet C, Boughdene-Stambouli O, et al. Lamellar ichthyosis: further narrowing, physical and expression mapping of the chromosome 2 candidate locus. *Eur J Hum Genet* 1999; 7: 77-87.
- [9] Virolainen E, Wessman M, Hovatta I, et al. Assignment of a novel locus for autosomal recessive congenital ichthyosis to chromosome 19p13.1 p13.2. *Am J Hum Genet* 2000; 66: 1132-7.

- [10] Fischer J, Faure A, Bouadjar B, et al. Two new loci for autosomal recessive ichthyosis on chromosomes 3p21 and 19p12 q12 and evidence for further genetic heterogeneity. *Am J Hum Genet* 2000; 66: 904-13.
- [11] Krebsova A, Kuster W, Lestringant GG, et al. Identification, by homozygosity mapping, of a novel locus for autosomal recessive congenital ichthyosis on chromosome 17p, and evidence for further genetic heterogeneity. *Am J Hum Genet* 2001; 69: 216-22.
- [12] Lefvire C, Bouadjar B, Karaduman A, et al. Mutations in ichthyin a new gene on chromosome 5q33 in a new form of autosomal recessive congenital ichthyosis. *Hum Mol Genet* 2004; 13: 2473-82.
- [13] Mizrahi-Koren M, Geiger D, Indelman M, Bitterman-Deutsch O, Bergman R, Sprecher E. Identification of a novel locus associated with congenital recessive ichthyosis on 12p11.2-q13. *J Invest Dermatol* 2005; 125: 456-62.
- [14] Annilo T, Shulenin S, Chen ZQ, et al. Identification and characterization of a novel ABCA subfamily member, ABCA12, located in the lamellar ichthyosis region on 2q34. *Cytogenet Genome Res* 2002; 98: 169-76.
- [15] Jobard F, Lefvire C, Karaduman A et al. Lipoygenase-3 (ALOXE3) and 12(R)-lipoygenase (ALOX12B) are mutated in non-bullous congenital ichthyosiform erythroderma (NCIE) linked to chromosome 17p13.1. *Hum Mol Genet* 2002; 11: 107-13.
- [16] Williams ML, Schmuth M, Crumrine D, Hachem J-P, Bruckner AL, Demerjian M, Elias PM. Pathogenesis of the ichthyoses: update and therapeutic implications. *J Skin Barrier Res* 2005; 7: 122-33.
- [17] Akiyama M. Pathomechanisms of harlequin ichthyosis and ABC transporters in human diseases. *Arch Dermatol* (in press).

- [18] Kelsell DP, Norgett EE, Unsworth H, et al. Mutations in ABCA12 underlie the severe congenital skin disease harlequin ichthyosis. *Am J Hum Genet* 2005; 76: 794-803.
- [19] Yu Z, Schneider C, Boeglin WE, Marnett LJ, Brash AR. The lipoxygenase gene ALOXE3 implicated in skin differentiation encodes a hydroperoxide isomerase. *Proc Natl Acad Sci U S A* 2003; 100: 9162-7.
- [20] Yu Z, Schneider C, Boeglin WE, Brash AR. Mutations associated with a congenital form of ichthyosis (NCIE) inactivate the epidermal lipoxygenases 12R-LOX and eLOX3. *Biochim Biophys Acta* 2005; 1686: 238-47.
- [21] Akiyama M, Sawamura D, Nomura Y, Sugawara M, Shimizu H. Truncation of CGI-58 protein causes malformation of lamellar granules resulting in ichthyosis in Dorfman-Chanarin syndrome. *J Invest Dermatol* 121: 1029-34, 2003.
- [22] Shibaki A, Akiyama M, Shimizu H. Novel ALDH3A2 heterozygous mutations are associated with defective lamellar granule formation in a Japanese family of Sjgren-Larsson syndrome. *J Invest Dermatol* 123: 1197-9, 2004.
- [23] Marekov LN, Steinert PM. Ceramides are bound to structural proteins of the human foreskin epidermal cornified cell envelope. *J Biol Chem* 273: 17763-70, 1998.
- [24] Nemes Z, Marekov LN, Fesus L, Steinert PM. A novel function for transglutaminase 1: attachment of long-chain omega-hydroxyceramides to involucrin by ester bond formation. *Proc Natl Acad Sci USA* 96: 8402-7, 1999.
- [25] Elias PM, Schmuth M, Uchida Y, Rice RH, Behne M, Crumrine D, Feingold KR, Holleran WM, Pharm D. Basis for the permeability barrier abnormality in lamellar ichthyosis. *Exp Dermatol* 11: 248-56, 2002.
- [26] Kuramoto N, Takizawa T, Takizawa T, et al. Development of ichthyosiform skin

compensates for defective permeability barrier function in mice lacking transglutaminase 1. *J Clin Invest* 2002; 109: 243-50.

- [27] Akiyama M, Takizawa Y, Suzuki Y, Shimizu H. A novel homozygous mutation 371delA in TGM1 leads to a classic lamellar ichthyosis phenotype. *Br J Dermatol* 2003; 148: 149-53.
- [28] Laiho E, Ignatius J, Mikkola H, et al. Transglutaminase 1 mutations in autosomal recessive congenital ichthyosis: private and recurrent mutations in an isolated population. *Am J Hum Genet* 1997; 61: 529-38.
- [29] Akiyama M, Takizawa Y, Kokaji T, Shimizu H. Novel mutations of TGM1 in a child with congenital ichthyosiform erythroderma. *Br J Dermatol* 2001; 144: 401-7.
- [30] Becker K, Csikos M, Sardy M, et al. Identification of two novel nonsense mutations in the transglutaminase 1 gene in a Hungarian patient with congenital ichthyosiform erythroderma. *Exp Dermatol* 2003; 12: 324-9.
- [31] Akiyama M, Takizawa Y, Suzuki Y, Ishiko A, Matsuo I, Shimizu H. Compound heterozygous TGM1 mutations including a novel missense mutation L204Q in a mild form of lamellar ichthyosis. *J Invest Dermatol* 2001; 116, 992-5.
- [32] Descargues P, Deraison C, Bonnart C, Kreft M, Kishibe M, Ishida-Yamamoto A, Elias P, Barrandon Y, Zambruno G, Sonnenberg A, Hovnanian A. Spink5-deficient mice mimic Netherton syndrome through degradation of desmoglein 1 by epidermal protease hyperactivity. *Nat Genet* 2005; 37: 56-65.
- [33] Borst P, Elferink RO. Mammalian ABC transporters in health and disease. *Annu Rev Biochem* 2002; 71: 537-92.
- [34] Peelman F, Labeur C, Vanloo B, et al. Characterization of the ABCA transporter subfamily: identification of prokaryotic and eukaryotic members, phylogeny and

topology. *J Mol Biol* 2003; 325: 259-74.

- [35] Dale BA, Holbrook KA, Fleckman P, Kimball JR, Brumbaugh S, Sybert VP. Heterogeneity in harlequin ichthyosis, an inborn error of epidermal keratinization: variable morphology and structural protein expression and a defect in lamellar granules. *J Invest Dermatol* 1990; 94: 6-18.
- [36] Milner ME, O'Guin WM, Holbrook KA, Dale BA. Abnormal lamellar granules in harlequin ichthyosis. *J Invest Dermatol* 1992; 99: 824-9.
- [37] Akiyama M, Kim D-K, Main DM, Otto CE, Holbrook KA. Characteristic morphologic abnormality of harlequin ichthyosis detected in amniotic fluid cells. *J Invest Dermatol* 1994; 102: 210-3.
- [38] Akiyama M, Dale BA, Smith LT, Shimizu H, Holbrook KA. Regional difference in expression of characteristic abnormality of harlequin ichthyosis in affected fetuses. *Prenat Diagn* 1998; 18: 425-36.
- [39] Akiyama M, Yoneda K, Kim S-Y, Koyama H, Shimizu H. Cornified cell envelope proteins and keratins are normally distributed in harlequin ichthyosis. *J Cutan Pathol* 1996; 23: 571-5.
- [40] Akiyama M, Kim S-Y, Yoneda K, Shimizu H. Expression of transglutaminase 1 (transglutaminase K) in harlequin ichthyosis. *Arch Dermatol Res* 1997; 289: 116-9.
- [41] Virolainen E, Niemi K-M, Ganemo A, Kere J, Vahlquist A, Saarialho-Kere U. Ultrastructural features resembling those of harlequin ichthyosis in patients with severe congenital ichthyosiform erythroderma. *Br J Dermatol* 2001; 145: 480-3.
- [42] Kawashima J, Akiyama M, Takizawa Y, Takahashi S, Matsuo I, Shimizu H. Structural, enzymatic and molecular studies in a series of non-bullous congenital ichthyosiform erythroderma patients. *Clin Exp Dermatol* 2005; 30: 429-31.

- [43] Yamano G, Funahashi H, Kawanami O, et al. ABCA3 is a lamellar body membrane protein in human lung alveolar type II cells. *FEBS Lett* 2001; 508: 221-5.
- [44] Blanchet-Bardon C, Dumez Y, Labbe F, et al. Prenatal diagnosis of harlequin fetus. *Lancet* 1983; I (8316), 132.
- [45] Akiyama M, Suzumori K, Shimizu H. Prenatal diagnosis of harlequin ichthyosis by the examinations of keratinized hair canals and amniotic fluid cells at 19 weeks' estimated gestational age. *Prenat Diagn* 1999; 19: 167-71.
- [46] Tsuji-Abe Y, Akiyama M, Nakamura H, et al. DNA-based prenatal exclusion of bullous congenital ichthyosiform erythroderma at the early stage, 10-11 weeks' of pregnancy in two consequent siblings. *J Am Acad Dermatol* 2004; 51: 1008-11.
- [47] Schorderet DF, Huber M, Laurini RN, et al. Prenatal diagnosis of lamellar ichthyosis by direct mutational analysis of the keratinocyte transglutaminase gene. *Prenat Diagn* 1997; 17: 483-6.
- [48] Bichakjian CK, Nair RP, Wu WW, Goldberg S, Elder JT. Prenatal exclusion of lamellar ichthyosis based on identification of two new mutations in the transglutaminase 1 gene. *J Invest Dermatol* 1998; 110: 179-82.
- [49] Uitto J, Pfendner E, Jackson LG. Probing the fetal genome: progress in non-invasive prenatal diagnosis. *Trends Mol Med* 2003; 9, 339-43.
- [50] Wells D, Delhantry JDA. Preimplantation genetic diagnosis: Applications of molecular medicine. *Trends Mol Med* 2001; 7, 23-30.

Figure Legends

Figure 1. (a) A newborn patient with HI, harboring ABCA12 mutations. (b) A LI patient with transglutaminase 1 mutations in the neonatal period.

Figure 2. Hypothetical pathomechanisms of ARCI due to ABCA12 and transglutaminase 1 mutations: deficiency of intercellular lipid barrier in the stratum corneum.

(a) Schematic model showing the formation of the normal intercellular lipid layers in the stratum corneum. An intact cornified cell envelope is essential for the correct formation of intercellular lipid layers in the stratum corneum. ABCA12 works in lipid transport via lamellar granules to form an intercellular lipid coat. Transglutaminase 1 (TGase 1) crosslinks the cornified cell envelope precursor proteins.

(b) Schematic model of the malformation of stratum corneum barrier by ABCA12 defects. A loss of ABCA12 lipid transporter function results in abnormal lamellar granules and defective intercellular lipid layers.

(c) Schematic model of the malformation of intercellular lipid layers in the stratum corneum caused by an transglutaminase (TGase 1) enzyme deficiency. Defective cornified cell envelope formation also leads to the collapse of stratum corneum lipid barrier.

Table 1. Causative molecules or loci for ARCI includng HI, LI and NBCIE

Molecule	Gene (locus)	Type of Mutations	Phenotype [Reference]	Pathomechanism
ABCA12	<i>ABCA12</i> (2q34)	truncation/deletion/missense	HI [6]	severe deficiency of LG lipid transport
ABCA12	<i>ABCA12</i> (2q34)	missense	LI or NBCIE [5]	mild deficiency of LG lipid transport
transglutaminase 1	<i>TGMI</i> (14q11.2)	missense/deletion/insertion/truncation	LI [2,3] or NBCIE [28, 29]	defective CCE formation
lipoxygenase-3	<i>ALOXE3</i> (17p13.1)	missense/truncation	LI or NBCIE [15]	unknown (abnormal lipid metabolism in keratinocytes?)
12R-lipoxygenase	<i>ALOX12B</i> (17p13.1)	missense/truncation	LI or NBCIE [15]	unknown (abnormal lipid metabolism in keratinocytes?)
ichthyin	<i>ichthyin</i> (5q33)	missense/truncation	NBCIE or LI [12]	unknown
unknown	unknown (3p21)	---	NBCIE [10]	unknown
unknown	unknown (19p12-q12)	---	LI [10]	unknown
unknown	unknown (12p11.2-q13)	---	NBCIE [13]	unknown
unknown	unknown (19p13.2-p13.1)	---	NNCI [9]	unknown

ARCI, autosomal recessive congenital ichthyosis; HI, harlequin ichthyosis; LI, lamellar ichthyosis; NBCIE, non-bullous congenital ichthyosiform erythroderma; NNCI, non-lamellar non-erythrodermic congenital ichthyosis; LG, lamellar granule; CCE, cornified cell envelope

(a)

(b)

(c)