Title	Generalized Weak Weyl Relation and Decay of Quantum Dynamics
Author(s)	Arai, Asao
Citation	Reviews in Mathematical Physics, 17(9), 1071-1109 https://doi.org/10.1142/S0129055X05002479
Issue Date	2005
Doc URL	http://hdl.handle.net/2115/1390
Type	article (author version)
Note	Reprinted from Publication, Reviews in Mathematical Physics, Vol 17-9, Arai, A, Generalized Weak Weyl Relation and Decay of Quantum Dynamics, p.1071-1109, c2005, with permission from World Scientific Publishing Co. Pte. Ltd, Singapore
File Information	RMP17-9.pdf

[Instructions for use](#)

Generalized Weak Weyl Relation and Decay of Quantum Dynamics

Asao Arai

Department of Mathematics, Hokkaido University

Sapporo 060-0810, Japan

E-mail: arai@math.sci.hokudai.ac.jp

January 30, 2006

This paper was published in *Reviews in Mathematical Physics* Vol. 17 (2005), 1071-1109.

Abstract

Let H be a self-adjoint operator on a Hilbert space $\mathcal{H}$, T be a symmetric operator on $\mathcal{H}$ and $K(t)$ ($t \in \mathbb{R}$) be a bounded self-adjoint operator on $\mathcal{H}$. We say that (T, H, K) obeys the *generalized weak Weyl relation* (GWWR) if $e^{-itH}D(T) \subset D(T)$ for all $t \in \mathbb{R}$ and $Te^{-itH}\psi = e^{-itH}(T + K(t))\psi, \forall \psi \in D(T)$ ($D(T)$ denotes the domain of T). In the context of quantum mechanics where H is the Hamiltonian of a quantum system, we call T a *generalized time operator* of H . We first investigate, in an abstract framework, mathematical structures and properties of triples (T, H, K) obeying the GWWR. These include the absolute continuity of the spectrum of H restricted to a closed subspace of $\mathcal{H}$, an uncertainty relation between H and T (a “time-energy uncertainty relation”), the decay property of transition probabilities $\left| \langle \psi, e^{-itH} \phi \rangle \right|^2$ as $|t| \rightarrow \infty$ for all vectors ψ and ϕ in a subspace of $\mathcal{H}$, where $\langle \cdot, \cdot \rangle$ denotes the inner product of $\mathcal{H}$. We describe methods to construct various examples of triples (T, H, K) obeying the GWWR. In particular we show that there exist generalized time operators of second quantization operators on Fock spaces (full Fock spaces, boson Fock spaces, fermion Fock spaces) which may have applications to quantum field models with interactions.

Keywords: Generalized weak Weyl relation; time operator; canonical commutation relation; Hamiltonian; quantum dynamics; survival probability; decay in time; time-energy uncertainty relation; Schrödinger operator; Dirac operator; Fock space; second quantization

Contents

1	Introduction	2
2	Fundamental Properties of the GWWR	5
2.1	Elementary facts	5
2.2	Nonself-adjointness of generalized time operators	7
2.3	Construction of triples obeying the GWWR in direct sums of Hilbert spaces	9
2.4	Perturbations	10
3	Transition Probability Amplitudes and the Point Spectra of Hamiltonians	11
4	Generalized Weak CCR and Time-Energy Uncertainty Relations	12
5	The Point Spectra of Generalized Time Operators	14
6	Commutation Formulas and Absolute Continuity	14
6.1	General cases	15
6.2	A special case	16
7	Absence of Minimum-Uncertainty States	19
8	Power Decays of Transition Probability Amplitudes in Quantum Dynamics	21
8.1	A simple case	22
8.2	Higher order dcays in smaller subspaces	22
8.3	Correlation functions	25
8.4	Heat semi-groups	27
9	Abstract Version of Wigner's Time-Energy Uncertainty Relation	27
10	Structure producing successively triples obeying the GWWR	29
11	Generalized Time Operators of Partial Differential Operators	31
11.1	Constructions from the Schrödinger representation of the CCR with d degrees of freedom	31
11.2	Abstract Dirac operators	34
12	Representations of the GWWR in Fock Spaces	35
12.1	Tensor Representations of the GWWR	35
12.2	Constructions of triples obeying the GWWR in Fock Spaces	36

1 Introduction

In this paper we develop, in an abstract framework, an operator theory of a commutation relation, which is a generalization of a variant of the canonical commutation relation

(CCR) with one degree of freedom, and put a basis for applications of the theory to quantum mechanics and quantum field theory.

To explain new features of the present work, we first recall some of the basic aspects on the representation theory of the CCR. As is well-known, a representation of the CCR with one degree of freedom is defined to be a triple $(\mathcal{H}, \mathcal{D}, (Q, P))$ consisting of a complex Hilbert space $\mathcal{H}$, a dense subspace $\mathcal{D}$ of $\mathcal{H}$ and a pair (Q, P) of symmetric operators on $\mathcal{H}$ such that $\mathcal{D} \subset D(QP) \cap D(PQ)$ ($D(\cdot)$ denotes operator domain) and

$$QP - PQ = iI \quad (1.1)$$

on $\mathcal{D}$, where $i := \sqrt{-1}$ and I denotes the identity on $\mathcal{H}^1$. If both Q and P are self-adjoint and $\mathcal{D}$ is a common core of Q and P , then we say that the representation $(\mathcal{H}, \mathcal{D}, (Q, P))$ is self-adjoint².

A typical example of self-adjoint representations of the CCR is the Schrödinger representation $(L^2(\mathbb{R}), C_0^\infty(\mathbb{R}), (q, p))$ with q being the multiplication operator by the function $x \in \mathbb{R}$ acting in $L^2(\mathbb{R})$ and $p := -iD_x$ the generalized differential operator in the variable x acting in $L^2(\mathbb{R})$.

There is a stronger form of representation of the CCR: A double $(\mathcal{H}, (Q, P))$ consisting of a complex Hilbert space $\mathcal{H}$ and a pair (Q, P) of self-adjoint operators on $\mathcal{H}$ is called a Weyl representation of the CCR with one degree of freedom if

$$e^{itQ}e^{isP} = e^{-its}e^{isP}e^{itQ}, \quad \forall t, s \in \mathbb{R}.$$

This relation is called the Weyl relation (e.g., [19, pp.274–275]). It is easy to see that the Schrödinger representation $(L^2(\mathbb{R}), (q, p))$ is a Weyl representation. Von Neumann [16] proved that every Weyl representation on a separable Hilbert space is unitarily equivalent to a direct sum of the Schrödinger representation. This theorem—the von Neumann uniqueness theorem—implies that each Weyl representation of the CCR is a self-adjoint representation of the CCR (for details, see, e.g., [6, 18]). But *a self-adjoint representation of the CCR is not necessarily a Weyl representation of the CCR*, namely there are self-adjoint representations of the CCR that are *not* Weyl representations (e.g., [8]). Physically interesting examples of self-adjoint representations of the CCR's with two degrees of freedom which are not necessarily unitarily equivalent to the Schrödinger representation

¹One can generalize the concept of the representation of the CCR by taking the commutation relation (1.1) in the sense of sesquilinear form, i.e., $\mathcal{D} \subset D(Q) \cap D(P)$ and $\langle Q\psi, P\phi \rangle - \langle P\psi, Q\phi \rangle = i \langle \psi, \phi \rangle$, $\psi, \phi \in \mathcal{D}$, where $\langle \cdot, \cdot \rangle$ denotes the inner product of $\mathcal{H}$.

²There are representations of the CCR which cannot be self-adjoint. For example, consider the Hilbert space $L^2(\mathbb{R}_+)$ with $\mathbb{R}_+ := (0, \infty)$ and define operators q_+, p_+ on $L^2(\mathbb{R}_+)$ as follows:

$$\begin{aligned} D(q_+) &:= \left\{ f \in L^2(\mathbb{R}_+) \mid \int_{\mathbb{R}_+} |rf(r)|^2 dr < \infty \right\}, \quad (q_+f)(r) := rf(r), \quad f \in D(q_+), \quad \text{a.e. } r \in \mathbb{R}_+, \\ D(p_+) &:= C_0^\infty(\mathbb{R}_+), \quad (p_+f)(r) := -i \frac{df(r)}{dr}, \quad f \in D(p_+), \quad \text{a.e. } r \in \mathbb{R}_+. \end{aligned}$$

Then q_+ is self-adjoint, p is symmetric and $(L^2(\mathbb{R}_+), C_0^\infty(\mathbb{R}_+), (q_+, p_+))$ is a representation of the CCR with one degree of freedom. It is not so difficult to prove that p_+ has no self-adjoint extensions (e.g., see [6, Chapter 2, Example D.1]). Therefore (q_+, p_+) cannot be extended to a self-adjoint representation of the CCR on $L^2(\mathbb{R}_+)$.

of the CCR's appear in two-dimensional gauge quantum mechanics with singular gauge potentials. These representations, which are closely related to the so-called Aharonov-Bohm effect [1], have been studied extensively by the present author in a series of papers (see [5] and references therein).

Schmüdgen [21] presented and studied a weaker version of the Weyl relation with one degree of freedom: Let T be a symmetric operator and H be a self-adjoint operator on a Hilbert space $\mathcal{H}$. We say that (T, H) obeys the *weak Weyl relation* (WWR) if $e^{-itH}D(T) \subset D(T)$ for all $t \in \mathbb{R}$ and

$$Te^{-itH}\psi = e^{-itH}(T+t)\psi, \quad \forall \psi \in D(T), \forall t \in \mathbb{R},$$

where, for later convenience, we use the symbols (T, H) instead of (Q, P) . We call $(\mathcal{H}, (T, H))$ a weak Weyl representation of the CCR with one degree of freedom. It is easy to see that every Weyl representation of the CCR is a weak Weyl representation of the CCR. But *the converse is not true* [21]³. It should be remarked also that the WWR implies the CCR, but *a representation of the CCR is not necessarily a weak Weyl representation of the CCR*. In this sense the WWR is between the CCR and the Weyl relation.

The WWR was used to study a time operator with application to survival probabilities in quantum dynamics [11, 12] (in the article [11], the WWR is called the T -weak Weyl relation), where H is taken to be the Hamiltonian of a quantum system. It was proven in [11] that, if (T, H) obeys the WWR, then H has no point spectrum and its spectrum is purely absolutely continuous [11, Corollary 4.3, Theorem 4.4]. This kind class of H , however, is somewhat restrictive. From this point of view, it would be natural to investigate a general version of the WWR (if any) such that H is not necessarily purely absolutely continuous. This is one of the motivations of the present work.

The general version of the WWR we consider in the present paper is defined as follows:

Definition 1.1 Let T be a symmetric operator on a Hilbert space $\mathcal{H}$, H be a self-adjoint operator on $\mathcal{H}$ and $K(t)$ ($t \in \mathbb{R}$) be a bounded self-adjoint operator on $\mathcal{H}$ with $D(K(t)) = \mathcal{H}$, $\forall t \in \mathbb{R}$. We say that (T, H, K) obeys the *generalized weak Weyl relation* (GWWR) in $\mathcal{H}$ if $e^{-itH}D(T) \subset D(T)$ for all $t \in \mathbb{R}$ and

$$Te^{-itH}\psi = e^{-itH}(T + K(t))\psi, \quad \forall \psi \in D(T), \forall t \in \mathbb{R}. \quad (1.2)$$

We call the operator-valued function K the commutation factor in the GWWR. Also we sometimes say that (T, H, K) is a representation of the GWWR.

Obviously the case $K(t) = t$ in the GWWR gives the WWR. Hence the GWWR is certainly a generalization of the WWR. One can show also that the GWWR implies a generalized version of the CCR (Proposition 4.3 in the present paper). Since the (G)WWR is a weaker version of the Weyl relation, the strong properties arising from the Weyl relation (e.g., spectral properties) may be weakened by the (G)WWR. It is very interesting to investigate this aspect. Thus triples (T, H, K) obeying the GWWR become

³The pair $(p_+, -q_+)$ in the preceding footnote obeys the WWR. But it cannot be a Weyl representation, since p_+ is not self-adjoint (or $-q_+$ is nonpositive).

the main objects of the present paper. As suggested above, in applications to quantum mechanics and quantum field theory, we have in mind the case where H is the Hamiltonian of a quantum system. In this realization of H , we call T a *generalized time operator*. We show that the GWR implies a “time-energy uncertainty relation” between H and T (for physical discussions related to this aspect, see [15] and references therein). Mathematically rigorous studies for time-energy uncertainty relations, which, however, do not use time operators, are given in [17]. In the present paper we construct generalized time operators for Hamiltonians in both relativistic and nonrelativistic quantum mechanics including Dirac type operators as well as in quantum field theory.

The outline of the present paper is as follows. In Section 2 we discuss fundamental properties of representations (T, H, K) of the GWR. In Section 3 we derive a decay property (in time $t \in \mathbb{R}$) of transition probability amplitudes $\langle \psi, e^{-itH} \phi \rangle$ for vectors ψ, ϕ in a suitable subspace, where $\langle \cdot, \cdot \rangle$ denotes the inner product of $\mathcal{H}$. In Section 4, we introduce a concept of the generalized weak CCR and show that the GWR implies the generalized weak CCR. We derive also a time-energy uncertainty relation. Section 5 is concerned with properties of the point spectrum of T . In Section 6 we develop functional calculus for the GWR. In a special case where $K(t) = K_C(t) := tC$ with C a bounded self-adjoint operator, we prove the absolute continuity of H restricted to the closure of $\text{Ran}(C)$, the range of C . In Section 7, we prove *absence* of minimum-uncertainty states for each representation (T, H, K_C) with T being closed and H being bounded from below. This is an interesting aspect in the sense that it shows a difference from the Weyl representation (on a separable Hilbert space) or the Fock representation of the CCR in which minimum-uncertainty states exist (see Remark 7.1 for more details). In Section 8, for representations (T, H, K_C) of the GWR, we derive power laws for decays (in time) of transition probability amplitudes as well as two-point correlation functions. Heat semi-groups $e^{-\beta H}$ ($\beta > 0$) generated by H (under the condition that H is bounded from below) are also considered. In Section 9 we present an abstract version of Wigner’s time-energy uncertainty relation [23]. In Section 10, we show that there exists a structure producing successively representations of the GWR. In Section 11, we construct concrete classes of representations of the GWR, using partial differential operators acting in $L^2(\mathbb{R}^d)$ ($d \in \mathbb{N}$). We also find generalized time operators for an abstract Dirac operator. In the last section we present a tensor representation of the GWR and construct generalized time operators for second quantization operators on Fock spaces (full Fock spaces, boson Fock spaces, fermion Fock spaces). This puts a basis to investigations of quantum field models with interactions.

2 Fundamental Properties of the GWR

Throughout this section, we assume that (T, H, K) obeys the GWR in a Hilbert space $\mathcal{H}$ (Definition 1.1). We denote the inner product and the norm of $\mathcal{H}$ by $\langle \cdot, \cdot \rangle$ and $\| \cdot \|$ respectively.

2.1 Elementary facts

Proposition 2.1 For all $t \in \mathbb{R}$, $e^{-itH}D(T) = D(T)$ and the operator equality

$$Te^{-itH} = e^{-itH}(T + K(t)) \quad (2.1)$$

holds. Moreover

$$K(0) = 0. \quad (2.2)$$

Proof: Taking $-t$ as the t in Definition 1.1, we have $e^{itH}D(T) \subset D(T)$ for all $t \in \mathbb{R}$. Hence $D(T) \subset e^{-itH}D(T) \subset D(T)$ for all $t \in \mathbb{R}$, which implies that $e^{-itH}D(T) = D(T)$ for all $t \in \mathbb{R}$. By definition, we have $e^{-itH}(T + K(t)) \subset Te^{-itH}$ for all $t \in \mathbb{R}$. On the other hand, the preceding result implies that $D(Te^{-itH}) = D(T)$ for all $t \in \mathbb{R}$. Thus (2.1) follows. Letting $t = 0$ in (2.1), we have $K(0) = 0$ on $D(T)$. Since $D(T)$ is dense and $K(0)$ is bounded, we obtain (2.2). ■

Proposition 2.2 Let $\bar{T}$ be the closure of T . Then $(\bar{T}, H, K)$ obeys the GWWR.

Proof: For each $\psi \in D(\bar{T})$, there exists a sequence $\{\psi_n\}_{n=1}^\infty \subset D(T)$ such that $\lim_{n \rightarrow \infty} \psi_n = \psi$ and $\lim_{n \rightarrow \infty} T\psi_n = \bar{T}\psi$. We have $Te^{-itH}\psi_n = e^{-itH}(T + K(t))\psi_n$. The right hand side (r.h.s.) strongly converges to $e^{-itH}(\bar{T} + K(t))\psi$ as $n \rightarrow \infty$. We have $e^{-itH}\psi_n \rightarrow e^{-itH}\psi$ as $n \rightarrow \infty$. Hence $e^{-itH}\psi \in D(\bar{T})$ and $\bar{T}e^{-itH}\psi = e^{-itH}(\bar{T} + K(t))\psi$. Thus $(\bar{T}, H, K)$ obeys the GWWR. ■

For a linear operator A , we denote by $\sigma(A)$ (resp. $\sigma_p(A)$) the spectrum (resp. the point spectrum) of A .

Corollary 2.3 For all $t \in \mathbb{R}$, $\sigma(T + K(t)) = \sigma(T)$ and $\sigma_p(T + K(t)) = \sigma_p(T)$, where the multiplicity of each $\lambda \in \sigma_p(T)$ is equal to that of $\lambda \in \sigma_p(T + K(t))$.

Proof: Operator equality (2.1) means the unitary equivalence of $T + K(t)$ and T . Hence, by a general theorem, the desired results follow. ■

Definition 2.4 We say that a linear operator L on $\mathcal{H}$ strongly commutes with H if $e^{-itH}D(L) \subset D(L)$ for all $t \in \mathbb{R}$ and $e^{-itH}L \subset Le^{-itH}$.

Remark 2.1 In the same way as in the proof of Proposition 2.1, one can show that L strongly commutes with H if and only if operator equality $e^{-itH}L = Le^{-itH}$ holds for all $t \in \mathbb{R}$.

Proposition 2.5 Let S be a symmetric operator on $\mathcal{H}$ strongly commuting with H such that $D(S) \cap D(T)$ is dense (hence $T + S$ is a symmetric operator with $D(T + S) := D(T) \cap D(S)$). Then $(T + S, H, K)$ obeys the GWWR.

Proof: A simple calculation. ■

We denote by $\mathcal{B}(\mathcal{H})$ the Banach space of all bounded linear operators on $\mathcal{H}$ with domains equal to $\mathcal{H}$.

Proposition 2.6 For all $t \in \mathbb{R}$,

$$e^{itH}K(-t) + K(t)e^{itH} = 0. \quad (2.3)$$

In particular

$$\sigma(K(t)) = \sigma(-K(-t)), \quad \sigma_p(K(t)) = \sigma_p(-K(-t)), \quad \forall t \in \mathbb{R}. \quad (2.4)$$

Proof : Let $t \in \mathbb{R}$. In general, for all $W \in \mathcal{B}(\mathcal{H})$ and every densely defined linear operator A on $\mathcal{H}$, we have $(WA)^* = A^*W^*$ (operator equality). Hence, by taking the adjoint of (2.1), we have $e^{itH}T^* \subset (T^* + K(t))e^{itH}$ for all $t \in \mathbb{R}$. Hence, for all $\psi \in D(T)$,

$$e^{itH}T\psi = (T + K(t))e^{itH}\psi = e^{itH}(T + K(-t))\psi + K(t)e^{itH}\psi,$$

where we have used (2.1) to obtain the second equality. This implies that $e^{itH}K(-t)\psi + K(t)e^{itH}\psi = 0$ for all $t \in \mathbb{R}$. Since $D(T)$ is dense, we obtain (2.3). Operator equality (2.3) implies the unitary equivalence of $K(t)$ and $-K(-t)$. Hence (2.4) follows. ■

2.2 Nonself-adjointness of generalized time operators

In this subsection we prove the following theorem:

Theorem 2.7 Assume that $K : \mathbb{R} \rightarrow \mathcal{B}(\mathcal{H})$ is strongly differentiable on $\mathbb{R}$ and let

$$K'(t) := s\text{-}\frac{dK(t)}{dt}, \quad (2.5)$$

the strong derivative of K in $t \in \mathbb{R}$. Suppose that $K'(0) \neq 0$, H is semi-bounded (i.e., bounded from below or bounded from above) and

$$K(t)T \subset TK(t) \quad (2.6)$$

for all $t \in \mathbb{R}$. Then T is not self-adjoint.

Remark 2.2 In the simple case $K(t) = t$, the fact stated in Theorem 2.7 has been pointed out in [11].

We need some preliminary results.

Proposition 2.8 Suppose that T is self-adjoint. Then, for all $t \in \mathbb{R}$, $T + K(t)$ is self-adjoint and

$$e^{-isT}e^{-itH} = e^{-itH}e^{-is(T+K(t))}, \quad \forall s, \forall t \in \mathbb{R}. \quad (2.7)$$

Proof : The self-adjointness of $T + K(t)$ follows from a simple application of the Kato-Rellich theorem, since $K(t)$ is bounded and self-adjoint. By (2.1), we have

$$e^{itH}Te^{-itH} = T + K(t) \quad (2.8)$$

as operator equality. Hence, by the functional calculus, we have for all $s, t \in \mathbb{R}$

$$e^{itH}e^{-isT}e^{-itH} = e^{-is(T+K(t))},$$

which is equivalent to (2.7). ■

Definition 2.9 We say that two self-adjoint operators A and B on a Hilbert space strongly commute if their spectral measures commute : $E_A(J_1)E_B(J_2) = E_B(J_2)E_A(J_1)$ for all Borel sets J_1, J_2 in $\mathbb{R}$, where E_A (resp. E_B) denotes the spectral measure of A (resp. B).

For characterizations of the strong commutativity of two self-adjoint operators, we refer the reader to [19, Theorem VIII.13].

Lemma 2.10 *Let A and B be self-adjoint operators on a Hilbert space. Suppose that B is bounded and $BA \subset AB$. Then A and B strongly commute.*

Proof : The assumption implies that $BD(A) \subset D(A)$ and $BA\psi = AB\psi$, $\forall \psi \in D(A)$. This implies that, for all $n \in \mathbb{N}$ and $\psi \in D(A)$, $B^n\psi \in D(A)$ and $B^nA\psi = AB^n\psi$. For $N \in \mathbb{N}$ and $t \in \mathbb{R}$, we set $e_N := \sum_{n=0}^N (it)^n B^n / n!$. let $\psi \in D(A)$. Then $e_N A\psi = A e_N \psi$. It is easy to see that, for all $\phi \in \mathcal{H}$, $e_N \phi \rightarrow e^{itB} \phi$ as $N \rightarrow \infty$. Hence $A e_N \psi \rightarrow e^{itB} A\psi$ as $N \rightarrow \infty$. Since A is closed, it follows that $e^{itB} \psi \in D(A)$ and $A e^{itB} \psi = e^{itB} A\psi$. In particular, $e^{itB} D(A) \subset D(A)$ for all $t \in \mathbb{R}$ and hence, in fact, $e^{itB} D(A) = D(A)$. Therefore operator equality $e^{itB} A e^{-itB} = A$ follows. By the functional calculus, we have $e^{itB} e^{isA} e^{-itB} = e^{isA}$, $s, t \in \mathbb{R}$. This implies the strong commutativity of A and B (apply [19, Theorem VIII.13]). ■

Lemma 2.11 *Let A and B be strongly commuting self-adjoint operators on a Hilbert space. Then $A + B$ is essentially self-adjoint and, for all $t \in \mathbb{R}$,*

$$e^{it\overline{(A+B)}} = e^{itA} e^{itB} = e^{itB} e^{itA}. \quad (2.9)$$

Proof : The essential self-adjointness of $A + B$ follows from the two variable functional calculus (note that $\overline{A+B} = \int_{\mathbb{R}^2} (\lambda + \mu) dE_{A,B}(\lambda, \mu)$, where $E_{A,B}$ is the two dimensional spectral measure such that $E_{A,B}(J_1 \times J_2) = E_A(J_1)E_B(J_2)$ for all Borel sets J_1, J_2 in $\mathbb{R}$). Formula (2.9) follows from the Trotter product formula [19, Theorem VIII.31] and [19, Theorem VIII.13(c)]. ■

Proof of Theorem 2.7

Suppose that T were self-adjoint. Then it follows from (2.6) and Lemma 2.10 that $K(t)$ and T strongly commute. Hence, by (2.7) and Lemma 2.11, we have for all $s \in \mathbb{R}, t \in \mathbb{R} \setminus \{0\}$

$$\left(\frac{e^{-itH} - 1}{t} \right) e^{-isT} \psi = e^{-isT} \left(\frac{e^{-itH} - 1}{t} \right) \psi - e^{-itH} e^{-isT} \left(\frac{e^{-isK(t)} - 1}{t} \right) \psi \quad (2.10)$$

for all $\psi \in \mathcal{H}$. We can write

$$\frac{e^{-isK(t)} - 1}{t} = -is \frac{K(t)}{t} + M_s(t)$$

with

$$M_s(t) := \sum_{n=2}^{\infty} \frac{(-is)^n}{n!} \frac{K(t)}{t} K(t)^{n-1}$$

in the operator norm topology. By the strong differentiability of K with $K(0) = 0$ (Proposition 2.1) and the principle of uniform boundedness, we have for each $\delta > 0$

$$C_\delta := \sup_{|t| < \delta} \|K(t)/t\| < \infty, \quad D_\delta := \sup_{|t| < \delta} \|K(t)\| < \infty,$$

where, for a bounded operator $A \in \mathcal{B}(\mathcal{H})$, $\|A\|$ denotes the operator norm of A . Hence, for all $\psi \in \mathcal{H}$ and $0 < |t| < \delta$,

$$\|M_s(t)\psi\| \leq C_\delta \left(\sum_{n=2}^{\infty} \frac{|s|^n}{n!} D_\delta^{n-2} \right) \|K(t)\psi\|.$$

Using this estimate and the fact that $K(0) = 0$, we obtain $\lim_{t \rightarrow 0} M_s(t)\psi = 0$. Hence

$$\lim_{t \rightarrow 0} \frac{e^{-isK(t)} - 1}{t} \psi = -isK'(0)\psi.$$

Let $\psi \in D(H)$. Then, by (2.10),

$$\lim_{t \rightarrow 0} \left(\frac{e^{-itH} - 1}{t} \right) e^{-isT} \psi = e^{-isT} (-iH)\psi - e^{-isT} (-is)K'(0)\psi,$$

which implies that $e^{-isT}\psi \in D(H)$ and

$$(-iH)e^{-isT}\psi = e^{-isT}(-iH)\psi - e^{-isT}(-is)K'(0)\psi,$$

i.e.,

$$e^{isT}He^{-isT}\psi = H\psi - sK'(0)\psi.$$

Hence

$$\langle e^{-isT}\psi, He^{-isT}\psi \rangle = \langle \psi, H\psi \rangle - s \langle \psi, K'(0)\psi \rangle. \quad (2.11)$$

Since $K'(0)$ is a non-zero bounded self-adjoint operator by the present assumption, there exists a vector $\eta \in D(H)$ such that $\|\eta\| = 1$ and $\langle \eta, K'(0)\eta \rangle \neq 0$. Equation (2.11) implies that

$$\sup_{s \in \mathbb{R}} \langle e^{-isT}\eta, He^{-isT}\eta \rangle = \infty, \quad \inf_{s \in \mathbb{R}} \langle e^{-isT}\eta, He^{-isT}\eta \rangle = -\infty.$$

Hence H is not semi-bounded (note that $\|e^{-isT}\eta\| = 1$ for all $s \in \mathbb{R}$). Thus we are led to a contradiction. $\blacksquare$

2.3 Construction of triples obeying the GWWR in direct sums of Hilbert spaces

Let $\mathcal{H}_1$ be a Hilbert space and $\mathcal{F} := \mathcal{H} \oplus \mathcal{H}_1$. Let (T_1, H_1, K_1) be a triple obeying the GWWR in $\mathcal{H}_1$. We define

$$\widetilde{H} := H \oplus H_1 = \begin{pmatrix} H & 0 \\ 0 & H_1 \end{pmatrix}. \quad (2.12)$$

Proposition 2.12 *Let A be a bounded linear operator from $\mathcal{H}$ to $\mathcal{H}_1$ with $D(A) = \mathcal{H}$ and*

$$\tilde{T} := \begin{pmatrix} T & A^* \\ A & T_1 \end{pmatrix}, \quad \tilde{K}(t) := \begin{pmatrix} K(t) & e^{itH} A^* e^{-itH_1} - A^* \\ e^{itH_1} A e^{-itH} - A & K_1(t) \end{pmatrix}. \quad (2.13)$$

Then $(\tilde{T}, \tilde{H}, \tilde{K})$ obeys the GWWR in $\mathcal{F}$.

Proof: By the functional calculus, we have $e^{-it\tilde{H}} = e^{-itH} \oplus e^{-itH_1}$ for all $t \in \mathbb{R}$. Then direct computations yield the desired result. $\blacksquare$

Note that, in Proposition 2.12, $\tilde{T}$ is not diagonal if $A \neq 0$. This procedure of construction of a new triple obeying the GWWR obviously yields an algorithm to obtain a triple obeying the GWWR in the N direct sum $\oplus_{n=1}^N \mathcal{H}_n$ of Hilbert spaces $\mathcal{H}_n$ ($N \geq 2$), provided that, for each n , a triple (T_n, H_n, K_n) obeying the GWWR in $\mathcal{H}_n$ is given.

2.4 Perturbations

Let V be a symmetric operator on $\mathcal{H}$ and assume that

$$H(V) := H + V \quad (2.14)$$

is essentially self-adjoint. It is an interesting problem to investigate if there exist a symmetric operator T_V and an operator-valued function $K_V : \mathbb{R} \rightarrow \mathcal{B}(\mathcal{H})$ such that $(T_V, \overline{H(V)}, K_V)$ obeys the GWWR. Here we present only an abstract answer to this problem.

Proposition 2.13 *Assume that the following conditions (i)–(iii) hold:*

- (i) *The operators T, H and $K(t)$ ($t \in \mathbb{R}$) are reduced by a closed subspace $\mathcal{M}$ of $\mathcal{H}$. We denote their reduced part by $T_{\mathcal{M}}, H_{\mathcal{M}}$ and $K_{\mathcal{M}}(t)$ respectively.*
- (ii) *The operator $\overline{H(V)}$ is reduced by a closed subspace $\mathcal{N}$ of $\mathcal{H}$.*
- (iii) *There exists a unitary operator $U : \mathcal{M} \rightarrow \mathcal{N}$ such that $U H_{\mathcal{M}} U^{-1} = \overline{H(V)}_{\mathcal{N}}$.*

Let

$$T_V := (U T_{\mathcal{M}} U^{-1}) \oplus 0, \quad K_V(t) := (U K_{\mathcal{M}}(t) U^{-1}) \oplus 0 \quad (2.15)$$

relative to the orthogonal decomposition $\mathcal{H} = \mathcal{N} \oplus \mathcal{N}^\perp$. Then $(T_V, \overline{H(V)}, K_V)$ obeys the GWWR.

Proof: It is obvious that T_V is symmetric and $K_V(t)$ is a bounded self-adjoint operator. By direct computations, one sees that $(T_V, \overline{H(V)}, K_V)$ obeys the GWWR. $\blacksquare$

Remark 2.3 A method to find the unitary operator U in Proposition 2.13 is to use the method of wave operators with respect to the pair $(H, \overline{H(V)})$. In that case, U would be one of the wave operators $W_\pm := s\text{-}\lim_{t \rightarrow \pm\infty} e^{itH(V)} J e^{-itH} P_{\text{ac}}(H)$ (if they exist) ($P_{\text{ac}}(H)$ is the orthogonal projection onto the absolutely continuous space of H and J is a linear operator), $\mathcal{M} = (\ker W_\pm)^\perp$ and $\mathcal{N} = \text{Ran}(W_\pm)$ (e.g., [9, §4.2], [20, p.34, Proposition 4]). This method was taken in [11, 12] in the case where H is the 1-dimensional Laplacian and V is a real-valued function on $\mathbb{R}$ (hence $H(V)$ is a one-dimensional Schrödinger operator). In the present paper we do not develop this aspect.

3 Transition Probability Amplitudes and the Point Spectra of Hamiltonians

If a self-adjoint operator H on a Hilbert space represents the Hamiltonian of a quantum system, then the transition probability amplitude of an initial state $\psi \in \mathcal{H}$ with $\|\psi\| = 1$ to a state $\phi \in \mathcal{H}$ with $\|\phi\| = 1$ at time $t \in \mathbb{R}$ is given by $\langle \phi, e^{-itH}\psi \rangle$. The square $|\langle \phi, e^{-itH}\psi \rangle|^2$ of its modulus is called the transition probability from ψ at time 0 to ϕ at time t . In particular $|\langle \psi, e^{-itH}\psi \rangle|^2$ is referred to as the survival probability, at time t , of the state ψ .

Let (T, H, K) be a triple obeying the GWWR in a Hilbert space $\mathcal{H}$. The following proposition is concerned with upper bounds of the modulus of a transition probability amplitude in time t .

Proposition 3.1 *Suppose that there is a constant $\alpha > 0$ such that the strong limit*

$$L_\alpha := \text{s-}\lim_{t \rightarrow \infty} \frac{K(t)}{t^\alpha} \in \mathcal{B}(\mathcal{H}) \quad (3.1)$$

exists. Let S be a symmetric operator strongly commuting with H . Then, for all $\psi, \phi \in D(T) \cap D(S)$ and $t > 0$,

$$|\langle \psi, e^{-itH} L_\alpha \phi \rangle| \leq \frac{\|(T+S)\psi\| \|\phi\| + \|\psi\| \|(T+S)\phi\|}{t^\alpha} + \|\psi\| \left\| \left(L_\alpha - \frac{K(t)}{t^\alpha} \right) \phi \right\|. \quad (3.2)$$

Proof: Let $L = L_\alpha$ and $L(t) := K(t)/t^\alpha$. Then

$$\begin{aligned} |\langle \psi, e^{-itH} L \phi \rangle| &\leq |\langle \psi, e^{-itH} (L - L(t)) \phi \rangle| + |\langle \psi, e^{-itH} L(t) \phi \rangle| \\ &\leq \|\psi\| \|(L - L(t)) \phi\| + |\langle \psi, e^{-itH} L(t) \phi \rangle|. \end{aligned}$$

By Proposition 2.5, we have $e^{-itH} L(t) \phi = t^{-\alpha} [(T+S)e^{-itH} \phi - e^{-itH} (T+S) \phi]$. Using this relation, we have

$$|\langle \psi, e^{-itH} L(t) \phi \rangle| \leq \frac{\|(T+S)\psi\| \|\phi\| + \|\psi\| \|(T+S)\phi\|}{t^\alpha}.$$

Hence (3.2) follows. ■

Remark 3.1 Proposition 3.1 is a generalization of [11, Theorem 4.1] where the special case $K(t) = t$ is considered.

The following corollary is a generalized version of [11, Corollary 4.3]:

Corollary 3.2 *Suppose that the assumption of Proposition 3.1 holds. Then, for all $\psi, \phi \in \mathcal{H}$,*

$$\lim_{t \rightarrow \infty} \langle \psi, e^{-itH} L_\alpha \phi \rangle = 0. \quad (3.3)$$

Proof: The proof is similar to that of [11, Corollary 4.3]. By the polarization identity, we need only to prove (3.3) with $\phi = \psi$. Since $D(T)$ is dense, there exists a sequence $\psi_n \in D(T)$ such that $\lim_{n \rightarrow \infty} \psi_n = \psi$. Then, in the same way as in the proof of the preceding proposition, we have

$$\begin{aligned} \left| \langle \psi, e^{-itH} L_\alpha \psi \rangle \right| &\leq \|\psi - \psi_n\| \|L_\alpha \psi\| + \|\psi_n\| \|L_\alpha\| \|\psi - \psi_n\| \\ &\quad + \left| \langle \psi_n, e^{-itH} L_\alpha \psi_n \rangle \right|. \end{aligned}$$

By (3.2) with $\psi = \phi = \psi_n$ and $S = 0$, we have $\lim_{t \rightarrow \infty} \left| \langle \psi_n, e^{-itH} L_\alpha \psi_n \rangle \right| = 0$. Hence

$$\limsup_{t \rightarrow \infty} \left| \langle \psi, e^{-itH} L_\alpha \psi \rangle \right| \leq \|\psi - \psi_n\| \|L_\alpha \psi\| + \|\psi_n\| \|L_\alpha\| \|\psi - \psi_n\|.$$

Then, taking $n \rightarrow \infty$, we obtain (3.3) with $\phi = \psi$. ■

This corollary implies an interesting structure of the point spectrum of H :

Corollary 3.3 *Suppose that the assumption of Proposition 3.1 holds. Then, for all $E \in \mathbb{R}$, $\ker(H - E) \subset \ker L_\alpha$. In particular, if $\ker L_\alpha = \{0\}$, then $\sigma_p(H) = \emptyset$.*

Proof: Let $\psi_E \in \ker(H - E)$. Then $e^{itH} \psi_E = e^{itE} \psi_E$. Taking $\psi = \psi_E$ in (3.3), we obtain $\langle \psi_E, L_\alpha \phi \rangle = 0$ for all $\phi \in \mathcal{H}$. This implies that $L_\alpha \psi_E = 0$, i.e., $\psi_E \in \ker L_\alpha$. ■

Remark 3.2 Corollary 3.3 is a generalization of [11, Corollary 4.3] where the case $K(t) = t$ is considered.

4 Generalized Weak CCR and Time-Energy Uncertainty Relations

Let A, B be symmetric operators on a Hilbert space $\mathcal{H}$ and $C \in \mathcal{B}(\mathcal{H})$ be a self-adjoint operator. We say that (A, B, C) obeys the *generalized weak CCR* (GWCCR) if

$$\langle A\psi, B\phi \rangle - \langle B\psi, A\phi \rangle = \langle \psi, iC\phi \rangle, \quad \forall \psi, \phi \in D(A) \cap D(B). \quad (4.1)$$

The case $C = I$ (the identity on $\mathcal{H}$) is the usual CCR with one degree of freedom in the sense of sesquilinear form.

For a symmetric operator A on a Hilbert space, a constant $a \in \mathbb{R}$ and a unit vector $\psi \in D(A)$, we define

$$(\Delta A)_\psi(a) := \|(A - a)\psi\|, \quad (4.2)$$

an *uncertainty of A in the state vector ψ* . The quantity $(\Delta A)_\psi(a)$ with $a = \langle \psi, A\psi \rangle$ is the usual uncertainty of A in the state vector ψ . We set

$$(\Delta A)_\psi := (\Delta A)_\psi(\langle \psi, A\psi \rangle). \quad (4.3)$$

We also introduce

$$\delta_C := \inf_{\psi \in (\ker C)^\perp, \|\psi\|=1} |\langle \psi, C\psi \rangle|. \quad (4.4)$$

Proposition 4.1 *Suppose that (A, B, C) obeys the GWCCR. Then, for all $\psi \in D(A) \cap D(B) \cap (\ker C)^\perp$ with $\|\psi\| = 1$ and all $a, b \in \mathbb{R}$,*

$$(\Delta A)_\psi(a)(\Delta B)_\psi(b) \geq \frac{\delta_C}{2}. \quad (4.5)$$

Proof: It is easy to see that $(A - a, B - b, C)$ also obeys the GWCCR. Hence, applying the Cauchy-Schwarz inequality on a non-negative, sesquilinear form on a space of linear operators [17, Lemma 1], we have for all $\psi \in D(A) \cap D(B)$

$$(\Delta A)_\psi(a)(\Delta B)_\psi(b) \geq \frac{1}{2} |\langle \psi, C\psi \rangle|.$$

Thus (4.5) follows. ■

Proposition 4.2 *Suppose that (A, B, C) obeys the GWCCR with $C \geq 0$. Then, for all $\lambda \in \mathbb{R}$, $\ker(B - \lambda) \cap D(A) \subset \ker C$ and $\ker(A - \lambda) \cap D(B) \subset \ker C$.*

Proof: Let $\psi \in \ker(B - \lambda) \cap D(A)$. Then, taking $\phi = \psi$ in (4.1), we have $\langle \psi, C\psi \rangle = 0$. Since C is nonnegative, it follows that $C\psi = 0$, i.e., $\psi \in \ker C$. ■

The following proposition gives a connection of the GWWR with the GWCCR:

Proposition 4.3 *Let (T, H, K) be a triple obeying the GWWR in $\mathcal{H}$. Assume that K is strongly differentiable on $\mathbb{R}$. Then $(T, H, K'(0))$ obeys the GWCCR:*

$$\langle T\psi, H\phi \rangle - \langle H\psi, T\phi \rangle = \langle \psi, iK'(0)\phi \rangle, \quad \psi, \phi \in D(T) \cap D(H). \quad (4.6)$$

Proof: Let $\psi, \phi \in D(T) \cap D(H)$. Then we have by (1.2)

$$\langle T\psi, e^{-itH}\phi \rangle = \langle e^{itH}\psi, T\phi \rangle + \langle e^{itH}\psi, K(t)\phi \rangle. \quad (4.7)$$

It is well known that, for all $\eta \in D(H)$, $e^{isH}\eta$ is strongly differentiable with $s \frac{d}{ds} e^{isH}\eta = iHe^{isH}\eta = ie^{isH}H\eta$. Hence the both sides of (4.7) are differentiable in t . Evaluating the derivatives at $t = 0$ and using (2.2), we obtain (4.6). ■

Propositions 4.3 and 4.1 yield the following result:

Corollary 4.4 *Suppose that the same assumption as in Proposition 4.3 holds. Then, for all $\psi \in D(T) \cap D(H) \cap (\ker K'(0))^\perp$ with $\|\psi\| = 1$ and all $t, E \in \mathbb{R}$,*

$$(\Delta T)_\psi(t)(\Delta H)_\psi(E) \geq \frac{\delta_{K'(0)}}{2}. \quad (4.8)$$

In applications to quantum theory, (4.8) gives a time-energy uncertainty relation if H is the Hamiltonian of a quantum system.

5 The Point Spectra of Generalized Time Operators

For a linear operator L on a Hilbert space $\mathcal{H}$, we introduce a subset of $\mathcal{H}$:

$$N_0(L) := \{\psi \in D(L) \mid \langle \psi, L\psi \rangle = 0\}. \quad (5.1)$$

It is obvious that $\ker L \subset N_0(L)$.

Remark 5.1 If L is a non-negative self-adjoint operator, then $N_0(L) = \ker L$.

Proposition 5.1 Assume that (T, H, K) obeys the GWR and K is strongly differentiable on $\mathbb{R}$. Then, for all $E \in \mathbb{R}$,

$$\ker(T - E) \subset N_0(K'(0)). \quad (5.2)$$

Proof: Let $\psi_0 \in \ker(T - E)$. Then $T\psi_0 = E\psi_0$. Taking the inner product of ψ_0 with the vector obtained from the operation of (2.1) to ψ_0 , we have $\langle \psi_0, e^{-itH} K(t) \psi_0 \rangle = 0$, $t \in \mathbb{R}$. Dividing the both sides by $t \neq 0$ and taking the limit $t \rightarrow 0$, we obtain $\langle \psi_0, K'(0)\psi_0 \rangle = 0$, where we have used (2.2). Hence $\psi_0 \in N_0(K'(0))$. Thus (5.2) follows. ■

Corollary 5.2 Assume that (T, H, K) obeys the GWR and K is strongly differentiable on $\mathbb{R}$. Then:

- (i) If $N_0(K'(0)) = \{0\}$, then $\sigma_p(T) = \emptyset$.
- (ii) If $K'(0) \geq 0$ or $K'(0) \leq 0$, then $\sigma_p(T|_{[D(T) \cap (\ker K'(0))^\perp]}) = \emptyset$.

Remark 5.2 Corollary (5.2) is a generalization of [11, Corollary 4.2] where the case $K(t) = t$ is considered.

Remark 5.3 It may be interesting to note that the behavior of $K(t)$ at $t = 0$ and $t = \infty$ is respectively related to $\sigma_p(T)$ (Corollary 5.2) and $\sigma_p(H)$ (Corollary 3.3).

6 Commutation Formulas and Absolute Continuity

In this section we prove commutation relations derived from the GWR. Moreover, in the special case where the commutation factor $K(t)$ is of the form tC with C a bounded self-adjoint operator, we show that H is reduced by $\overline{\text{Ran}(C)}$ ($\text{Ran}(C)$ denotes the range of C) and its reduced part is absolutely continuous.

6.1 General cases

For $p \geq 0$, we introduce a class of Borel measurable functions on $\mathbb{R}$:

$$L_p^1(\mathbb{R}) := \left\{ F : \mathbb{R} \rightarrow \mathbb{C}, \text{ Borel measurable} \mid \int_{\mathbb{R}} |F(t)|(1 + |t|^p) dt < \infty \right\}. \quad (6.1)$$

It is easy to see that $L_p^1(\mathbb{R})$ includes the space $\mathcal{S}(\mathbb{R})$ of rapidly decreasing C^∞ -functions on $\mathbb{R}$.

We say that a Borel measurable function f is in the set $\mathcal{M}_p$ if it is the Fourier transform of an element $F_f \in L_p^1(\mathbb{R})$:

$$f(\lambda) = \frac{1}{\sqrt{2\pi}} \int_{\mathbb{R}} F_f(t) e^{-it\lambda} dt, \quad \lambda \in \mathbb{R}. \quad (6.2)$$

Note that, for each $f \in \mathcal{M}_p$, F_f is uniquely determined. We have

$$\mathcal{S}(\mathbb{R}) \subset \mathcal{M}_p. \quad (6.3)$$

Moreover, every element f of $\mathcal{M}_p$ is bounded, $[p]$ times continuously differentiable ($[p]$ denotes the largest integer not exceeding p) and, for $j = 1, \dots, [p]$, $d^j f / d\lambda^j$ is bounded.

Let H be a self-adjoint operator on a Hilbert space $\mathcal{H}$ and $S : \mathbb{R} \rightarrow \mathcal{B}(\mathcal{H})$ be Borel measurable such that, for all $\psi \in \mathcal{H}$,

$$\|S(t)\psi\| \leq c(1 + |t|^p)\|\psi\|, \quad t \in \mathbb{R}$$

with constants $c > 0$ and $p \geq 0$ independent of ψ . Then, for all $\psi \in \mathcal{H}$ and $f \in \mathcal{M}_p$, the strong integral

$$f(H, S)\psi := \frac{1}{\sqrt{2\pi}} \int_{\mathbb{R}} F_f(t) e^{-itH} S(t)\psi dt \quad (6.4)$$

exists and $f(H, S) \in \mathcal{B}(\mathcal{H})$.

Theorem 6.1 *Assume that (T, H, K) obeys the GWWR. Suppose that K is strongly continuous and, for all $\psi \in \mathcal{H}$,*

$$\|K(t)\psi\| \leq c(1 + |t|^p)\|\psi\|, \quad (6.5)$$

where $c > 0$ and $p \geq 0$ are constants independent of ψ . Let $f \in \mathcal{M}_p$. Then, for all $\psi \in D(\bar{T})$, we have $f(H)\psi \in D(\bar{T})$ and

$$\bar{T}f(H)\psi = f(H)\bar{T}\psi + f(H, K)\psi, \quad (6.6)$$

where $f(H) := \int_{\mathbb{R}} f(\lambda) dE_H(\lambda)$.

Proof: Let $f \in \mathcal{M}_p$ and $\psi \in D(T^*)$, $\phi \in D(\bar{T})$. Then, by the functional calculus of the self-adjoint operator H , we have

$$\langle T^*\psi, f(H)\phi \rangle = \frac{1}{\sqrt{2\pi}} \int_{\mathbb{R}} F_f(t) \langle T^*\psi, e^{-itH}\phi \rangle dt.$$

Since $e^{-itH}\phi \in D(\bar{T}) = D(T^{**})$ by Proposition 2.2, it follows that $\langle T^*\psi, e^{-itH}\phi \rangle = \langle \psi, \bar{T}e^{-itH}\phi \rangle$. Using the fact that $(\bar{T}, H, K)$ obeys the GWWR (Proposition 2.2), we see that the r.h.s. is equal to $\langle \psi, e^{-itH}\bar{T}\phi \rangle + \langle \psi, e^{-itH}K(t)\phi \rangle$. Hence we obtain

$$\langle T^*\psi, f(H)\phi \rangle = \langle \psi, f(H)\bar{T}\phi \rangle + \frac{1}{\sqrt{2\pi}} \int_{\mathbb{R}} F_f(t) \langle \psi, e^{-itH}K(t)\phi \rangle dt.$$

Therefore $\langle T^*\psi, f(H)\phi \rangle = \langle \psi, f(H)\bar{T}\phi + f(H, K)\phi \rangle$. Since $\psi \in D(T^*)$ is arbitrary, it follows that $f(H)\phi \in D(T^{**}) = D(\bar{T})$ and (6.6) holds. $\blacksquare$

6.2 A special case

In this section we consider a special case of a triple (T, H, K) obeying the GWWR in a Hilbert space $\mathcal{H}$: We assume that K is of the form

$$K_C(t) := tC, \quad t \in \mathbb{R} \quad (6.7)$$

with C being a bounded self-adjoint operator on $\mathcal{H}$. In this case a more detailed analysis is possible as shown below.

We set

$$C_b^1(\mathbb{R}) := \{f \in C^1(\mathbb{R}) | f \text{ and } f' \text{ are bounded}\}, \quad (6.8)$$

$$C_{b,+}^1(\mathbb{R}) := \{f \in C^1(\mathbb{R}) | \text{for some } a \in \mathbb{R}, \sup_{\lambda \geq a} |f(\lambda)| < \infty \text{ and } \sup_{\lambda \geq a} |f'(\lambda)| < \infty\}. \quad (6.9)$$

Theorem 6.2 *Let C be a bounded self-adjoint operator on $\mathcal{H}$ and suppose that (T, H, K_C) obeys the GWWR.*

(i) *Let $f \in C_b^1(\mathbb{R})$. Then $f(H)D(\bar{T}) \subset D(\bar{T})$ and*

$$\bar{T}f(H)\psi - f(H)\bar{T}\psi = if'(H)C\psi \quad (6.10)$$

for all $\psi \in D(\bar{T})$.

(ii) *Suppose that H is bounded from below. Then, for all $f \in C_{b,+}^1(\mathbb{R})$, the same conclusion as that of part (i) holds. In particular, for all $z \in \mathbb{C}$ with $\Re z > 0$, $e^{-zH}D(\bar{T}) \subset D(\bar{T})$ and, for all $\psi \in D(\bar{T})$*

$$\bar{T}e^{-zH}\psi - e^{-zH}\bar{T}\psi = -ize^{-zH}C\psi. \quad (6.11)$$

Proof: (i) We first consider the case where $f \in \mathcal{M}_1$. Then it is easy to see that $f(H, K_C) = if'(H)C$. By this fact and Theorem 6.1, (6.10) holds.

We next consider the case where f is an arbitrary element in the set $C_0^1(\mathbb{R}) := \{f \in C^1(\mathbb{R}) | \text{supp } f \text{ is compact}\}$, where $\text{supp } f$ means the support of f . For a function g on $\mathbb{R}$, we set

$$\|g\|_{\infty} := \sup_{\lambda \in \mathbb{R}} |g(\lambda)|.$$

Using the Friedrichs mollifier, we can find a sequence $\{f_n\}_{n=1}^\infty \subset C_0^\infty(\mathbb{R})$ (the set of infinitely differentiable functions on $\mathbb{R}$ with compact support) such that $\sup_{n \geq 1} \|f_n\|_\infty < \infty$, $\sup_{n \geq 1} \|f'_n\|_\infty < \infty$ and, for all $\lambda \in \mathbb{R}$, $f_n(\lambda) \rightarrow f(\lambda)$, $f'_n(\lambda) \rightarrow f'(\lambda)$ as $n \rightarrow \infty$. Then, by using the functional calculus for H , we see that

$$f_n(H) \rightarrow f(H), \quad f'_n(H) \rightarrow f'(H)$$

strongly as $n \rightarrow \infty$. By the fact that $C_0^\infty(\mathbb{R}) \subset \mathcal{M}_1$ and the preceding result, we have

$$\bar{T}f_n(H)\psi = f_n(H)\bar{T}\psi + if'_n(H)C\psi$$

for all $\psi \in D(\bar{T})$. Hence $\bar{T}f_n(H)\psi \rightarrow f(H)\bar{T}\psi + if'(H)C\psi$ as $n \rightarrow \infty$. Since $\bar{T}$ is closed, it follows that $f(H)\psi \in D(\bar{T})$ and (6.10) holds.

Finally we consider the case where $f \in C_b^1(\mathbb{R})$. Let $\chi \in C_0^\infty(\mathbb{R})$ such that $\chi(0) = 1$ and set $f_n(\lambda) := \chi(\lambda/n)f(\lambda)$. Then $f_n \in C_0^1(\mathbb{R})$ and

$$f'_n(\lambda) = \frac{1}{n}\chi'\left(\frac{\lambda}{n}\right)f(\lambda) + \chi\left(\frac{\lambda}{n}\right)f'(\lambda).$$

It is easy to see that, for all $\lambda \in \mathbb{R}$, $f_n(\lambda) \rightarrow f(\lambda)$, $f'_n(\lambda) \rightarrow f'(\lambda)$ as $n \rightarrow \infty$ and

$$|f_n(\lambda)| \leq \|\chi\|_\infty \|f\|_\infty, \quad |f'_n(\lambda)| \leq \|\chi'\|_\infty \|f\|_\infty + \|\chi\|_\infty \|f'\|_\infty.$$

Hence, by the functional calculus for H , we obtain that $f_n(H) \rightarrow f(H)$, $f'_n(H) \rightarrow f'(H)$ strongly as $n \rightarrow \infty$. Then, in the same manner as in the preceding paragraph, we obtain the desired conclusion.

(ii) Let $H \geq -M$ with a constant $M \geq 0$ and $f \in C_{b,+}^1(\mathbb{R})$. Let χ be as above and define $f_n(\lambda) := \chi(\lambda/n)f(\lambda)$. Then $f_n \in C_0^1(\mathbb{R})$. Hence, by part (i),

$$\bar{T}f_n(H)\psi = f_n(H)\bar{T}\psi + if'_n(H)C\psi \tag{6.12}$$

for all $\psi \in D(\bar{T})$. We have for all $\psi \in \mathcal{H}$

$$\|f_n(H)\psi - f(H)\psi\|^2 = \int_{[-M,\infty)} F_n(\lambda) d\|E_H(\lambda)\psi\|^2. \tag{6.13}$$

with

$$F_n(\lambda) := \left| \chi\left(\frac{\lambda}{n}\right) - 1 \right|^2 |f(\lambda)|^2.$$

We have

$$|F_n(\lambda)| \leq (\|\chi\|_\infty + 1)^2 \left(\sup_{\lambda \geq -M} |f(\lambda)| \right)^2, \quad \lambda \in [-M, \infty).$$

Hence, by the Lebesgue dominated convergence theorem, the r.h.s. of (6.13) converges to 0 as $n \rightarrow \infty$. Hence $f_n(H) \rightarrow f(H)$ strongly as $n \rightarrow \infty$. Similarly we can show that $f'_n(H) \rightarrow f'(H)$. Thus, in the same way as in part (i), we obtain (6.10).

If $\Re z > 0$ ($z \in \mathbb{C}$), then the function $f_z : \mathbb{R} \rightarrow \mathbb{C}$ defined by $f_z(\lambda) := e^{-z\lambda}$ is in $C_{b,+}^1(\mathbb{R})$. Hence, by applying the preceding result, we obtain (6.11). $\blacksquare$

Remark 6.1 Theorem 6.2 is a slight generalization of [21, 3.1 Proposition 1].

Proposition 6.3 *Let C be a bounded self-adjoint operator on $\mathcal{H}$ and suppose that (T, H, K_C) obeys the GWWR. Then H and C strongly commute.*

Proof: We can apply Proposition 2.6 to obtain that $e^{-itH}Ce^{itH} = C$, $\forall t \in \mathbb{R}$. This implies the strong commutativity of H and C . ■

Lemma 6.4 *Let A and B be strongly commuting self-adjoint operators on a Hilbert space $\mathcal{H}$. Then $\overline{\text{Ran}(B)}$ reduces A .*

Proof: Since we have the orthogonal decomposition $\mathcal{H} = \ker B \oplus \overline{\text{Ran}(B)}$, it is sufficient to prove that $\ker B$ reduces A . Let P be the orthogonal projection onto $\ker B$. Then we have $P = E_B(\{0\})$. Hence, by the strong commutativity of A and B , $e^{itA}P = Pe^{itA}$ for all $t \in \mathbb{R}$. This implies that $PA \subset AP$. Thus $\ker B$ reduces A . ■

This lemma and Proposition 6.3 imply the following fact:

Corollary 6.5 *Under the same assumption as in Proposition 6.3, H is reduced by $\overline{\text{Ran}(C)}$.*

As in the case of [21, 3.2 Corollary 2], we have from Theorem 6.2 and Corollary 6.5 the following theorem. For a self-adjoint operator H , we set

$$E_H(\lambda) := E_H((-\infty, \lambda]), \quad \lambda \in \mathbb{R}.$$

Theorem 6.6 *Suppose that (T, H, K_C) obeys the GWWR. Then H is reduced by $\overline{\text{Ran}(C)}$ and the reduced part $H|_{\overline{\text{Ran}(C)}}$ is absolutely continuous. Moreover, for all $\psi, \phi \in D(\bar{T})$, the Radon-Nikodym derivative $d\langle \psi, E_H(\lambda)C\phi \rangle / d\lambda$ is given by*

$$\frac{d\langle \psi, E_H(\lambda)C\phi \rangle}{d\lambda} = i \left(\langle \bar{T}\psi, E_H(\lambda)\phi \rangle - \langle E_H(\lambda)\psi, \bar{T}\phi \rangle \right). \quad (6.14)$$

Proof: The reducibility of H by $\overline{\text{Ran}(C)}$ has already been proved in Corollary 6.5. Let $f \in \mathcal{S}(\mathbb{R})$ and $\psi, \phi \in D(\bar{T})$. It is easy to see that $\mathcal{S}(\mathbb{R}) \subset C_b^1(\mathbb{R})$. Hence, by Theorem 6.2, we have

$$\langle \bar{T}\psi, f(H)\phi \rangle - \langle \psi, f(H)\bar{T}\phi \rangle = i \langle \psi, f'(H)C\phi \rangle. \quad (6.15)$$

Let $\mu(\lambda) := \langle \psi, E_H(\lambda)C\phi \rangle$ and $\sigma(\lambda) := \langle \bar{T}\psi, E_H(\lambda)\phi \rangle - \langle E_H(\lambda)\psi, \bar{T}\phi \rangle$. Then, by (6.15) and the spectral theorem, we have

$$\int_{\mathbb{R}} f(\lambda) d\sigma(\lambda) = i \int_{\mathbb{R}} f'(\lambda) d\mu(\lambda).$$

Applying to the l.h.s. the integration by parts formula on the Stieltjes integral, we have

$$- \int_{\mathbb{R}} f'(\lambda) \sigma(\lambda) d\lambda = i \int_{\mathbb{R}} f'(\lambda) d\mu(\lambda). \quad (6.16)$$

It is easy to see that the functional $\Phi : \mathcal{S}(\mathbb{R}) \rightarrow \mathbb{C}$ defined by

$$\Phi(g) := i \int_{\mathbb{R}} g(\lambda) d\mu(\lambda) + \int_{\mathbb{R}} g(\lambda) \sigma(\lambda) d\lambda, \quad g \in \mathcal{S}(\mathbb{R})$$

is a tempered distribution on $\mathbb{R}$. Eq.(6.16) implies that $\Phi(f') = 0$ for all $f \in \mathcal{S}(\mathbb{R})$. Hence $\Phi(f) = \alpha \int_{\mathbb{R}} f(\lambda) d\lambda$ with α being a constant. This implies that the measure μ is absolutely continuous and its Radon-Nikodym derivative $d\mu/d\lambda$ is given by

$$\frac{d\mu(\lambda)}{d\lambda} = -i\alpha + i\sigma(\lambda). \quad (6.17)$$

By a limiting argument, the absolute continuity of μ can be extended to that of $\langle \psi, E_H(\cdot)C \times \phi \rangle$ for all $\psi, \phi \in \mathcal{H}$. Hence, in particular, $H|_{\text{Ran}(C)}$ is absolutely continuous. It follows from (6.17) that, for all $\lambda \in \mathbb{R}$

$$|\alpha| \leq |\mu(\lambda + 1) - \mu(\lambda)| + \sup_{\lambda' \in (\lambda, \lambda+1]} |\sigma(\lambda')|.$$

Noting the fact that $\lim_{\lambda \rightarrow -\infty} \mu(\lambda) = 0, \lim_{\lambda \rightarrow -\infty} \sigma(\lambda) = 0$, we obtain $\alpha = 0$. Hence (6.14) follows. $\blacksquare$

7 Absence of Minimum-Uncertainty States

Let (A, B, C) be a triple obeying the GWCCR. A vector $\psi_0 \in D(A) \cap D(B) \cap (\ker C)^\perp$ with $\|\psi_0\| = 1$ which attains the equality $(\Delta A)_{\psi_0}(\Delta B)_{\psi_0} = \delta_C/2 > 0$ in the uncertainty relation (4.5) with $a = \langle \psi_0, A\psi_0 \rangle$ and $b = \langle \psi_0, B\psi_0 \rangle$ is called a *minimum-uncertainty state* for (A, B, C) .

Remark 7.1 It is well-known that the Schrödinger representation (q, p) of the CCR has minimum-uncertainty states. For example, the vector $f_0 \in L^2(\mathbb{R})$ given by $f_0(x) := (2\pi)^{-1/4} \sigma^{-1/2} e^{-(x-a)^2/(4\sigma^2)}$, $x \in \mathbb{R}$ with $a \in \mathbb{R}$ and $\sigma > 0$ being constants is a minimum-uncertainty state for (q, p) : $(\Delta q)_{f_0}(\Delta p)_{f_0} = 1/2$. It follows from this fact that every representation (Q, P) of the CCR unitarily equivalent to the Schrödinger one has minimum-uncertainty states. In particular, by the von Neumann uniqueness theorem mentioned in Introduction of the present paper, every Weyl representation on a separable Hilbert space has minimum-uncertainty states. Also the Fock representation of the CCR with one degree of freedom has minimum-uncertainty states (see, e.g., [10, 11.5.1, 11.5.2]).

In this section, in contrast to the facts stated in Remark 7.1, we give a sufficient condition for a triple (T, H, C) to have *no* minimum-uncertainty states.

Theorem 7.1 (Absence of minimum-uncertainty state) *Suppose that (T, H, K_C) obeys the GWWR with T being closed. Assume that H is bounded from below and that $C \geq 0$ with $\delta_C > 0$. Then there exist no vectors $\psi_0 \in D(H) \cap D(T) \cap (\ker C)^\perp$ with $\|\psi_0\| = 1$ such that*

$$(\Delta T)_{\psi_0}(\Delta H)_{\psi_0} = \frac{\delta_C}{2} > 0. \quad (7.1)$$

Remark 7.2 An essential condition in this theorem is the boundedness below of H (note that the operators q and p in the Schrödinger representation of the CCR are unbounded both above and below with $\sigma(q) = \sigma(p) = \mathbb{R}$).

Remark 7.3 Theorem 7.1 is an extension of [11, Theorem 5.1], where the case $C = I$ is considered. A new point here is that one does not need to assume the analytic continuation property of the weak Weyl relation (the GWWR with $C = I$) as is done in [11, Theorem 5.1].

To prove Theorem 7.1, we need two lemmas.

Lemma 7.2 Assume that (A, B, C) obeys the GWCCR with $C \geq 0$. Suppose that there exists a unit vector $\psi \in D(A) \cap D(B) \cap (\ker C)^\perp$ such that

$$\begin{aligned} (A + aB + b)\psi &= 0, \\ \langle A\psi, B\psi \rangle + \langle B\psi, A\psi \rangle - 2\langle \psi, A\psi \rangle \langle \psi, B\psi \rangle &= 0, \end{aligned}$$

where a and b are complex constants. Then $\Re a = 0$ and $\Im a > 0$.

Proof : In the same way as in the proof of [11, Lemma 5.3], we can show that $i\langle \psi, C\psi \rangle = 2a(\Delta B)_\psi^2$. By Proposition 4.2, ψ cannot be an eigenvector of B . Hence $(\Delta B)_\psi \neq 0$. Thus $\Re a = 0$ and $\langle \psi, C\psi \rangle = 2(\Im a)(\Delta B)_\psi^2$. The l.h.s. on the second equation is strictly positive, since $C \geq 0$ and $\psi \in (\ker C)^\perp$. Hence $\Im a > 0$. ■

For a self-adjoint operator A on a Hilbert space, we define

$$E_0(A) := \inf \sigma(A), \tag{7.2}$$

called the *ground state energy* of A , provided that A is bounded from below. If $E_0(A)$ is an eigenvalue of A , then a non-zero vector in $\ker(A - E_0(A))$ is called a *ground state* of A .

By the variational principle, one has

$$E_0(A) = \inf_{\psi \in D(A), \|\psi\|=1} \langle \psi, A\psi \rangle. \tag{7.3}$$

The following lemma is well known (e.g., [4, Theorem 6.16]).

Lemma 7.3 Let A be a self-adjoint operator on a Hilbert space and bounded from below. Suppose that there exists a unit vector $\psi_0 \in D(A)$ such that $\langle \psi_0, A\psi_0 \rangle = E_0(A)$. Then $A\psi_0 = E_0(A)\psi_0$, i.e., ψ_0 is a ground state.

Proof of Theorem 7.1

It is sufficient to prove Theorem 7.1 in the case where $H \geq 0$. Suppose that there existed a unit vector $\psi_0 \in D(T) \cap D(H) \cap (\ker C)^\perp$ satisfying (7.1). Set $\hat{T} := T - \langle \psi_0, T\psi_0 \rangle$, $\hat{H} := H - \langle \psi_0, H\psi_0 \rangle$. Then, from the derivation of (4.5) [17, Lemma 1], we have

$$\begin{aligned} \|\hat{T}\psi_0\| \|\hat{H}\psi_0\| &= |\langle \hat{T}\psi_0, \hat{H}\psi_0 \rangle| = |\Im \langle \hat{T}\psi_0, \hat{H}\psi_0 \rangle| \\ &= \frac{1}{2} \langle \psi_0, C\psi_0 \rangle = \frac{1}{2} \delta_C. \end{aligned} \tag{7.4}$$

The first equality is the equality in the Schwarz inequality for $|\langle \hat{T}\psi_0, \hat{H}\psi_0 \rangle|$. By this fact and the condition $\delta_C > 0$, there exists a constant $c \in \mathbb{C}, c \neq 0$ such that

$$\hat{T}\psi_0 + c\hat{H}\psi_0 = 0. \quad (7.5)$$

The second equality in (7.4) implies that $\Re \langle \hat{T}\psi_0, \hat{H}\psi_0 \rangle = 0$. Hence

$$\langle T\psi_0, H\psi_0 \rangle + \langle H\psi_0, T\psi_0 \rangle - 2\langle \psi_0, T\psi_0 \rangle \langle \psi_0, H\psi_0 \rangle = 0. \quad (7.6)$$

The forth equality in (7.4) and Lemma 7.3 imply that

$$C\psi_0 = \delta_C\psi_0. \quad (7.7)$$

By (7.5), (7.6) and Lemma 7.2, we have $\Re c = 0$ and $\Im c > 0$. Hence, let $c = ia$ with $a > 0$. Then (7.5) implies that

$$T\psi_0 + iaH\psi_0 + b\psi_0 = 0, \quad (7.8)$$

where b is a constant. Hence $\langle \psi_0, T\psi_0 \rangle + ia\langle \psi_0, H\psi_0 \rangle + b = 0$, which implies that $\Im b = -a\langle \psi_0, H\psi_0 \rangle < 0$, since $H \geq 0$ and $H^{1/2}\psi_0 \neq 0$ (see (7.5)). Let $z \in \mathbb{C}$ with $\Re z > 0$. Then, by (6.11), (7.8) and (7.7), we have

$$Te^{-zH}\psi_0 = e^{-zH}(-iaH\psi_0 - b\psi_0) - iz\delta_C e^{-zH}\psi_0.$$

Since $\Re(ib/\delta_C) > 0$, we can take $z = ib/\delta_C$ so that $Te^{-zH}\psi_0 = -iae^{-zH}H\psi_0$. Hence

$$\langle e^{-zH}\psi_0, Te^{-zH}\psi_0 \rangle = -ia\langle e^{-zH}\psi_0, He^{-zH}\psi_0 \rangle.$$

The l.h.s. is real and the r.h.s. is pure imaginary. Hence the both sides must vanish. Hence $\langle e^{-zH}\psi_0, He^{-zH}\psi_0 \rangle = 0$, which implies that $H^{1/2}\psi_0 = 0$. But this is a contradiction. ■

8 Power Decays of Transition Probability Amplitudes in Quantum Dynamics

In Section 3 we have derived an estimate for transition probability amplitudes in time t . In this section we consider a triple (T, H, K_C) obeying the GWWR (discussed in Section 6.2) and show that, for state vectors in “smaller” subspaces, transition probability amplitudes decay in powers of t as $|t| \rightarrow \infty$. We apply the results to two-point correlation functions of Heisenberg operators. We also discuss decays of heat semi-groups $e^{-\beta H}$ on $\beta > 0$ the inverse of the absolute temperature.

Let H be a self-adjoint operator on a Hilbert space $\mathcal{H}$ and $C \neq 0$ be a bounded self-adjoint operator on $\mathcal{H}$. We introduce a set of generalized time operators:

$$\mathbb{T}(H, C) := \{T | (T, H, K_C) \text{ obeys the GWWR}\}. \quad (8.1)$$

By Proposition 2.5, if $T \in \mathbb{T}(H, C)$, then $T + S \in \mathbb{T}(H, C)$ for all symmetric operators S on $\mathcal{H}$ strongly commuting with H such that $D(T) \cap D(S)$ is dense in $\mathcal{H}$.

8.1 A simple case

Theorem 8.1 *Let $T \in \mathcal{T}(H, C)$ and $\psi, \phi \in D(T)$. Then, for all $t \in \mathbb{R} \setminus \{0\}$,*

$$\left| \langle \phi, e^{-itH} C \psi \rangle \right| \leq \frac{1}{|t|} (\|T\phi\| \|\psi\| + \|\phi\| \|T\psi\|). \quad (8.2)$$

Proof: In the present case, we have $L_\alpha = C$ with $\alpha = 1$. Hence Proposition 3.1 gives the desired result. $\blacksquare$

Remark 8.1 For vectors $\phi, \psi \in \mathcal{H}$, we can define a set of operators

$$\mathcal{T}_{\phi, \psi}(H, C) := \{T \in \mathcal{T}(H, C) \mid \phi, \psi \in D(T)\}$$

and put

$$c_{\phi, \psi} := \inf_{T \in \mathcal{T}_{\phi, \psi}(H, C)} (\|T\phi\| \|\psi\| + \|\phi\| \|T\psi\|),$$

then (8.2) implies that

$$\left| \langle \phi, e^{-itH} C \psi \rangle \right| \leq \frac{c_{\phi, \psi}}{|t|}. \quad (8.3)$$

Remark 8.2 Let $T \in \mathcal{T}(H, C)$. Then, for all $\psi \in D(T)$ with $\|\psi\| = 1$, $T - \langle \psi, T\psi \rangle$ is in the set $\mathcal{T}(H, C)$. Hence (8.2) implies that

$$\left| \langle \psi, e^{-itH} C \psi \rangle \right|^2 \leq \frac{4(\Delta T)_\psi^2}{t^2}. \quad (8.4)$$

Hence Theorem 8.1 gives a generalization of [11, Theorem 4.1].

8.2 Higher order decays in smaller subspaces

As demonstrated in a concrete example [11, Proposition 3.2], the modulus of a transition probability amplitude $\left| \langle \phi, e^{-itH} \psi \rangle \right|$ may decay faster than $|t|^{-1}$ as $|t| \rightarrow \infty$ for a class of vectors ϕ and ψ . In this subsection we investigate this aspect in an abstract framework and show that $\left| \langle \phi, e^{-itH} \psi \rangle \right|$ decays faster than $|t|^{-1}$ for all ϕ and ψ in smaller subspaces.

Lemma 8.2 *Let $T \in \mathcal{T}(H, C)$. Suppose that*

$$CT \subset TC. \quad (8.5)$$

Then, for all $n \in \mathbb{N}$ and $t \in \mathbb{R}$,

$$e^{-itH} D(T^n) = D(T^n) \quad (8.6)$$

$$T^n e^{-itH} = e^{-itH} (T + tC)^n. \quad (8.7)$$

Proof : By (2.8) we have the operator equality $e^{itH}T^n e^{-itH} = (T + tC)^n$. Condition (8.5) implies that $D((T + tC)^n) = D(T^n)$. Hence 8.6) follows and (8.7) holds. ■

Theorem 8.3 *Let $T \in \mathcal{T}(H, C)$. Assume (8.5). Let $n \in \mathbb{N}$ and $\psi, \phi \in D(T^n)$. We define constants $d_k^T(\phi, \psi), k = 1, \dots, n$ by the following recursion relation:*

$$d_1^T(\phi, \psi) := \|T\phi\|\|\psi\| + \|\phi\|\|T\psi\|, \quad (8.8)$$

$$d_n^T(\phi, \psi) := \|T^n\phi\|\|\psi\| + \|\phi\|\|T^n\psi\| + \sum_{r=1}^{n-1} {}_nC_r d_{n-r}^T(\phi, T^r\psi), \quad n \geq 2, \quad (8.9)$$

where ${}_nC_r := n!/[(n-r)!r!]$. Then, for all $t \in \mathbb{R} \setminus \{0\}$,

$$\left| \langle \phi, e^{-itH} C^m \psi \rangle \right| \leq \frac{d_n^T(\phi, \psi)}{|t|^n}. \quad (8.10)$$

Proof : We prove (8.10) by induction in n . The case $n = 1$ holds by Theorem 8.1. Suppose that (8.10) holds for $n = 1, \dots, m-1$ ($m \geq 2$). Let $\psi, \phi \in D(T^m)$. Condition (8.5) implies that, for all $k, r \in \mathbb{N}$, $C^k T^r \subset T^r C^k$. By this fact and Lemma 8.2, we have

$$\begin{aligned} T^m e^{-itH} \psi &= e^{-itH} (T + tC)^m \psi \\ &= e^{-itH} T^m \psi + \sum_{r=1}^{m-1} {}_mC_r t^{m-r} e^{-itH} C^{m-r} T^r \psi + t^m e^{-itH} C^m \psi. \end{aligned}$$

Hence

$$\begin{aligned} |t|^m \left| \langle \phi, e^{-itH} C^m \psi \rangle \right| &\leq \|T^m \phi\|\|\psi\| + \|\phi\|\|T^m \psi\| \\ &\quad + \sum_{r=1}^{m-1} {}_mC_r |t|^{m-r} \left| \langle \phi, e^{-itH} C^{m-r} T^r \psi \rangle \right|. \end{aligned}$$

By the induction hypothesis, we have

$$|t|^{m-r} \left| \langle \phi, e^{-itH} C^{m-r} T^r \psi \rangle \right| \leq d_{m-r}^T(\phi, T^r \psi).$$

Hence (8.10) with $n = m$ follows. ■

Theorem 8.3 can be generalized. We need a lemma.

Lemma 8.4 *Let $T_1, \dots, T_n \in \mathcal{T}(H, C)$ ($n \in \mathbb{N}$) such that $CT_j \subset T_j C$, $j = 1, \dots, n$. Then, for all $t \in \mathbb{R}$,*

$$e^{-itH} D((T_1 + tC) \cdots (T_n + tC)) \subset D(T_1 \cdots T_n) \quad (8.11)$$

and, for all $\psi \in D((T_1 + tC) \cdots (T_n + tC))$,

$$T_1 \cdots T_n e^{-itH} \psi = e^{-itH} (T_1 + tC) \cdots (T_n + tC) \psi. \quad (8.12)$$

Proof: We prove the assertion by induction in n . The case $n = 1$ obviously holds. Suppose that, for an $m \in \mathbb{N}$, (8.11) and (8.12) hold. Let $\phi \in D((T_1 + tC) \cdots (T_{m+1} + tC))$. Then the vector $(T_{m+1} + tC)\phi$ is in $D((T_1 + tC) \cdots (T_m + tC))$. Hence, by the induction hypothesis, $e^{-itH}(T_{m+1} + tC)\phi$ is in $D(T_1 \cdots T_m)$ and

$$T_1 \cdots T_m e^{-itH}(T_{m+1} + tC)\phi = e^{-itH}(T_1 + tC) \cdots (T_{m+1} + tC)\phi.$$

On the other hand, $e^{-itH}\phi$ is in $D(T_{m+1})$ and $T_{m+1}e^{-itH}\phi = e^{-itH}(T_{m+1} + tC)\phi$. Hence $T_{m+1}e^{-itH}\phi$ is in $D(T_1 \cdots T_m)$, i.e., $e^{-itH}\phi \in D(T_1 \cdots T_{m+1})$, and (8.12) with $n = m + 1$ holds. Thus the assertion holds for $n = m + 1$. $\blacksquare$

Theorem 8.5 *Let $T, T_1, \dots, T_n \in \mathfrak{T}(H, C)$ such that $CT \subset TC, CT_j \subset T_j C$, $j = 1, \dots, n$. Let $\phi \in D(T_n \cdots T_1) \cap D(T^{n-1})$ and $\psi \in \cap_{r=1}^{n-1} \cap_{1 \leq i_1 < \dots < i_r \leq n} D(T^{n-r} T_{i_1} \cdots T_{i_r})$. For $k = 1, \dots, n$, we define a constant*

$$\delta_n^T(\phi, \psi; T_1, \dots, T_n) := \|T_n \cdots T_1 \phi\| \|\psi\| + \|\phi\| \|T_1 \cdots T_n \psi\| \quad (8.13)$$

$$+ \sum_{r=1}^{n-1} \sum_{1 \leq i_1 < \dots < i_r \leq n} d_{n-r}^T(\phi, T_{i_1} \cdots T_{i_r} \psi). \quad (8.14)$$

Then, for all $t \in \mathbb{R} \setminus \{0\}$,

$$\left| \langle \phi, e^{-itH} C^n \psi \rangle \right| \leq \frac{\delta_n^T(\phi, \psi; T_1, \dots, T_n)}{|t|^n}. \quad (8.15)$$

Proof: By Lemma 8.4, we have

$$\begin{aligned} T_1 \cdots T_n e^{-itH} \psi &= e^{-itH} T_1 \cdots T_n \psi \\ &+ \sum_{r=1}^{n-1} t^{n-r} \sum_{1 \leq i_1 < \dots < i_r \leq n} e^{-itH} C^{n-r} T_{i_1} \cdots T_{i_r} \psi + t^n e^{-itH} C^n \psi. \end{aligned}$$

Hence

$$\begin{aligned} |t|^n \left| \langle \phi, e^{-itH} C^n \psi \rangle \right| &\leq \|T_n \cdots T_1 \phi\| \|\psi\| + \|\phi\| \|T_1 \cdots T_n \psi\| \\ &+ \sum_{r=1}^{n-1} |t|^{n-r} \sum_{1 \leq i_1 < \dots < i_r \leq n} \left| \langle \phi, e^{-itH} C^{n-r} T_{i_1} \cdots T_{i_r} \psi \rangle \right| \end{aligned}$$

Note that $\phi, T_{i_1} \cdots T_{i_r} \psi \in D(T^{n-r})$ for $r = 1, \dots, n-1$. Hence, by Theorem 8.3, we have

$$|t|^{n-r} \left| \langle \phi, e^{-itH} C^{n-r} T_{i_1} \cdots T_{i_r} \psi \rangle \right| \leq d_{n-r}^T(\phi, T_{i_1} \cdots T_{i_r} \psi).$$

Thus (8.15) follows. $\blacksquare$

Finally we discuss the case where condition (8.5) is not necessarily satisfied. For $n \geq 2$ and $r = 1, \dots, n-1$, we introduce a set

$$\mathbf{J}_{n,r} := \{j := (j_1, \dots, j_{r+1}) \in \{0, 1\}^{r+1} \mid j_1 + \dots + j_{r+1} = n - r\} \quad (8.16)$$

and, for each $j \in \mathbf{J}_{n,r}$, we define

$$K_{n,r}^{(j)} := T^{j_1} C T^{j_2} C \dots C T^{j_r} C T^{j_{r+1}}. \quad (8.17)$$

Let

$$\begin{aligned} \mathcal{D}_n(T, C) &:= \left\{ \psi \in D(T^n) \cap \left(\bigcap_{r=1}^{n-1} \bigcap_{j \in \mathbf{J}_{n,r}} D(K_{n,r}^{(j)}) \right) \mid K_{n,r}^{(j)} \psi \in \text{Ran}(C^r | D(T^r)), \right. \\ &\quad \left. j \in \mathbf{J}_{n,r}, r = 1, \dots, n-1 \right\} \quad (n \geq 2). \end{aligned} \quad (8.18)$$

We set $\mathcal{D}_1(T, C) := D(T)$.

Remark 8.3 If (8.5) holds, then $\mathcal{D}_n(T, C) = D(T^n)$.

Theorem 8.6 Let $T \in \mathfrak{T}(H, C)$. Then, for all $\phi \in D(T^n)$ and $\psi \in \mathcal{D}_n(T, C)$ and $t \in \mathbb{R} \setminus \{0\}$,

$$|\langle \phi, e^{-itH} C^n \psi \rangle| \leq \frac{d_n(\phi, \psi)}{|t|^n}, \quad (8.19)$$

where $d_n(\phi, \psi) > 0$ is a constant independent of t .

Proof : We prove (8.19) by induction in n . The case $n = 1$ is just Theorem 8.1. Suppose that (8.19) holds with $n = 1, \dots, m-1$ ($m \geq 2$). For all $\psi \in \mathcal{D}_m(T, C)$ we have

$$T^m e^{-itH} \psi = e^{-itH} T^m \psi + t^m e^{-itH} C^m \psi + \sum_{r=1}^{m-1} t^r \sum_{j \in \mathbf{J}_{m,r}} e^{-itH} K_{m,r}^{(j)} \psi.$$

Hence, for all $\phi \in D(T^m)$,

$$\begin{aligned} |t|^m |\langle \phi, e^{-itH} C^m \psi \rangle| &\leq \|T^m \phi\| \|\psi\| + \|\phi\| \|T^m \psi\| \\ &\quad + \sum_{r=1}^{m-1} |t|^r \sum_{j \in \mathbf{J}_{m,r}} |\langle \phi, e^{-itH} K_{m,r}^{(j)} \psi \rangle|. \end{aligned}$$

Since $K_{m,r}^{(j)} \psi \in \text{Ran}(C^r | D(T^r))$, there is a vector $\chi_j \in D(T^r)$ such that $K_{m,r}^{(j)} \psi = C^r \chi_j$. By this fact and the induction hypothesis, we have $|t|^r |\langle \phi, e^{-itH} K_{m,r}^{(j)} \psi \rangle| \leq c_{m,r}$ with a constant $c_{m,r}$ independent of t . Thus (8.19) with $n = m$ follows. $\blacksquare$

8.3 Correlation functions

In this subsection, we show that the existence of generalized time-operators gives upper bounds for correlation functions for a class of linear operators. For a linear operator A on $\mathcal{H}$ and a self-adjoint operator H on $\mathcal{H}$, we define

$$A(t) := e^{itH} A e^{-itH}, \quad t \in \mathbb{R}, \quad (8.20)$$

the Heisenberg operator of A with respect to H . Let B be a linear operator on $\mathcal{H}$. Let

$$\psi \in \cap_{t \in \mathbb{R}} [D(Ae^{-itH}) \cap D(Be^{-itH})]$$

with $\|\psi\| = 1$. Then we can define

$$W(t, s; \psi) := \langle A(t)\psi, B(s)\psi \rangle, \quad s, t \in \mathbb{R}. \quad (8.21)$$

We call it the *two-point correlation function* of A and B with respect to the vector ψ .

Theorem 8.7 *Let $T \in \mathcal{T}(H, C)$. Suppose that ψ is an eigenvector of H such that $A\psi \in D(T)$ and $B\psi \in \text{Ran}(C|D(T))$. Then, for all $t, s \in \mathbb{R}$ with $t \neq s$,*

$$|W(t, s; \psi)| \leq \frac{c_{A,B,T}}{|t - s|}, \quad (8.22)$$

where

$$c_{A,B,T} := \inf_{\chi \in D(T), B\psi = C\chi} \|TA\psi\| \|\chi\| + \|A\psi\| \|T\chi\|.$$

Proof: Let E be the eigenvalue of H with eigenvector ψ . Then we have

$$W(t, s) = e^{i(t-s)E} \langle A\psi, e^{-i(t-s)H} B\psi \rangle. \quad (8.23)$$

There exists a vector $\chi \in D(T)$ such that $B\psi = C\chi$. Hence, applying Theorem 8.1, we obtain

$$|W(t, s; \psi)| \leq \frac{\|TA\psi\| \|\chi\| + \|A\psi\| \|T\chi\|}{|t - s|}.$$

Thus (8.22) follows. ■

Theorem 8.7 can be strengthened:

Theorem 8.8 *Let $T \in \mathcal{T}(H, C)$ with (8.5). Suppose that ψ is an eigenvector of H such that $\psi \in D(A)$ and $A\psi \in D(T^n)$ and $B\psi \in \text{Ran}(C^n|D(T^n))$. Then, for all $t, s \in \mathbb{R}$ with $t \neq s$,*

$$|W(t, s)| \leq \frac{c_{A,B,T}^{(n)}}{|t - s|^n}, \quad (8.24)$$

where

$$c_{A,B,T}^{(n)} := \inf_{\chi \in D(T^n), B\psi = C^n\chi} d_n^T(A\psi, \chi).$$

Proof: This follows from (8.23) and an application of Theorem 8.3. ■

8.4 Heat semi-groups

In this subsection we assume the following:

Hypothesis (H) The self-adjoint operator H is bounded from below and there exists a closed symmetric operator T such that $T \in \mathcal{T}(H, C)$.

Then (6.11) holds. We set

$$\hat{H} := H - E_0(H). \quad (8.25)$$

We have for all $\beta \geq 0$

$$\|e^{-\beta\hat{H}}\| = 1.$$

Hence, in the same way as in Subsections 8.1 and 8.2, we obtain the following results on the decay (in β) of the quantity $\langle \phi, e^{-\beta\hat{H}} C \psi \rangle$.

Theorem 8.9 *Let $\phi, \psi \in D(T)$. Then, for all $\beta > 0$,*

$$\left| \langle \phi, e^{-\beta\hat{H}} C \psi \rangle \right| \leq \frac{1}{\beta} (\|T\phi\| \|\psi\| + \|\phi\| \|T\psi\|). \quad (8.26)$$

Theorem 8.10 *Assume (8.5). Then, for all $\phi, \psi \in D(T^n)$ and $\beta > 0$,*

$$\left| \langle \phi, e^{-\beta\hat{H}} C^n \psi \rangle \right| \leq \frac{d_n^T(\phi, \psi)}{\beta^n}. \quad (8.27)$$

Theorem 8.11 *For all $\phi \in D(T^n)$, $\psi \in \mathcal{D}_n(T, C)$ and $\beta > 0$,*

$$\left| \langle \phi, e^{-\beta\hat{H}} C^n \psi \rangle \right| \leq \frac{c_n}{\beta^n}, \quad (8.28)$$

where c_n is a constant independent of β .

9 Abstract Version of Wigner's Time-Energy Uncertainty Relation

In this section we apply Theorems 8.1 and 8.6 to establish an abstract version of Wigner's time-energy uncertainty relation [23].

Let H be a self-adjoint operator on a Hilbert space $\mathcal{H}$. In the context of quantum mechanics where H is interpreted as the Hamiltonian of a quantum system, the state vector at time $t \in \mathbb{R}$ is given by

$$\psi(t) := e^{-itH} \psi \quad (9.1)$$

with $\psi \in \mathcal{H}$ being the initial state. Let $\phi_0 \in \mathcal{H}$ and suppose that

$$\int_{\mathbb{R}} t^2 |\langle \phi_0, \psi(t) \rangle|^2 dt < \infty. \quad (9.2)$$

Then

$$N_0 := \int_{\mathbb{R}} |\langle \phi_0, \psi(t) \rangle|^2 dt \quad (9.3)$$

is finite and we can define

$$\langle t \rangle := \frac{\int_{\mathbb{R}} t |\langle \phi_0, \psi(t) \rangle|^2 dt}{N_0}, \quad (9.4)$$

$$\Delta t := \left(\frac{\int_{\mathbb{R}} (t - \langle t \rangle)^2 |\langle \phi_0, \psi(t) \rangle|^2 dt}{N_0} \right)^{1/2}. \quad (9.5)$$

Physically $\langle t \rangle$ may be interpreted as the expectation value of the “arrival time”, i.e., the time t at when the state $\psi(t)$ “arrives” at ϕ_0 . In this interpretation, Δt expresses the standard deviation of the arrival time of the initial state ψ to the state ϕ_0 .

We define

$$f_H(t; \phi_0, \psi) := \begin{cases} \langle \phi_0, H\psi(t) \rangle & ; \text{if } \psi \in D(H) \\ \langle H\phi_0, \psi(t) \rangle & ; \text{if } \phi_0 \in D(H) \end{cases}. \quad (9.6)$$

We assume that $\phi_0 \in D(H)$ or $\psi \in D(H)$ and

$$\int_{\mathbb{R}} |f_H(t; \phi_0, \psi)|^2 dt < \infty. \quad (9.7)$$

Then we can define

$$\langle \mathcal{E}_H \rangle := \frac{\int_{\mathbb{R}} \langle \psi(t), \phi_0 \rangle f_H(t; \phi_0, \psi) dt}{N_0}, \quad (9.8)$$

$$\Delta \mathcal{E}_H := \left(\frac{\int_{\mathbb{R}} |f_H(t; \phi_0, \psi)|^2 dt}{N_0} - \langle \mathcal{E}_H \rangle^2 \right)^{1/2}. \quad (9.9)$$

Theorem 9.1 *Suppose that $\phi_0 \in D(H)$ or $\psi \in D(H)$ and that (9.2) and (9.7) hold. Then*

$$\Delta t \cdot \Delta \mathcal{E}_H \geq \frac{1}{2}. \quad (9.10)$$

Proof: From the Schrödinger representation $(\hat{E}, \hat{t})$ of the CCR in $L^2(\mathbb{R})$, where $\hat{t}$ is the multiplication operator by the coordinate function $t \in \mathbb{R}$ and $\hat{E} := iD_t$ with D_t being the generalized differential operator in the variable t , we have the standard uncertainty relation

$$\|(\hat{E} - \langle u, \hat{E}u \rangle)u\| \cdot \|(\hat{t} - \langle u, \hat{t}u \rangle)u\| \geq \frac{1}{2}$$

for all $u \in D(\hat{E}) \cap D(\hat{t})$ with $\|u\| = 1$. We define $f : \mathbb{R} \rightarrow \mathbb{C}$ by $f(t) := \langle \phi_0, \psi(t) \rangle$. Then, by (9.2) and (9.7), f is in $D(\hat{t}) \cap D(\hat{E})$ with $\hat{E}f(t) = f_H(t; \phi_0, \psi)$. Let $\tilde{f}(t) := f(t)/\|f\|$. Then it is easy to see that

$$\Delta \mathcal{E}_H = \|(\hat{E} - \langle \tilde{f}, \hat{E}\tilde{f} \rangle)\tilde{f}\|, \quad \Delta t := \|(\hat{t} - \langle \tilde{f}, \hat{t}\tilde{f} \rangle)\tilde{f}\|.$$

Hence (9.10) follows. ■

Theorem 9.2 *Let $T \in \mathcal{T}(H, C)$. Suppose that $\psi = C^2\chi$ with $\chi \in D(T^2) \cap D(TC)$ satisfying $TC\chi \in \text{Ran}(C|D(T))$ and that $\phi_0 \in D(T^2) \cap D(TH)$. Then (9.10) holds.*

Proof : By the present assumption, ϕ_0 is in $D(H)$ with $H\phi_0 \in D(T)$ and $\psi = C(C\chi)$ with $C\chi \in D(T)$. Hence we can apply Theorem 8.1 to obtain

$$|\langle H\phi_0, \psi(t) \rangle| \leq \frac{1}{|t|} (\|TH\phi_0\| \|C\chi\| + \|H\phi_0\| \|TC\chi\|).$$

Using this estimate, we see that (9.7) holds. We have $\phi_0 \in D(T^2)$ and $\chi \in \mathcal{D}_2(T, C)$, where $\mathcal{D}_n(T, C)$ is defined by (8.18). Hence, by Theorem 8.6, $|\langle \phi_0, \psi(t) \rangle| \leq c/|t|^2$ with $c > 0$ a constant. This implies that (9.2) holds. Thus, by Theorem 9.1, we obtain the desired result. $\blacksquare$

10 Structure producing successively triples obeying the GWR

Let (Q, P, K_C) be a triple obeying the GWR in a Hilbert space $\mathcal{H}$, i.e.,

$$Qe^{-itP} = e^{-itP}(Q + tC), \quad \forall t \in \mathbb{R}. \quad (10.1)$$

(See Proposition 2.1.)

Lemma 10.1

(i) For all $\psi \in D(\bar{Q})$ and all $z \in \mathbb{C}$ with $\Re z > 0$, we have $e^{-zP^2}\psi \in D(\bar{Q})$ and

$$\bar{Q}e^{-zP^2}\psi = e^{-zP^2}\bar{Q}\psi - 2izPe^{-zP^2}C\psi. \quad (10.2)$$

(ii) For all $\psi \in D(\bar{Q}) \cap D(P)$ and all $t \in \mathbb{R}$, we have $e^{-itP^2}\psi \in D(\bar{Q}) \cap D(P)$ and

$$\bar{Q}e^{-itP^2}\psi = e^{-itP^2}(\bar{Q} + 2tPC)\psi. \quad (10.3)$$

Proof : (i) Let $\Re z > 0$. Then the function $t \mapsto f(t) := e^{-zt^2}$, $t \in \mathbb{R}$ is in the class $C_b^1(\mathbb{R})$ and $f'(t) = -2zte^{-zt^2}$. Hence, applying Theorem 6.2-(i), we conclude that $e^{-zP^2}D(\bar{Q}) \subset D(\bar{Q})$ and (10.2) holds for all $\psi \in D(\bar{Q})$.

(ii) Let $\varepsilon > 0$. By part (i), we have for all $\phi \in D(Q^*) = D((\bar{Q})^*)$ and $\psi \in D(\bar{Q}) \cap D(P)$

$$\langle Q^*\phi, e^{-(\varepsilon+it)P^2}\psi \rangle = \langle \phi, e^{-(\varepsilon+it)P^2}\bar{Q}\psi \rangle - \langle \phi, 2i(\varepsilon+it)Pe^{-(\varepsilon+it)P^2}C\psi \rangle.$$

Applying Proposition 6.3 with $H = P$ and $T = Q$, we see that $C\psi \in D(P)$. Hence, taking $\varepsilon \rightarrow 0$, we obtain

$$\langle Q^*\phi, e^{-itP^2}\psi \rangle = \langle \phi, e^{-itP^2}\bar{Q}\psi \rangle + \langle \phi, 2tPe^{-itP^2}C\psi \rangle.$$

This implies that $e^{-itP^2}\psi \in D(Q^{**}) = D(\bar{Q})$ and (10.3) holds. $\blacksquare$

In the rest of this section, we consider a simple case such that

$$\ker C = \{0\}. \quad (10.4)$$

Then, by Corollary 3.3 applied to $H = P$, $T = Q$ and $L_\alpha = C$, we have $\sigma_p(P) = \emptyset$. In particular, P is injective. Hence we can define

$$T(Q, P) := \frac{1}{4} [\bar{Q}P^{-1} + (\bar{Q}P^{-1})^*], \quad (10.5)$$

provided that $D(\bar{Q}P^{-1})$ is dense in $\mathcal{H}$.

Theorem 10.2 *Assume (10.4) and that $D(\bar{Q}P^{-1})$ is dense in $\mathcal{H}$. Then, for all $t \in \mathbb{R}$, $e^{-itP^2}D(T(Q, P)) \subset D(T(Q, P))$ and*

$$T(Q, P)e^{-itP^2}\psi = e^{-itP^2}(T(Q, P) + tC)\psi, \quad \psi \in D(T(Q, P)). \quad (10.6)$$

Proof: Let $\psi \in D(\bar{Q}P^{-1})$. Then $P^{-1}\psi \in D(\bar{Q}) \cap D(P)$. Hence, by Lemma 10.1(ii), $e^{-itP^2}P^{-1}\psi \in D(\bar{Q}) \cap D(P)$ for all $t \in \mathbb{R}$ and

$$\bar{Q}P^{-1}e^{-itP^2}\psi = e^{-itP^2}(\bar{Q}P^{-1} + 2tC)\psi,$$

which implies that, for all $t \in \mathbb{R}$, $e^{-itP^2}(\bar{Q}P^{-1} + 2tC) \subset \bar{Q}P^{-1}e^{-itP^2}$. Hence

$$e^{itP^2}(\bar{Q}P^{-1})^* \subset [(\bar{Q}P^{-1})^* + 2tC]e^{itP^2}$$

for all $t \in \mathbb{R}$. This implies that $e^{-itP^2}[(\bar{Q}P^{-1})^* + 2tC] \subset (\bar{Q}P^{-1})^*e^{-itP^2}$ for all $t \in \mathbb{R}$. Thus the desired result follows. $\blacksquare$

Theorem 10.2 shows that, if $D(\bar{Q}P^{-1}) \cap D((\bar{Q}P^{-1})^*)$ is dense in addition to the assumption there, then $(T(Q, P), P^2, K_C)$ obeys the GWWR. This structure is very interesting, since, apart from domain problems, it produces a series of triples obeying the GWWR. Indeed, let

$$T_1(Q, P) := T(Q, P), \quad (10.7)$$

$$T_n(Q, P) := T(T_{n-1}(Q, P), P^{2^{n-1}}), \quad n \geq 2. \quad (10.8)$$

Then $(T_n(Q, P), P^{2^n}, K_C)$ obeys the GWWR for all $n = 1, \dots, N$ ($N \in \mathbb{N}$), provided that each $n = 1, \dots, N$, $T_n(Q, P)$ is symmetric.

Example 10.1 A simple example of (Q, P) is given by the Schrödinger representation (q, p) on $L^2(\mathbb{R})$ of the CCR with one degree of freedom. It is easy to see that (q, p, K_I) obeys the GWWR. The operator $T(q, p)$ in this case is called the *Aharonov-Bohm time operator* [2, 11, 12]. Theorem 10.2 clarifies a general mathematical structure behind this operator. A simple application of Theorem 10.2 to this special case produces generalized time operators to $H = (-\Delta)^{2^{n-1}}$ ($n \in \mathbb{N}$).

We can extend the theory presented above to the case of finitely many degrees of freedom. Let Q_j ($j = 1, \dots, n$, $n \in \mathbb{N}$) be a symmetric operator on $\mathcal{H}$ and P_j be a self-adjoint operator on $\mathcal{H}$ such that P_j strongly commutes with Q_k and P_k ($j, k = 1, \dots, n$, $j \neq k$).

k). Suppose that each (Q_j, P_j, K_C) obeys the GWR with C satisfying (10.4). Then, as already shown, each P_j is injective. Suppose that $D(\bar{Q}_j P_j^{-1})$ is dense. Then we can define

$$T_j := \frac{1}{4}[\bar{Q}_j P_j^{-1} + (\bar{Q}_j P_j^{-1})^*], \quad j = 1, \dots, n. \quad (10.9)$$

By the strong commutativity of P_j 's ($j = 1, \dots, n$), the operator

$$H := \sum_{j=1}^n P_j^2 \quad (10.10)$$

is self-adjoint and nonnegative.

Theorem 10.3 *Let the assumption stated above on (Q_j, P_j) ($j = 1, \dots, n$) be satisfied. Then, for all $t \in \mathbb{R}$ and $j = 1, \dots, n$, $e^{-itH}D(T_j) \subset D(T_j)$ and*

$$T_j e^{-itH} \psi = e^{-itH} (T_j + tC) \psi, \quad \psi \in D(T_j). \quad (10.11)$$

Proof : By the strong commutativity of Q_j with P_k ($k \neq j$), we have $Q_j e^{-itH} = e^{-itH_j} Q_j e^{-itP_j^2}$ on $D(P_j)$, where $H_j := \sum_{k \neq j} P_k^2$. Hence, applying Theorem 10.2 with $Q = Q_j, P = P_j$ we obtain the desired result. ■

In this case too, a remark similar to the one after Theorem 10.2 is applicable.

11 Generalized Time Operators of Partial Differential Operators

In this section we construct classes of generalized time operators of partial differential operators.

11.1 Constructions from the Schrödinger representation of the CCR with d degrees of freedom

Let $\mathbf{q} = (q_1, \dots, q_d)$ and $\mathbf{p} = (p_1, \dots, p_d)$ be the Schrödinger representation of the CCR with d degrees of freedom. Namely, q_j is the multiplication operator by x_j , the j -th coordinate function ($x = (x_1, \dots, x_d) \in \mathbb{R}^d$) acting in $L^2(\mathbb{R}^d)$ and $p_j = -iD_j$ on $L^2(\mathbb{R}^d)$ (D_j is the generalized partial differential operator in x_j). The following properties are well known:

- (i) p_j and p_l are strongly commuting self-adjoint operators.
- (ii) For all $t \in \mathbb{R}$,

$$q_j e^{-itp_l} = e^{-itp_l} (q_j + \delta_{jl}t), \quad j, l = 1, \dots, d.$$

We denote by

$$\hat{\mathbb{R}}^d := \{k = (k_1, \dots, k_d) | k_j \in \mathbb{R}, j = 1, \dots, d\} \quad (11.1)$$

the d -dimensional momentum space and by $C^r(\hat{\mathbb{R}}^d)$ the set of r times continuously differentiable functions on $\hat{\mathbb{R}}^d$

Let $F \in C^1(\hat{\mathbb{R}}^d)$ be real-valued. Then the operators

$$H_F := F(\mathbf{p}) \quad (11.2)$$

and

$$F_j^{(1)} := (\partial_j F)(\mathbf{p}) \quad (11.3)$$

are self-adjoint. Let $N_F^{(j)}$ be a closed subset of $\hat{\mathbb{R}}^d$ with Lebesgue measure zero and $G \in C^1(\hat{\mathbb{R}}^d \setminus N_F^{(j)})$ such that

$$\sup_{k \in \hat{\mathbb{R}}^d \setminus N_F^{(j)}} |G(k) \partial_j F(k)| < \infty. \quad (11.4)$$

Then we can define a linear operator $T_{F,G}^{(j)}$ on $L^2(\mathbb{R}^d)$ by

$$T_{F,G}^{(j)} := G(\mathbf{p}) q_j + q_j G(\mathbf{p})^*, \quad (11.5)$$

with domain

$$D(T_{F,G}^{(j)}) := \mathcal{F}^{-1} C_0^1(\hat{\mathbb{R}}^d \setminus N_F^{(j)}), \quad (11.6)$$

where $\mathcal{F} : L^2(\mathbb{R}^d) \rightarrow L^2(\hat{\mathbb{R}}^d)$ is the Fourier transform:

$$(\mathcal{F}\psi)(k) := \frac{1}{(2\pi)^{d/2}} \int_{\mathbb{R}^d} e^{-ikx} \psi(x) dx, \quad \psi \in L^2(\mathbb{R}^d) \quad (11.7)$$

in the L^2 -sense. Note that, for all $u \in C_0^1(\hat{\mathbb{R}}^d \setminus N_F^{(j)})$,

$$(\mathcal{F} T_{F,G}^{(j)} \mathcal{F}^{-1} u)(k) = G(k) i \frac{\partial u(k)}{\partial k_j} + i \frac{\partial}{\partial k_j} [G(k)^* u(k)], \quad \text{a.e. } k \in \hat{\mathbb{R}}^d \setminus N_F^{(j)}. \quad (11.8)$$

It is easy to see that $T_{F,G}^{(j)}$ is symmetric [note that $D(T_{F,G}^{(j)})$ is dense in $L^2(\mathbb{R}^d)$, since $C_0^\infty(\hat{\mathbb{R}}^d \setminus N_F^{(j)})$ is dense in $L^2(\hat{\mathbb{R}}^d)$].

Lemma 11.1 *For all $t \in \mathbb{R}$ and $\psi \in D(T_{F,G}^{(j)})$, one has $e^{-itH_F} \psi \in D(T_{F,G}^{(j)})$.*

Proof: We have by the functional calculus $\mathcal{F} e^{-itH_F} \mathcal{F}^{-1} = e^{-itM_F}$, where M_F is the multiplication operator by the function F . It is easy to see that, for all $u \in C_0^1(\hat{\mathbb{R}}^d \setminus N_F^{(j)})$, $e^{-itM_F} u \in C_0^1(\hat{\mathbb{R}}^d \setminus N_F^{(j)})$. Thus the assertion follows. ■

For the functions F and G as above, the operator

$$C_{F,G}^{(j)} := [G(\mathbf{p}) + G(\mathbf{p})^*] F_j^{(1)} \quad (11.9)$$

is bounded and self-adjoint.

Proposition 11.2 *Let F and G be as above. Then $(T_{F,G}^{(j)}, H_F, K_{C_{F,G}^{(j)}})$ obeys the GWWR.*

Proof: By using the Fourier transform, it is easy to see that

$$q_j e^{-itH_F} \psi = e^{-itH_F} q_j \psi + t e^{-itH_F} F_j^{(1)} \psi$$

for all $\psi \in D(q_j) \cap D(F_j^{(1)})$, which, together with Lemma 11.1, implies that

$$\begin{aligned} G(\mathbf{p}) q_j e^{-itH_F} \psi &= e^{-itH_F} G(\mathbf{p}) q_j \psi + t e^{-itH_F} G(\mathbf{p}) F_j^{(1)} \psi, \\ q_j G(\mathbf{p})^* e^{-itH_F} \psi &= e^{-itH_F} q_j G(\mathbf{p})^* \psi + t e^{-itH_F} F_j^{(1)} G(\mathbf{p})^* \psi \end{aligned}$$

for all $\psi \in D(T_{F,G}^{(j)})$. Adding these equations, we obtain

$$T_{F,G}^{(j)} e^{-itH_F} \psi = e^{-itH_F} T_{F,G}^{(j)} \psi + t e^{-itH_F} C_{F,G}^{(j)} \psi.$$

Thus the desired result follows. ■

Example 11.1 (The free Hamiltonian of a nonrelativistic quantum particle) Consider the case where $F(k) = k^2/(2m)$ ($m > 0$ is a constant denoting the mass of a quantum particle) and $G(k) = v(k)/\partial_j F(k) = mv(k)k_j^{-1}$ with $v \in C^1(\hat{\mathbb{R}}^d)$ being bounded. Then

$$\begin{aligned} H_F &= H_{\text{NR}} := -\frac{\Delta}{2m}, \quad T_{F,G}^{(j)} = T_v^{(j)} := m \left(v(\mathbf{p}) p_j^{-1} q_j + q_j v(\mathbf{p})^* p_j^{-1} \right), \\ C_{F,G}^{(j)} &= C_v := v(\mathbf{p}) + v(\mathbf{p})^* \end{aligned}$$

with $N_F^{(j)} = \{k \in \hat{\mathbb{R}}^d | k_j = 0\}$ (hence its d -dimensional Lebesgue measure is zero), where

$$\Delta := \sum_{j=1}^d D_j^2 \tag{11.10}$$

is the d -dimensional generalized Laplacian on $L^2(\mathbb{R}^d)$. Then $(T_v^{(j)}, H_{\text{NR}}, K_{C_v})$ obeys the GWWR. The operator $T_v^{(j)}$ with $d = 1$, $m = 1/2$ and $v = 1/2$ is just the Aharonov-Bohm time operator mentioned in Example 10.1).

Example 11.2 (The free Hamiltonian of a relativistic quantum particle) Consider the case where $F(k) = \sqrt{k^2 + m^2}$ with $m > 0$ (a constant). Let $G(k) = v(k)/\partial_j F(k) = v(k)\sqrt{k^2 + m^2}k_j^{-1}$. Then

$$\begin{aligned} H_F &= H_{\text{rel}} := \sqrt{-\Delta + m^2}, \\ T_{F,G}^{(j)} &= T_{\text{rel}}^{(j)} := v(\mathbf{p}) \sqrt{-\Delta + m^2} p_j^{-1} q_j + q_j v(\mathbf{p})^* \sqrt{-\Delta + m^2} p_j^{-1}, \\ C_{F,G}^{(j)} &= C_v \end{aligned}$$

with $N_F^{(j)} = \{k \in \hat{\mathbb{R}}^d | k_j = 0\}$. Then $(T_{\text{rel}}^{(j)}, H_{\text{rel}}, K_{C_v})$ obeys the GWWR.

Example 11.3 (Powers of the d -dimensional Laplacian) Let $\alpha > 0$ be a constant and consider the case where

$$F(k) = F_\alpha(k) := |k|^{2\alpha}, \quad k \in \hat{\mathbb{R}}^d.$$

In this case we have

$$H_{F_\alpha} = (-\Delta)^\alpha.$$

Let $M_j := \mathbb{R}^d \setminus \{k \in \hat{\mathbb{R}}^d | k_j = 0\}$ and

$$T_j^{(\alpha)} := \frac{1}{2\alpha} \left\{ v(\mathbf{p})(-\Delta)^{-\alpha+1} p_j^{-1} q_j + q_j v(\mathbf{p})^* (-\Delta)^{-\alpha+1} p_j^{-1} \right\},$$

with $D(T_j^{(\alpha)}) := \mathcal{F}^{-1} C_0^1(M_j)$. Then $(T_j^{(\alpha)}, H_{F_\alpha}, K_{C_v})$ obeys the GWWR.

11.2 Abstract Dirac operators

Let $\mathcal{K}$ be a Hilbert space and A_j ($j = 1, \dots, d$) and B be bounded self-adjoint operators satisfying the anticommutation relations

$$\begin{aligned} \{A_j, A_l\} &= 2\delta_{jl}, \quad \{A_j, B\} = 0, \\ B^2 &= I, \quad j, l = 1, \dots, d, \end{aligned}$$

where $\{X, Y\} := XY + YX$. Let M be a strictly positive, continuously differentiable function on $\mathbb{R}^d$ such that $\partial_j M$ is bounded ($j = 1, \dots, d$) and the set

$$N_j := \{k \in \hat{\mathbb{R}}^d | k_j + M(k)(\partial_j M)(k) = 0\}$$

is closed with Lebesgue measure zero. Then the operator

$$p_j(M) := p_j + M(\mathbf{p})(\partial_j M)(\mathbf{p}) \tag{11.11}$$

is injective and $\mathcal{F}^{-1} C_0^\infty(\hat{\mathbb{R}}^d \setminus N_j) \subset D(p_j(M)^{-1})$.

We define an operator of Dirac type

$$H_D := \sum_{j=1}^d A_j \otimes p_j + B \otimes M(\mathbf{p}) \tag{11.12}$$

acting on $\mathcal{K} \otimes L^2(\mathbb{R}^d)$. This is an abstract Dirac operator.

Example 11.4 The usual free Dirac operator [22, p.7] is given by taking $d = 3$, $\mathcal{K} = \mathbb{C}^4$, $A_j = \alpha_j$, $M = m$ (a positive constant), $B = \beta$ with α_j and β being 4×4 -Hermitian matrices satisfying $\{\alpha_j, \alpha_l\} = 2\delta_{jl}$, $\{\beta, \alpha_j\} = 0$, $\beta^2 = I$, $j, l = 1, 2, 3$.

By a general theorem [3, Theorem 4.3], H_D is self-adjoint and

$$H_D^2 = I \otimes (-\Delta + M(\mathbf{p})^2).$$

In what follows, under the natural identification $\mathcal{K} \otimes L^2(\mathbb{R}^d) = L^2(\mathbb{R}^d; \mathcal{K})$ (the Hilbert space of $\mathcal{K}$ -valued square integrable functions on $\mathbb{R}^d$), we write $A_j \otimes p_j$ as $A_j p_j$. Let $v \in C^1(\hat{\mathbb{R}}^d)$ be bounded and

$$T_j := v(\mathbf{p}) H_D p_j(M)^{-1} q_j + q_j v(\mathbf{p})^* H_D p_j(M)^{-1} \quad (11.13)$$

with $D(T_j) := \mathcal{K} \hat{\otimes} \mathcal{F}^{-1} C_0^\infty(\hat{\mathbb{R}}^d \setminus N_j)$, where $\hat{\otimes}$ denotes algebraic tensor product.

Theorem 11.3 *The triple (T_j, H_D, K_{C_v}) obeys the GWWR.*

Proof: The self-adjoint operator $L := (-\Delta + M(\mathbf{p})^2)^{1/2}$ is strictly positive. Hence it has a bounded inverse. Let

$$U := \frac{1}{\sqrt{2}} B(H_D + BL) L^{-1/2} (L + M(\mathbf{p})^{-1/2}).$$

Then U is unitary and

$$U H_D U^* = BL.$$

(This is an abstract structure of the usual free Dirac operator, see, [22, §1.4].) We also have

$$\hat{T}_j := U T_j U^* = v(\mathbf{p}) B L p_j(M)^{-1} q_j + q_j v(\mathbf{p})^* B L p_j(M)^{-1}$$

with $D(\hat{T}_j) := U D(T_j)$ (note that $U C_v = C_v U$). Using the Fourier analysis, we can show that $(\hat{T}_j, BL, K_{C_v})$ obeys the GWWR. This implies that (T_j, H_D, K_{C_v}) obeys the GWWR. $\blacksquare$

Remark 11.1 We can apply general results established in Sections 2–10 to the Hamiltonians $H_F, H_{\text{NR}}, H_{\text{rel}}, H_{F_\alpha}$ and the Dirac operator H_D . In particular, we can derive decay estimates (in time t) for transition probability amplitudes of states with these Hamiltonians. But we do not write down them here.

12 Representations of the GWWR in Fock Spaces

In this section we show that, given a triple obeying the GWWR in a Hilbert space $\mathcal{H}$, there exist triples obeying the GWWR in Fock spaces (full Fock spaces, boson Fock spaces, fermion Fock spaces) over $\mathcal{H}$.

12.1 Tensor Representations of the GWWR

Let $n \in \mathbb{N}$ and $\mathcal{H}_j$ ($j = 1, \dots, n, n \in \mathbb{N}$) be a Hilbert space. Suppose that (T_j, H_j, K_{C_j}) obeys the GWWR in $\mathcal{H}_j$ such that T_j is closed symmetric. We consider an operator

$$L := \sum_{j=1}^n \overbrace{I \otimes \cdots \otimes I \otimes H_j \otimes I \otimes \cdots \otimes I}^n | \hat{\otimes}_{j=1}^n D(H_j), \quad (12.1)$$

where $\hat{\otimes}$ denotes algebraic tensor product. It is well known that L is self-adjoint [19, §VIII.10] and

$$e^{-itL} = e^{-itH_1} \otimes \cdots \otimes e^{-itH_n}, \quad t \in \mathbb{R}. \quad (12.2)$$

Let

$$\tilde{T}_j := \underbrace{I \otimes \cdots \otimes I \otimes T_j \otimes I \otimes \cdots \otimes I}_n, \quad (12.3)$$

$$\tilde{C}_j := \underbrace{I \otimes \cdots \otimes I \otimes C_j \otimes I \otimes \cdots \otimes I}_n. \quad (12.4)$$

Proposition 12.1 *For all $j = 1, \dots, n$, $(\tilde{T}_j, L, K_{\tilde{C}_j})$ obeys the GWWR.*

Proof : Let $\Psi = \psi_1 \otimes \cdots \otimes \psi_n$ with $\psi_j \in D(T_j), j = 1, \dots, n$. Then, for all $t \in \mathbb{R}$, $e^{-itL}\Psi = \otimes_{l=1}^n e^{-itH_l}\psi_l \in \hat{\otimes}_{l=1}^n D(T_l)$ and

$$\begin{aligned} \tilde{T}_j e^{-itL}\Psi &= e^{-itH_1}\psi_1 \otimes \cdots \otimes e^{-itH_{j-1}}\psi_{j-1} \otimes T_j e^{-itH_j}\psi_j \otimes e^{-itH_{j+1}}\psi_{j+1} \cdots \otimes e^{-itH_n}\psi_n \\ &= e^{-itH_1}\psi_1 \otimes \cdots \otimes e^{-itH_j}(T_j + tC_j)\psi_j \otimes \cdots \otimes e^{-itH_n}\psi_n \\ &= e^{-itL}(\tilde{T}_j + t\tilde{C}_j)\Psi. \end{aligned}$$

Since $\hat{\otimes}_{j=1}^n D(T_j)$ is a core of $\tilde{T}_j$, the last equality extends to all $\Psi \in D(\tilde{T}_j)$ with $e^{-itL}\Psi \in D(\tilde{T}_j)$. Hence $(\tilde{T}_j, L, K_{\tilde{C}_j})$ obeys the GWWR. $\blacksquare$

12.2 Constructions of triples obeying the GWWR in Fock Spaces

Let $\mathcal{H}$ be a Hilbert space. Then the full Fock space over $\mathcal{H}$ is defined by

$$\mathcal{F}(\mathcal{H}) := \oplus_{n=0}^{\infty} \otimes^n \mathcal{H} \quad (12.5)$$

with $\otimes^0 \mathcal{H} := \mathbb{C}$ (see, e.g., [7, §5.2] or [19, §II.4, §VIII.10] for Fock space theory). For a densely defined closed linear operator A on $\mathcal{H}$ and $n \in \{0\} \cup \mathbb{N}$, we define a linear operator $A_{\Sigma}^{(n)}$ on $\otimes^n \mathcal{H}$ as follows:

$$A_{\Sigma}^{(0)} := 0, \quad (12.6)$$

$$A_{\Sigma}^{(n)} := \overline{\left(\sum_{j=1}^n \underbrace{I \otimes \cdots \otimes I \otimes \overset{j\text{-th}}{A} \otimes I \otimes \cdots \otimes I}_n \right)} | \hat{\otimes}^n D(A) \quad (12.7)$$

for $n \geq 1$. Then the operator

$$d\Gamma(A) := \oplus_{n=0}^{\infty} A_{\Sigma}^{(n)}, \quad (12.8)$$

called the second quantization of A , is densely defined and closed. If A is self-adjoint, then so is $d\Gamma(A)$.

Let H be a self-adjoint operator on $\mathcal{H}$, T be a closed symmetric operator on $\mathcal{H}$ and $C \in \mathcal{B}(\mathcal{H})$ be self-adjoint. In what follows we take the following hypothesis:

Hypothesis (I) The triple (T, H, K_C) obeys the GWR in $\mathcal{H}$.

We define

$$T_j^{(n)} := \begin{cases} a_{n,j} & ; 0 \leq n < j \\ \underbrace{I \otimes \cdots \otimes I \otimes \overbrace{T}^{j\text{-th}} \otimes I \otimes \cdots \otimes I}_n & ; n \geq j \end{cases}, \quad (12.9)$$

where $a_{n,j}$ is an arbitrary real constant. For $j \in \mathbb{N}$, we define a linear operator T_j on $\mathcal{F}(\mathcal{H})$ by

$$T_j := \bigoplus_{n=0}^{\infty} T_j^{(n)}. \quad (12.10)$$

Then T_j is a closed symmetric operator. Let

$$C_j^{(n)} := \begin{cases} 0 & ; 0 \leq n < j \\ \underbrace{I \otimes \cdots \otimes I \otimes \overbrace{C}^{j\text{-th}} \otimes I \otimes \cdots \otimes I}_n & ; n \geq j \end{cases} \quad (12.11)$$

and

$$C_j := \bigoplus_{n=0}^{\infty} C_j^{(n)}. \quad (12.12)$$

Then C_j is a bounded self-adjoint operator on $\mathcal{F}(\mathcal{H})$.

Proposition 12.2 Under Hypothesis (I), for all $j \in \mathbb{N}$, $(T_j, d\Gamma(H), K_{C_j})$ obeys the GWR.

Proof : It follows from Proposition 12.1 that, for all n , $(T_j^{(n)}, H_{\Sigma}^{(n)}, K_{C_j^{(n)}})$ obeys the GWR. It is easy to see that this implies the desired result. ■

We next construct a triple obeying the GWR in $\mathcal{F}(\mathcal{H})$ which is reduced by the boson Fock space over $\mathcal{H}$

$$\mathcal{F}_b(\mathcal{H}) := \bigoplus_{n=0}^{\infty} \otimes_s^n \mathcal{H},$$

where $\otimes_s^n \mathcal{H}$ denotes the n -fold symmetric tensor product of $\mathcal{H}$, and the fermion Fock space over $\mathcal{H}$

$$\mathcal{F}_f(\mathcal{H}) := \bigoplus_{n=0}^{\infty} \otimes_{as}^n \mathcal{H},$$

where $\otimes_{as}^n \mathcal{H}$ denotes the n -fold antisymmetric tensor product of $\mathcal{H}$. The operator

$$\hat{N} := d\Gamma(I) \quad (12.13)$$

on $\mathcal{F}(\mathcal{H})$ is called the number operator. The vector

$$\Omega := \{1, 0, 0, \dots\} \in \mathcal{F}(\mathcal{H}) \quad (12.14)$$

is called the *Fock vacuum*. We denote by P_0 the orthogonal projection onto the one-dimensional subspace $\{z\Omega | z \in \mathbb{C}\}$. We set

$$Q_0 := I - P_0. \quad (12.15)$$

Let

$$S := Q_0 \hat{N}^{-1/2} d\Gamma(C) \hat{N}^{-1/2} Q_0 \quad (12.16)$$

Lemma 12.3 *The operator S is bounded and self-adjoint with $\|S\| \leq \|C\|$.*

Proof: It is easy to see that, for all $\Psi \in D(S)$, $(S\Psi)^{(0)} = 0$ and, for $n \geq 1$,

$$(S\Psi)^{(n)} = \frac{(d\Gamma(C)\Psi)^{(n)}}{n}.$$

Hence $\|(S\Psi)^{(n)}\| \leq \|C\| \|\Psi^{(n)}\|$, which implies that $\|S\Psi\|^2 \leq \|C\|^2 \|\Psi\|^2$. Hence S is bounded with $\|S\| \leq \|C\|$ and $D(S) = \mathcal{F}(\mathcal{H})$. It is easy to see that S is symmetric. ■

Let

$$\mathcal{D}_T := \oplus_{n=0}^{\infty} \hat{\otimes}^n D(T) \quad (12.17)$$

and

$$\tau(T) := \left(Q_0 \hat{N}^{-1/2} d\Gamma(T) \hat{N}^{-1/2} Q_0 \right) | \mathcal{D}_T. \quad (12.18)$$

Then $\tau(T)$ is symmetric.

Theorem 12.4 *Assume Hypothesis (I). Then $(\tau(T), d\Gamma(H), K_S)$ obeys the GWWR.*

Proof: For all $\Psi \in \mathcal{D}_T$, $\hat{N}^{-1/2} Q_0 \Psi \in \mathcal{D}_T$, since

$$(\hat{N}^{-1/2} Q_0 \Psi)^{(0)} = 0, \quad (\hat{N}^{-1/2} Q_0 \Psi)^{(n)} = n^{-1/2} \Psi^{(n)}, \quad n \geq 1.$$

Using this property and the easily proven fact that $e^{-it d\Gamma(H)} \hat{N}^{-1/2} Q_0 = \hat{N}^{-1/2} Q_0 e^{-it d\Gamma(H)}$, we have by direct computations

$$d\Gamma(T) \hat{N}^{-1/2} Q_0 e^{-it d\Gamma(H)} \Psi = e^{-it d\Gamma(H)} d\Gamma(T) \hat{N}^{-1/2} Q_0 \Psi + t e^{-it d\Gamma(H)} d\Gamma(C) \hat{N}^{-1/2} Q_0 \Psi.$$

Operating $Q_0 \hat{N}^{-1/2}$ to the both sides, we obtain

$$\tau(T) e^{-it d\Gamma(H)} \Psi = e^{-it d\Gamma(H)} \tau(T) \Psi + t e^{-it d\Gamma(H)} S \Psi,$$

which implies the desired result. ■

Remark 12.1 Applying the results in Section 8, we can derive decay estimates for $|\langle \Phi, e^{-it d\Gamma(H)} \Psi \rangle|$ ($t \in \mathbb{R}$) for vectors Φ, Ψ in suitable subspaces. But we do not write down them here. The same applies to other results implied by the abstract theory.

It is easy to see that $d\Gamma(H)$, $\tau(T)$ and S are reduced by $\mathcal{F}_{\#}(\mathcal{H})$ ($\# = \text{b, f}$). Hence Theorem 12.4 holds for the reduced parts of them too.

It is more interesting and important to construct generalized time operators of perturbed Hamiltonians of the form $d\Gamma(H) + V$ on the boson Fock space $\mathcal{F}_b(\mathcal{H})$ or the fermion Fock space $\mathcal{F}_f(\mathcal{H})$ with V a symmetric operator. For this purpose, the method given in Section 2.4 can be applied. But in the present paper we leave this problem for future study.

References

- [1] Y. Aharonov and D. Bohm, Significance of electromagnetic potentials in the quantum theory, *Phys. Rev.* **115** (1959), 485–491.
- [2] Y. Aharonov and D. Bohm, Time in the quantum theory and the uncertainty relation for time and energy, *Phys. Rev.* **122** (1961), 1649–1658.
- [3] A. Arai, Characterization of anticommutativity of self-adjoint operators in connection with Clifford algebra and applications, *Integr. Equat. Oper. Th.* **17** (1993), 451–463.
- [4] A. Arai, *Hilbert Spaces and Quantum Mechanics*, Kyoritsu-shuppan, Tokyo, 1997 (in Japanese).
- [5] A. Arai, Representation-theoretic aspects of two-dimensional quantum systems in singular vector potentials: canonical commutation relations, quantum algebras, and reduction to lattice quantum systems, *J. Math. Phys.* **39** (1998), 2476–2498.
- [6] A. Arai, *Mathematical Principles of Quantum Phenomena*, Asakura-shoten, Tokyo, 2005, to be published (in Japanese).
- [7] O. Bratteli and D. W. Robinson, *Operator Algebras and Quantum Statistical Mechanics 2*, Second Edition, Springer, Berlin, Heidelberg, 1997.
- [8] B. Fuglede, On the relation $PQ - QP = -iI_d$, *Math. Scand.* **20** (1967), 79–88.
- [9] S. T. Kuroda, *Spectral Theory II*, Iwanami-shoten, 1979 (in Japanese).
- [10] L. Mandel and E. Wolf, *Optical Coherence and Quantum Optics*, Cambridge University Press, Cambridge, 1995.
- [11] M. Miyamoto, A generalized Weyl relation approach to the time operator and its connection to the survival probability, *J. Math. Phys.* **42** (2001), 1038–1052.
- [12] M. Miyamoto, Characteristic decay of the autocorrelation functions prescribed by the Aharonov-Bohm time operator, arXiv:quant-ph/0105033v2, 2001.
- [13] M. Miyamoto, The various power decays of the survival probability at long times for free quantum particle, *J. Phys. A* **35** (2002), 7159–7171.
- [14] M. Miyamoto, Difference between the decay forms of the survival and nonescape probabilities, arXiv:quant-ph/0207067v2, 2002.
- [15] J. G. Muga, R. Sala Mayato and I. L. Egusquiza (Eds.), *Time in Quantum Mechanics*, Springer, 2002.
- [16] J. von Neumann, Die Eindeutigkeit der Schrödingerschen Operatoren, *Math. Ann.* **104** (1931), 570–578.
- [17] P. Pfeifer and J. Fröhlich, Generalized time-energy uncertainty relations and bounds on lifetimes of resonances, *Rev. Mod. Phys.* **67** (1995), 759–779.

- [18] C. R. Putnam, *Commutation Properties of Hilbert Space Operators*, Springer, Berlin, 1967.
- [19] M. Reed and B. Simon, *Methods of Modern Mathematical Physics I: Functional Analysis*, Academic Press, New York, 1972.
- [20] M. Reed and B. Simon, *Methods of Modern Mathematical Physics III: Scattering Theory*, Academic Press, New York, 1979.
- [21] K. Schmüdgen, On the Heisenberg commutation relation. I, *J. Funct. Anal.* **50** (1983), 8–49.
- [22] B. Thaller, *The Dirac Equation*, Springer, Berlin Heidelberg, 1992.
- [23] E. P. Wigner, On the time-energy uncertainty relation, *in Aspects of Quantum Theory* (edited by A. Salam and E. P. Wigner, Cambridge University, Cambridge, UK), 237–247.