Title	Isolation and characterization of polymorphic microsatellite DNA markers in two shrew species, <i>Sorex unguiculatus</i> and <i>S. caecutiens</i>
Author(s)	Naitoh, Y.; Ishibashi, Y.; Abe, S.; Ohdachi, S. D.
Citation	Molecular Ecology Notes, 2(4), 434-436
Issue Date	2002
Doc URL	http://hdl.handle.net/2115/1471
Rights	Copyright 2002 Blackwell Publishing
Type	article (author version)
File Information	MEN2-4.pdf

[Instructions for use](#)

Isolation and characterization of polymorphic microsatellite DNA markers in two shrew species, *Sorex unguiculatus* and *S. caecutiens*

Y. NAITOH,^{*} Y. ISHIBASHI,[†] S. ABE,^{‡,¶} and S. D. OHDACHI^{*}

5 ^{*}*Institute of Low Temperature Science, Hokkaido University, Sapporo 060-0819, Japan,*

[†]*Hokkaido Research Center, Forestry and Forest Products Research Institute, Sapporo*

062-8516, Japan, [‡]*Laboratory of Animal Cytogenetics, Center for Advanced Science*

and Technology, [¶]*Laboratory of Cytogenetics, Graduate School of Environmental Earth*

Science, Hokkaido University, Sapporo 060-0810, Japan.

10

Keywords: cross-species amplification, shrew, simple sequence repeat, *Sorex*, Soricidae, microsatellites

15

Correspondence: Satoshi D. Ohdachi

Institute of Low Temperature Science, Hokkaido University, Sapporo 060-0819, Japan.

Fax: (+81) 11 706 7142; E-mail: ohd@pop.lowtem.hokudai.ac.jp

20 Running title: Microsatellites in two *Sorex* species

Abstract

We isolated six microsatellite markers from the partial genomic libraries of two *Sorex* shrews, *S. unguiculatus* and *S. caecutiens*, and examined their allelic variation. All loci showed high allelic variation ranging from 15 to 19 alleles and all but one locus

5 conformed to Hardy-Weinberg expectations in the species where the loci were isolated.

Cross-species amplifications showed that all primers derived from *S. unguiculatus* were useful for *S. caecutiens*, while among primer sets derived from *S. caecutiens*, only one was useful for *S. unguiculatus*. Accordingly, at least five microsatellite markers were useful in *S. caecutiens* and three in *S. unguiculatus*.

To infer biogeographical history of an organism in detail, comparison of genetic structure among various local populations is essential. Recently, microsatellite markers have been routinely used to investigate genetic structure of natural populations in many species (Balloux and Lugon-Moulin 2002). With regard to soricine shrew some

5 microsatellite primers have been designed for *S. araneus* (Wyttenbach *et al.* 1997).

However, according to our preliminary study, only a few of their markers were useful for two *Sorex* species in north eastern Asia, *S. unguiculatus* and *S. caecutiens*, which are common in Hokkaido, the northernmost island of Japan (Ohdachi and Maekawa 1990).

Here, we isolated new microsatellite loci from the genomes of these two Asian *Sorex*
10 species and assessed their allelic variation in a natural population of each species.

Cross-species amplification was also performed using the primer sets designed for the species where the loci were originally isolated.

Total genomic DNAs were extracted from liver tissues preserved in 90% ethanol in one *S. unguiculatus* and one *S. caecutiens* captured in Sarobetsu moor, Hokkaido, by
15 the proteinase K/phenol/chloroform method (Sambrook *et al.* 1989). Partial genomic libraries were constructed from *Sau*3A I-digested genomic DNA, in which size-selected DNA fragments ranging from 0.5 to 1 kb were ligated into pBluescript II. For each shrew, approximately 8 000 transformed colonies were screened with a ³²P-5' end-labelled (CA)₈ probe. A second screening of the positive clones was performed,
20 from which 18 positive clones for *S. unguiculatus* and 23 positive clones for *S. caecutiens* were isolated. These isolated clones were sequenced using ABI PRISM dRhodamine Terminator Cycle Sequencing Ready Reaction Kit with an ABI PRISM 377 DNA Sequencer. Six and 11 primer sets for PCR amplification were designed for *S. unguiculatus* and *S. caecutiens*, respectively, based on the nucleotide sequences of the

clones containing a CA/GT-repeat.

Allelic variation and other characteristics at the cloned microsatellite loci were examined in both shrew species captured at Bibai in Hokkaido (*S. unguiculatus*; $N = 49$, and *S. caecutiens*; $N = 42$). Genomic DNA was extracted from liver or muscle tissue by

the proteinase K/phenol/chloroform method or the Chelex-100 method (Walsh *et al.* 1991). PCR amplification was carried out following Masuda and Yoshida (1994), or using icycler (BIO-RAD) in a 15 μ l reaction mixture containing 50 mM KCl, 1.5 mM $MgCl_2$, 10 mM Tris-HCl (pH 8.3), 0.1 % Triton X-100, 0.2 mM dNTPs, 0.5 μ M of each primer, 0.5 U rTaq DNA polymerase (TOYOBO) and 10 - 100 ng of genomic DNA.

After denaturation at 94 °C for 1 minute, cycling was performed for 30 - 35 cycles under the following conditions: 2 minutes at 94 °C, 1 minute at 55 or 58 °C and 1 minute at 72 °C. The PCR products were analysed on an ABI PRISM 310 Genetic Analyzer. Hardy-Weinberg equilibrium was tested at each of the loci using the program GENEPOP 3.1c (Raymond & Rousset 1995).

Among the synthesized primer sets, three sets for *S. unguiculatus* and eight sets for *S. caecutiens* were useless because they showed monomorphic or unclear peak patterns in the species where the primer sets were developed. As a result, three primer sets were useful for both species. These six loci showed allelic variation ranging from 15 - 19 alleles in the species where the loci were isolated. Observed heterozygosities were very high except for a locus, A10 (mean 0.787 for *S. unguiculatus* and 0.857 for *S. caecutiens*). In each species, all but one locus conformed to Hardy-Weinberg expectations (Table 1). In *S. unguiculatus*, non-amplification occurred in one of the 49 individuals at A10. Since the other two loci were amplified successfully in this individual, a null allele may exist at the locus A10 with a relatively high frequency in

the examined population. In *S. caecutiens*, the observed genotype frequencies deviated slightly from Hardy-Weinberg equilibrium at D2 ($P = 0.041$). Since both the observed and expected heterozygosities showed very high and similar values for this locus (Table 1), a sampling bias due to the small sample size in the present study might cause the deviation from Hardy-Weinberg equilibrium, although this remains to be clarified.

For the primer sets derived from *S. caecutiens*, only a locus, D2, showed high allelic variation and conformed to Hardy-Weinberg expectation in *S. unguiculatus* (Table 1). However, in *S. unguiculatus* PCR products of E1 showed monomorphic and those of D8 could not be typed because of a complex peak pattern. On the other hand, all the primer sets derived from *S. unguiculatus* provided analysable PCR products with high allelic variation ranging from 16 to 25 alleles and high observed heterozygosities (mean 0.928) in *S. caecutiens*, and all loci conformed to Hardy-Weinberg expectations (Table 1). Thus, among the isolated microsatellite markers, at least five will be useful in assessing population genetic structure in *S. caecutiens* and three in *S. unguiculatus*.

Acknowledgements

We wish to thank Dr. K. Nakata for generously providing samples and Ms. N. Etoh for technical advice.

References

- Balloux F, Lugon-Moulin N (2002) The estimation of population differentiation with microsatellite markers. *Molecular Ecology*, **11**, 155-165.
- Masuda R, Yoshida MC (1994) A molecular phylogeny of the family Mustelidae (Mammalia, Carnivora), based on comparison of mitochondrial cytochrome *b* nucleotide sequences. *Zoological Science*, **11**, 605-612.
- Ohdachi S and Maekawa K (1990) Geographic distribution and relative abundance of four species of soricine shrews in Hokkaido, Japan. *Acta Theriologica* , **35**, 261-267.
- 10 Raymond M, Rousset F (1995) GENEPOP (version 1.2): population genetics software for exact tests and ecumenicism. *Journal of Heredity* **86**, 248-249.
- Sambrook J, Fritsch EF, Maniatis T (1989) *Molecular Cloning: a Laboratory Manual*, 2nd edn. Cold spring Harbor Laboratory Press, New York.
- Walsh PS, Metzger DA, Higuchi R (1991) Chelex[®] 100 as a medium for simple extraction of DNA for PCR-based typing from forensic material. *BioTechniques*, **10**, 506-513.
- 15 Wyttenbach A, Favre L, Hausser J (1997) Isolation and characterization of simple sequence repeats in the genome of the common shrew. *Molecular Ecology*, **6**, 797-800.

Table 1 Characterization of six microsatellite loci isolated from *Sorex unguiculatus* and *S. caecutiens*. Repeat motifs are derived from the sequenced clones. Product size is the range of allele size detected in the origin species. Italicized letters are the results of cross-species amplification. Accession, EMBL/GenBank/DDBJ accession number; A, number of alleles; H_O, observed heterozygosity; H_E, expected heterozygosity; NA, not available.

5	Locus	Origin	Primer sets (5'-3')	Fluorescent	Repeat	Product	Annealing	<i>S. unguiculatus</i>			<i>S. caecutiens</i>		
	(Accession)				motif	size (bp)	temp. (°C)	(N = 49)			(N = 42)		
								A	H _O	H _E	A	H _O	H _E
5	A6	<i>S. unguiculatus</i>	GGCAGCTCTTCAGTGTTACAGG	TET	(AC) ₂₅	219-249	55	15	0.816	0.887	17	0.952	0.912
	(AB084159)		AGATTGCTGAAGGAGCCAGG	-									
10	A10	<i>S. unguiculatus</i>	CCTACATCTGGTTGAAGAGGC	6-FAM	(AC) ₂₉	181-267	55	19	0.646*	0.939	16	0.881	0.932
	(AB084160)		TGGCTTGTCATCTGGGTTCC	-									
10	B4	<i>S. unguiculatus</i>	GCCTCTGTGAACACTGAACG	TET	(AC) ₂₅	168-205	55	18	0.898	0.895	25	0.952	0.948
	(AB084161)		GACACATCACGATACAAAGGG	-									
15	D2	<i>S. caecutiens</i>	AAGTACTCCACAGCTGATGGG	HEX	(GT) ₂₃	204-236	58	18	0.878	0.912	15	0.905**	0.906
	(AB084162)		TGCATCTGGTACGTCATCGG	-									
15	D8	<i>S. caecutiens</i>	GTGTGGTTAAGGGTTGAGCC	HEX	(CA) ₁₆	174-206	55	NA	-	-	15	0.881	0.917
	(AB084163)		GCCAGAGACCAAAGTCTCGC	-									
15	E1	<i>S. caecutiens</i>	GAAAGCTTTGTGACTGTAGCTCAC	6-FAM	(CA) ₂₈	248-284	55	1	0	0	17	0.786	0.920
	(AB084164)		AGCACGGAGAAACCTATAGGAC	-									

*, ** Not conformed to Hardy-Weinberg equilibrium (* $P < 0.001$, ** $P < 0.05$).