



Title	Seroepidemiological survey of morbillivirus infection in Kuril harbor seals ( <i>Phoca vitulina stejnegeri</i> ) of Hokkaido, Japan
Author(s)	Fujii, Kei; Sato, Hiroki; Kakumoto, Chiharu; Kobayashi, Mari; Saito, Sachiko; Kariya, Tatsuya; Watanabe, Yukiko; Sakoda, Yoshihiro; Kai, Chieko; Kida, Hiroshi; Suzuki, Masatsugu
Citation	Japanese Journal of Veterinary Research, 54(2-3), 109-117
Issue Date	2006-11-30
DOI	10.14943/jjvr.54.2-3.109
Doc URL	<a href="http://hdl.handle.net/2115/16906">http://hdl.handle.net/2115/16906</a>
Type	bulletin (article)
File Information	JJVR54-2-3-2.pdf



[Instructions for use](#)

## Seroepidemiological survey of morbillivirus infection in Kuril harbor seals (*Phoca vitulina stejnegeri*) of Hokkaido, Japan

Kei Fujii<sup>1</sup>, Hiroki Sato<sup>2</sup>, Chiharu Kakumoto<sup>3</sup>, Mari Kobayashi<sup>4</sup>, Sachiko Saito<sup>5</sup>,  
Tatsuya Kariya<sup>6</sup>, Yukiko Watanabe<sup>7</sup>, Yoshihiro Sakoda<sup>8</sup>, Chieko Kai<sup>2</sup>,  
Hiroshi Kida<sup>8</sup> and Masatsugu Suzuki<sup>1</sup>\*

(Accepted for publication : November 6, 2006)

### Abstract

Serological analysis was performed to detect morbillivirus infection in Kuril harbor seals in Hokkaido, Japan. Serum samples were collected from the seals at Nosappu (231 sera), Akkeshi (16), and Erimo (75) between 1998 and 2005. Antibodies to phocine distemper virus (PDV) were detected by ELISA in seals from Nosappu and Erimo. Antibodies to PDV were found in 56% (5/9) of the sampled seals from Nosappu in 1998, versus only 5% (3/66) for 2003, 1% (1/79) for 2004, and 1% (1/77) for 2005. These suggest epidemic caused by the virus in or before 1998. As antibody-positive seals included juvenile seals in 2003 and 2005, sporadic infections of the virus are thought to have occurred in recent years. In Erimo, antibodies to PDV were found in 50% (14/28) of sampled seals in 2004, versus only 13% (1/8) for 1999, 7% (1/15) for 2003, and 0% (0/24) for 2005. These suggest sporadic infection

---

<sup>1</sup>Laboratory of Wildlife Biology, Graduate School of Veterinary Medicine, Hokkaido University, Sapporo 060-0818, Japan

<sup>2</sup>Laboratory Animal Research Center, The Institute of Medical Science, The University of Tokyo, Tokyo 108-8634, Japan

<sup>3</sup>Eco Friends, Sapporo 060-0818, Japan

<sup>4</sup>Laboratory of Aqua Resource Management, Department of Aqua Bioscience and Industry, Tokyo University of Agriculture, Abashiri 099-2422, Japan

<sup>5</sup>Laboratory of Veterinary Anatomy, Obihiro University of Agriculture and Veterinary Medicine, Obihiro 080-8555, Japan

<sup>6</sup>National Research Center for Protozoan Diseases, Obihiro University of Agriculture and Veterinary Medicine, Obihiro 080-8555, Japan

<sup>7</sup>Institute for Raptor Biomedicine Japan. Address : 2-2101 Hokuto, Kushiro 084-0922, Japan

<sup>8</sup>Laboratory of Microbiology, Department of Disease Control, Graduate School of Veterinary Medicine, Hokkaido University

\*Corresponding author : Masatsugu Suzuki, Laboratory of Wildlife Biology, Graduate School of Veterinary Medicine, Hokkaido University, Sapporo 060-0818, Japan

Tel. : +81-11-706-5210 ; Fax : +81-11-706-5569 ; E-mail : mszk@vetmed.hokudai.ac.jp

by the virus before 2003 and the epizootic between after autumn in 2003, when samples of 2003 were collected, and 2004. Since antibodies to canine distemper virus (CDV) were detected in one adult seal from Nosappu in each year from 2003 to 2005, sporadic infections of the virus were suggested. There were no difference in incidence of seals with antibodies to the viruses between males and females and between juveniles and adults.

**Key Words :** Canine distemper virus, Kuril harbor seal, Marine mammal, Morbillivirus, Phocine distemper virus

### Introduction

Canine distemper virus (CDV) and phocine distemper virus (PDV), two closely related viruses of the genus *Morbillivirus*, family *Paramyxoviridae*, are single-stranded negative RNA viruses. CDV is highly contagious pathogen that has a worldwide distribution<sup>3,12</sup> and wide host range that includes terrestrial carnivores<sup>5</sup> and phocids<sup>35</sup>.

Epizootics in Baikal seals (*Phoca sibirica*) in 1987<sup>13</sup> and Caspian seals (*P. caspica*) in 2000<sup>23</sup> were attributed to CDV. PDV was first recognized in 1988 after being isolated from an epizootic in which more than 17,000 Eastern Atlantic harbor seals (*P. vitulina vitulina*) and grey seals (*Halichoerus grypus*) died in the North Sea<sup>34</sup>. In 2002, more than 22,000 dead Eastern Atlantic harbor seals were again found in the North Sea<sup>15,37</sup>. The cause of death was a PDV infection that had infected the same populations in 1988<sup>21</sup>. Infections of CDV and PDV in seals resulted in polysynthetic disorders such as bronchopneumonia, lymphoid depletion, encephalitis, and dermatitis. Widespread screenings suggest that many populations of pinnipeds in the North Atlantic were exposed to these viruses before and after the 1988 PDV outbreak<sup>7-10,41</sup>. Serological investigation of CDV and PDV infection in seals of other areas including the Antarctic Ocean<sup>1</sup> and the Northwestern Pacific<sup>31</sup> revealed that infections occurred in the ani-

mals.

The Kuril harbor seal (*P. v. stejnegeri*) is one of five harbor seal subspecies<sup>2,25</sup>. Kuril harbor seals are distributed in the Northwestern Pacific from the coast of Hokkaido northward through the Kuril Islands and eastern Kamchatka as far north as the Commander Islands. The population of Kuril harbor seals in Hokkaido dramatically declined in the 1960s and 1970s from hunting<sup>20</sup>. The number of hauling-out sites also decreased during that period. Although the population has been rebounding for the past twenty years, exceeding 900 individuals in 2002, the number of hauling-out sites has not recovered<sup>40</sup>. Approximately 60% of the Kuril harbor seals in Hokkaido are concentrated at two hauling-out sites (Daikoku Island, Akkeshi and Erimo) of the seven<sup>4,22,40</sup>. Overpopulation at hauling-out sites may make the seals susceptible to infectious disease. Mass mortalities of marine mammals caused by viral infection have not been reported in Hokkaido, but morbillivirus infection in pinnipeds in Hokkaido, including Kuril harbor seals, was reported before 1998<sup>31</sup>. In this study, we surveyed morbillivirus infection in Kuril harbor seals on the coast of Hokkaido in recent years.

### Materials and Methods

#### Samples

Serum samples were collected from Kuril harbor seals at Nosappu (231 sera), Akkeshi




Fig. 1 Map of the sampling area  
Serum samples were collected from Kuril harbor seals at Nosappu (231 sera), Akkeshi (16), and Erimo (75) between 1998 and 2005.

(16), and Erimo (75) between 1998 and 2005 (Fig. 1). All samples from Nosappu and Akkeshi were obtained from seals by-caught in fixed salmon nets from late August to November. Of the samples from Erimo, 46 were collected from seals captured under investigative capture from late June to early July and 29 samples were collected from seals by-caught in fixed salmon nets from late August to November. Body length (nose to tip of tail) was measured as an index of age. Seals measuring 125 cm or less were judged to be juveniles (age < 2 years)<sup>11,29,42</sup>. Of the juvenile seals at Erimo, those captured from late June to early July and having glossy hair were judged to be newborns younger than 2 months, because birthing season is mid to late May<sup>29,30</sup> and hair of seals that are 1 year old before molting season (July-August) is not glossy<sup>11</sup>. The newborn seals are considered to contain maternal antibodies<sup>39,46</sup>.

#### Cells and viruses

Vero cells were cultivated in Eagle's MEM supplemented with 5% calf serum. PDV and the Onderstepoort strain of CDV

from the repository of the Laboratory Animal Research Center, The Institute of Medical Science, The University of Tokyo were propagated in Vero cells. Antigens of Vero cells infected with PDV or CDV were dissolved in lysis buffer (1% Triton X-100, 1 mM iodoacetamide, 0.2 U/ml of the trypsin inhibitor aprotinin, 1 mM phenylmethylsulfonyl fluoride, 1% sodium deoxycholate, 0.14 M NaCl, and 10 mM Tris-HCl at pH 8.0) and used as antigens in enzyme-linked immunosorbent assay (ELISA). Uninfected cell lysate was used as antigen for negative controls.

#### Enzyme-linked immunosorbent assay (ELISA)

Antibodies to PDV or CDV were detected by ELISA according to modified procedure described previously<sup>24,32</sup>. The lysate of Vero cells infected with PDV or CDV and uninfected cells was used as antigen to precoat the 96-well microtiter plates at 4 °C for at least 2 hr. After the wells were blocked with 1% bovine serum albumin (BSA) in PBS for 1 h at room temperature (RT), they were washed with PBST (PBS that contains 0.05% concentration of Tween 20). Fifty  $\mu$ l of serum diluted at 1 : 100 in PBS containing 0.5% BSA and 0.05% Tween 20 (BSA-PBST) was added to each well. After incubation at RT for 1 hr, the plates were washed with PBST. Fifty  $\mu$ l of Peroxidase-conjugated Protein G (Sigma) diluted at 1 : 200 in BSA-PBST was added to each well, and the plate was incubated for 1 hr at RT. The plates were washed with PBST, and 100  $\mu$ l of substrate solution (0.05 M citrate buffer pH 4.0, 0.008% hydrogen peroxide; 40 mM 2,2'-azino-di-3-ethyl-benzothiazobine-6-sulfuric acid) was added to each well. After RT incubation for 30 min, the optical density (OD) of each well was read by spectrophotometer using a 405 nm filter. Samples with OD values at least twice as great as the negative control were regarded as positive<sup>32</sup>.

Table 1 Number of anti-PDV antibodies positive sera collected from Kuril harbor seals in Nosappu, Akkeshi, and Erimo, Hokkaido, Japan.

	Nosappu					Akkeshi					Erimo				
	Juvenile <sup>a)</sup>		Adult		Total	Juvenile		Adult		Total	Juvenile (Juvenile excluding newborn <sup>b)</sup> )		Adult		Total
	Male	Female	Male	Female		Male	Female	Male	Female		Male	Female	Male	Female	
1998	0/0 <sup>c)</sup>	1/3	2/2	2/4	5/9	—	—	—	—	—	—	—	—	—	—
1999	—	—	—	—	—	—	—	—	—	—	0/3 (0/1)	1/2 (0/0)	0/3	—	1/8
2003	2/13	1/16	0/18	0/19	3/66	—	—	—	—	—	0/6 (0/4)	0/2 (0/1)	1/3	0/4	1/15
2004	0/12	0/6	0/35	1/26	1/79	—	—	0/2	0/2	0/4	8/13 (4/6)	4/10 (2/7)	1/3	1/2	14/28
2005	0/11	1/9	0/32	0/25	1/77	0/5	0/4	0/2	0/1	0/12	0/13 (0/3)	0/9 (0/2)	0/2	—	0/24

- a) Seals measuring 125 cm or less, were judged to be juvenile (age < 2 years). Juvenile seals of Nosappu and Akkeshi did not contain newborn seals.  
 b) Of the juvenile seals at Erimo, those captured from late June to early July and having glossy hair were judged to be newborn younger than 2 months.  
 c) Number of positive / number of sample.

Table 2 Number of anti-CDV antibodies positive sera collected from Kuril harbor seals in Nosappu, Akkeshi, and Erimo, Hokkaido, Japan.

	Nosappu					Akkeshi					Erimo				
	Juvenile <sup>a)</sup>		Adult		Total	Juvenile		Adult		Total	Juvenile (Juvenile excluding newborn <sup>b)</sup> )		Adult		Total
	Male	Female	Male	Female		Male	Female	Male	Female		Male	Female	Male	Female	
1998	0/0 <sup>c)</sup>	0/3	0/2	0/4	0/9	—	—	—	—	—	—	—	—	—	—
1999	—	—	—	—	—	—	—	—	—	—	0/3 (0/1)	0/2 (0/0)	0/3	—	0/8
2003	0/13	0/16	1/18	0/19	1/66	—	—	—	—	—	0/6 (0/4)	0/2 (0/1)	0/3	0/4	0/15
2004	0/12	0/6	0/35	1/26	1/79	—	—	0/2	0/2	0/4	0/13 (0/6)	0/10 (0/7)	0/3	0/2	0/28
2005	0/11	0/9	0/32	1/25	1/77	0/5	0/4	0/2	0/1	0/12	0/13 (0/3)	0/9 (0/2)	0/2	—	0/24

- a) Seals measuring 125 cm or less, were judged to be juvenile (age < 2 years). Juvenile seals of Nosappu and Akkeshi did not contain newborn seals.  
 b) Of the juvenile seals at Erimo, those captured from late June to early July and having glossy hair were judged to be newborn younger than 2 months.  
 c) Number of positive / number of sample.

ELISA was performed in duplicate.

## Results

Antibodies to morbillivirus were detected from serum of Kuril harbor seals from Nosappu and Erimo. Antibodies to PDV were found in 55.6% (5/9) of seals from Nosappu in 1998, versus 4.5% (3/66) for 2003, 1.3% (1/79) for 2004, and 1.3% (1/77) for 2005 (Table 1). There were statistical differences in the incidence between 1998 and other years (Fisher's exact test,  $P < 0.05$ ). Antibodies to PDV were found in 50% (14/28) of the seals from Erimo in 2004, versus 12.5% (1/8) for 1999, 6.7% (1/15) for 2003, and 0% (0/24) for 2005. There were statistical difference in the incidence between 2004 and 2003, and be-

tween 2004 and 2005 (Fisher's exact test,  $P < 0.05$ ). Incidence of seals with antibodies to PDV did not differ between males and females and between juveniles and adults at either areas in each year (Fisher's exact test,  $P > 0.05$ ). One out of 1 antibodies-positive juvenile seal in 1998 and 6 out of 12 antibodies-positive juvenile seals in 2004 from Erimo were considered to be newborn with maternal antibodies. There were no differences in incidence between juvenile males and juvenile females (excluding newborns), and there were no differences in incidence between juveniles excluding newborns and adults at Erimo in each year (Fisher's exact test,  $P > 0.05$ ). The range of OD for positive sera in PDV-ELISA was 0.22-0.56.

Antibodies to CDV were detected in the serum of 1 adult seal from Nosappu in each year from 2003 to 2005 (Table 2). There were no differences in incidence of antibodies to CDV between male and female and between juvenile and adult in each year (Fisher's exact test,  $P > 0.05$ ). The range of OD for positive sera in CDV-ELISA was 0.27-0.47. No seals had antibodies to both PDV and CDV.

### Discussion

We detected anti-PDV antibodies from the sera collected from Kuril harbor seals at Nosappu and Erimo. Ohashi and Kai<sup>31)</sup> reported antibodies to PDV in 83% (19/23) of Kuril harbor seals of Hokkaido in 1996 and 100% (2/2) in 1997. In addition, >50% incidence was reported in other pinnipeds, such as steller sea lions (*Eumetopias jubatus*) and larga seals (*P. largha*), around Hokkaido from 1994 to 1998. Those authors suggested that PDV infection in pinnipeds in Hokkaido had occurred. In our study, though the incidence of antibodies to PDV in seals at Nosappu in 1998 was as high as the incidence reported for pinnipeds of Hokkaido from 1994 to 1998, the incidences in 2003, 2004, and 2005 were low. These results suggested that the viral infection in the seals from Nosappu may have decreased during the last few years. However, because antibodies to PDV have been detected in juvenile seals in recent years, it is considered that sporadic infections have occurred in recent years. The seals of the southern Kuril Islands also may be exposed to the virus. This is suspected because most of the seals by-caught in fixed salmon nets at Nosappu are considered to be from Habomai Islands<sup>17,44)</sup>. In Erimo also, antibodies to PDV were detected from Kuril harbor seals. Sporadic infection was suggested by the samples of 1999 and 2003. The incidence of antibodies to PDV in 2004 was high, especially for juveniles. However, be-

cause 6 of 12 antibodies-positive juvenile seals in 2004 were newborn and may have had maternal antibodies, it should not be concluded that the antibody-positive rates translate directly into morbidity rates. The incidence of the antibodies in juvenile seals excluding newborns in 2004 was 46% (6/13), which was also high. The antibodies to PDV were not detected in sera in 2005. These results suggest that the epizootic of PDV occurred between after autumn 2003, when the sera of 2003 were collected, and 2004 in the population of Erimo. Antibodies to CDV were detected in 1 adult seal from Nosappu in each year from 2003 to 2004. These suggest CDV sporadic infection in the Kuril harbor seals from Nosappu. Kuril harbor seals in Hokkaido may be exposed to PDV more frequently than to CDV. For definitive diagnosis, detection of the viruses from the tissue or mucus of seals is required.

Kuril harbor seals haul out in dense concentrations<sup>30)</sup>. The hauling-out periods should be considered opportunities for transmission of the virus. Behavioral differences between male and female Kuril harbor seals have been reported, including frequency of observation at hauling-out sites<sup>30)</sup>, however, no difference in incidence of antibodies to PDV and CDV between males and females was confirmed in this study.

Whereas incidence of seals from Nosappu with antibodies to PDV tended to decrease in 2000s, the incidence of seals from Erimo tended to increase in 2004. The distance from Erimo to Akkeshi is about 170 km, there are no hauling-out sites between the two areas<sup>40)</sup>. Analysis of mtDNA has shown that there is little movement of Kuril harbor seals between Erimo and eastern Hokkaido (from Akkeshi to Nosappu)<sup>45)</sup>. The difference in the incidence trends between Nosappu and Erimo may relate to the exiguity of contact between the

Kuril harbor seals inhabiting eastern Hokkaido and the Kuril harbor seals inhabiting Erimo.

In North Sea, PDV caused mass mortality of Eastern Atlantic harbor seals<sup>15,34,37</sup>. In Hokkaido, it was suggested that PDV infection has occurred in Kuril harbor seals, however, mass mortality associated with the viral infection in this area has not been reported. The high mortality rate of harbor seals in the North Sea may be attributed in part to environmental pollutants that suppress their immunity<sup>6,14,33,38,43</sup>. The level of contamination may be one of the factors in the differing mortality of seals between the North Sea and Hokkaido.

Grey seals are probably asymptomatic carriers of PDV in North Sea<sup>14,36</sup>. These seals are thought to be carriers for PDV between Arctic seals and harbor seals in the 1988 and 2002 outbreak in the North Sea. In Hokkaido, there are five species of seal: Kuril harbor seal, large seal, ringed seal (*P. hispida*), ribbon seal (*P. fasciata*) and bearded seal (*Erignathus barbatus*). However Kuril harbor seals are found on Hokkaido coasts year round<sup>30</sup>, and the four other species visit in winter and spring, drifting south to Hokkaido with pack ice<sup>19,28</sup>. Seasonally migrating seals may be able to carry the virus to Kuril harbor seals, which are known for their sedentary behavior. Large seals are special note for this, because they often share hauling-out sites with Kuril harbor seals<sup>18,27</sup> and have great range of movement<sup>26</sup>.

Mass mortalities in animal populations lead to dramatic changes in abundance and community structure, and it has been suggested that these events play an important role in shaping long-term population dynamics and, thereby, evolutionary processes<sup>16</sup>. Continual epidemiological study, including sympatric animals, is required for manage-

ment and conservation of the marine mammals.

### Acknowledgement

We wish to thank the Erimo Seal Club, Kuril Harbor Seal Research Group, Ms. E. Nakagawa (Hokkaido University), Dr. K. Hattori (Hokkaido National Fisheries Research Institute) and the fishermen and Fishery Cooperatives of Habomai, Akkeshi and Erimo for their assistance in capturing and sampling. Seal capture was conducted under license from the Japanese Ministry of the Environment. This work was supported by the Japanese Ministry of the Environment.

### Reference

- 1) Bengston, J., Boveng, P., Franzen, U., Have, P., Heide-Jorgensen, M.P. and Härkönen, T. 1991. Antibodies to canine distemper virus in Arctic seals. *Mar. Mamm. Sci.*, 7 : 85-87.
- 2) Bigg, M.A. 1981. Harbour seal: *Phoca vitulina* Linnaeus, 1758 and *Phoca largha* Pallas, 1811. In: *Handbook of marine mammals*, Vol. 2, Seals, pp. 1-27, Ridgway, S. H. and Harrison, R.H. eds., Academic Press, New York.
- 3) Blixenkrone-Moller, M. 1993. Biological properties of phocine distemper virus and canine distemper virus. *APMIS. Supplementum*, 36 : 1-51.
- 4) Chishima, J. 1997. Census report of Kuril harbor seal along the coast of Eastern Hokkaido in 1996. *Wildlife Forum*, 3 : 113-118 (in Japanese).
- 5) Deem, S.L., Spelman, L.H., Yates, R.A. and Montali, R.J. 2000. Canine distemper in terrestrial carnivores: A review. *J. Zoo Wildl. Med.*, 31 : 441-451.
- 6) DeSwart, R.L., Ross, P.S., Vedder, L.J., Timmerman, H.H., Heisterkamp, S., Van Loveren, H., Vos, J.G., Reijnders, P.J.H.

- and Osterhaus, A.D.M.E. 1994. Impairment of immune function in harbor seals (*Phoca vitulina*) feeding on fish from polluted water. *Ambio*, 23 : 155-159.
- 7) Dietz, R., Hansen, C.T., Have, P. and Heide-Jorgen, M.P. 1989. Clue to seal epizootic?. *Nature*, 338 : 627.
  - 8) Duignan, P.J., Duffy, N., Rima, B.K. and Geraci, J.R. 1997. Comparative antibody response in harbour and grey seals naturally infected by a morbillivirus. *Vet. Immunol. Immunopathol.*, 55 : 341-349.
  - 9) Duignan, P.J., Nielsen, O., House, C., Kovacs, K.M., Duffy, N., Early, G., Sadove, S., St Aubin, D.J., Rima, B.K. and Geraci, J.R. 1997. Epizootiology of morbillivirus infection in harp, hooded, and ringed seals from the Canadian Arctic and western Atlantic. *J. Wildl. Dis.*, 33 : 7-19.
  - 10) Duignan, P.J., Saliki, J.T., St Aubin, D.J., Early, G., Sadove, S., House, J.A., Kovacs, K. and Geraci, J.R. 1995. Epizootiology of morbillivirus infection in North America harbor seals (*Phoca vitulina*) and grey seals (*Halichoerus grypus*). *J. Wildl. Dis.*, 31 : 491-501.
  - 11) Fujii, K., Ishikawa, T., Watanabe, Y., Saito, S., Nakagawa, E. and Kobayashi, Y. 2005. Tagging and measuring of Harbor seals (*Phoca vitulina stejnegeri*) in Erimo Cape, Japan. *Bulletin of the Erimo Town Museum*, 2 : 1-8 (in Japanese with English abstract).
  - 12) Gemma, T., Watari, T., Akiyama, K., Miyashima, N., Shin, Y.S., Iwatsuki, K., Kai, C. and Mikami, T. 1996. Epidemiological observations on recent outbreaks of canine distemper in Tokyo area. *J. Vet. Med. Sci.*, 58 : 547-550.
  - 13) Grachev, M.A., Kumarev, V.P., Mamaev, L. V., Zorin, V.L., Baranova, L.V., Denikina, N.N., Belikov, S.I., Petrov, E.A., Kolesnik, V.S., Kolesnik, R.S., Dorofeev, V.M., Nagieva, F.G. and Sidorov, V.N. 1989. Distemper virus in Baikal seals. *Nature*, 338 : 209-210.
  - 14) Hall, A.J., Jepson, P.D., Goodman, S.J. and Harkonen, T. 2006. Phocine distemper virus in the North and European Seals, Data and models, nature and nurture. *Biol. Conserv.*, 131 : 221-229.
  - 15) Harkonen, T., Dietz, R., Reijnders, P., Teilmann, J., Harding, K., Hall, A., Brasseur, S., Siebert, U., Goodman, S. J., Jepson, P. D., Rasmussen, T.D. and Thompson, P. 2006. A review of the 1988 and 2002 phocine distemper virus epidemics in European harbour seals. *Dis. of Aquat. Org.*, 68 : 115-130.
  - 16) Harwood, J. and Hall, A. 1990. Mass mortality in marine mammals : its implications for population dynamics and genetics. *Trends Ecol. Evol.*, 5 : 254-257.
  - 17) Hayama, S., Uno, H., and Wada, K. 1986. A migration model based on sex and age composition of Kuril seals captured along the coast of Nemuro Peninsula, Hokkaido. In : *Ecology and Protection of Kuril seal*, pp. 140-157, Wada, K. eds., Tokai University Press, Tokyo (in Japanese with English abstract).
  - 18) Ishikawa, S. 2006. Marine mammal watching at Cape Erimo. *Isana*, 44 : 11-13 (in Japanese).
  - 19) Ito, T. 1999. Pinniped, zoological character and Japanese Pinnipeds. In : *Pinniped, natural history of sea lion and seal*, pp. 3-40, Wada, K. and Ito, T. eds., Tokai University Press, Tokyo (in Japanese).
  - 20) Ito, T. and Shukunobe, T. 1986. Number and present of Kuril seal. In : *Ecology and Protection of Kuril seal*, pp. 18-58, Wada, K. eds., Tokai University Press, Tokyo. (in Japanese with English abstract).
  - 21) Jensen, T., van de Bildt, M., Dietz, H. H., Andersen, T. H., Hammer, A. S., Kuiken, T.



- and Osterhaus, A. 2002. Another phocine distemper outbreak in Europe. *Science*, 297 : 209.
- 22) Kariya, T., Kobayashi, Y., Fujii, K., Yamada, K., Nakaoka, T., Osa, Y., Chishima, J., Watanabe, Y., Saito, S., Nakagawa, E. and Wada, K. In Press. Present status and future problem on ecology and conservation of Kuril harbor seal. *Wild-life Forum* (in Japanese).
- 23) Kennedy, S., Kuiken, T., Jepson, P.D., Deaville, R., Forsyth, M., Barrett, T., van de Bildt, M.W., Osterhaus, A.D., Eybatov, I., Duck, C., Kydrmanov, A., Mitrofanov, I. and Wilson, S. 2000. Mass die-off of Caspian seals caused by canine distemper virus. *Emerg. Infect. Dis.*, 6 : 637-639.
- 24) Kida, H., Brown, L.E. and Webster, R. G. 1982. Biological activity of monoclonal antibodies to operationally defined antigenic regions on hemagglutinin molecule of A/Seal Massachusetts/ 1 /80 (H7N7) influenza virus. *Virology*, 122 : 38-47.
- 25) King, J.E. 1983. *Seals of the world*, pp. 79-85, Cornell University Press, Ithaca.
- 26) Mizuno, A.W., Onuma, M., Takahashi, M. and Ohtaishi, N. 2003. Population genetic strycture of the spotted seal *Phoca largha* along the coast of Hokkaido, based on mitochondrial DNA sequences. *Zool. Sci.*, 20 : 783-788.
- 27) Mizuno, A.W., Suzuki, M. and Ohtaishi, N. 2001. Distribution of the spotted seal *Phoca largha* along the coast of Hokkaido, Japan. *Mammal Study*, 26 : 109-118.
- 28) Mizuno, A.W., Wada, A., Ishinazaka, T., Hattori, K., Watanabe, Y. and Ohtaishi, N. 2002. Distribution and abundance of spotted seals *Phoca largha* and ribbon seals *Phoca fasciata* in the southern Sea of Okhotsk. *Ecological Research*, 17 : 79-96.
- 29) Naito, Y and Nishiwaki, M. 1972. The growth of two species of the harbor seal in the adjacent water of Hokkaido. *Sci. Rep. Whales Res. Inst.*, 24 : 127-144.
- 30) Niizuma, A. 1986. Socio-ecology and reproductive strategy of the Kuril seal. In : *Ecology and Protection of Kuril seal*, pp. 59-102, Wada, K. eds., Tokai University Press, Tokyo (in Japanese with English abstract).
- 31) Ohashi, K. and Kai, C. 2000. Morbillivirus Infections in wildlife of Japan. *Journal of the Veterinary Medicine*, 53 : 834-838 (in Japanese).
- 32) Ohashi, K., Miyazaki, N., Tanabe, S., Nakata, H., Miura, R., Fujita, K., Wakasa, C., Uema, M., Shiotani, M., Takahashi, E. and Kai, C. 2001. Seroepdemiological survey of distemper virus infection in Caspian Sea and Lake Baikal. *Veterinary Microbiology*, 82 : 203-210.
- 33) O'Shea, T. 2000. Cause of seal die-off in 1988 is still under debate. *Science*, 290 : 1097.
- 34) Osterhaus, A.D.M.E. and Vedder, E. J. 1988. Identification of virus causing recent seal deaths. *Nature*, 335 : 20.
- 35) Osterhaus, A.D.M.E., De Swart, R.L., Vos, H.W., Ross, P.S., Kenter, M.J.H. and Barrett, T. 1995. Morbilivirus infections in aquatic mammals : newly identified members of the genus. *Veterinary Microbiology*, 44 : 219-227.
- 36) Pomeroy, P.P., Hammond, J.A., Hall, A.J., Lonergan, M., Duck, C.D., Smith, V.J. and Thompson, H. 2005. Morbillivirus neutralizing antibodies in Scottish grey seals *Halichoerus grypus* : assessing the effects of the 1988 and 2002 PDV epizootics. *Mer. Ecol. Prog. Ser.*, 287 : 241-250.
- 37) Reineking, B. 2002. Seal reports 1 -43. In : *Information on dead seals in the Kattegat /Skagerrak area, the Wadden Sea and the North Sea in 2002*, Common Wadden Sea

- Secretariat : <http://waddensea-secretariat.org>
- 38) Ross, P.S., de Swart, R.L., Reijnders, P.J. H., Van Loveren, H., Vos, J.G. and Osterhaus, A.D.M.E. 1995. Contaminant-related suppression of delayed type hypersensitivity and antibody responses in harbor seals fed herring from the Baltic sea. *Environ. Health Perspect.*, 103 : 162-167.
- 39) Ross, P.S., de Swart, R.L., Visser, I.K.G., Vedder, L.J., Murk, W., Bowen, W.D. and Osterhaus, A.D.M.E. 1994. Relative immunocompetence of the newborn harbour seal, *Phoca vitulina*. *Vet. Immunol. Immunopathol.*, 42 : 331-348.
- 40) Saito, S. and Watanabe, Y. 2004. Outline and difficulty on Kuril harbor seal. In : *Management of marine mammals along the coast of Hokkaido, Japan*, pp. 23-28, Kobayashi, M., Isono T. and Hattori, K. eds., Marine Wildlife Center of Japan, Sapporo (in Japanese).
- 41) Stuenkel, S., Have, P., Osterhaus, A.D.M.E., Arnemo, J. M. and Moustgaard, A. 1994. Serological investigation of virus infections in harp seals (*Phoca groenlandica*) and hooded seals (*Cystophora cristata*). *Vet. Rec.*, 134 : 502-503.
- 42) Suzuki, M. and Yamashita, T. 1986. Sexual maturity and developmental stages of Kuril seals. In : *Ecology and Protection of Kuril seal*, pp. 179-194, Wada, K. eds., Tokai University Press, Tokyo (in Japanese with English abstract).
- 43) Vos, J.G., Birnbaum, L.S. and Osterhaus, A.D.M.E. 2000. PCBs are a health risk for human and wildlife. *Science*, 289 : 1878-1879.
- 44) Wada, K., Hayama, S., Nakaoka, T. and Uno, Y. 1991. Interaction between Kuril seals and Salmon trap net fisheries in the coastal waters of Southeastern Hokkaido. *Mar. Mamm. Sci.*, 7 : 75-84.
- 45) Watanabe, Y. 2000. Population Genetic Studies of Kuril Harbor Seal (*Phoca vitulina stejnegeri*) in Hokkaido, Japan, by mtDNA Sequences. The graduation thesis, Obihiro University of Agriculture and Veterinary Medicine, Japan.
- 46) Winters, W.D. 1981. Time dependent decreases of maternal canine virus antibodies in newborn pups. *Vet. Rec.*, 108 : 295-299.