



Title	STUDIES OF ESCHERICHIA COLI ISOLATED FROM CHICKENS, SEROLOGICAL TYPES AND SENSITIVITY TO ANTIBIOTICS
Author(s)	MIKAMI, Takeshi
Citation	Japanese Journal of Veterinary Research, 12(4), 92-92
Issue Date	1964-12
Doc URL	http://hdl.handle.net/2115/1801
Type	bulletin (article)
File Information	KJ00002369115.pdf



[Instructions for use](#)

It is concluded that the phage of *S. pullorum*, as well as those of several other bacteria, may be sufficiently available to the studies on typing of *S. pullorum* and epizootiological investigation of pullorum disease.

STUDIES OF *ESCHERICHIA COLI* ISOLATED FROM CHICKENS, SEROLOGICAL TYPES AND SENSITIVITY TO ANTIBIOTICS

Takeshi MIKAMI

*Department of Epizootiology,
Faculty of Veterinary Medicine,
Hokkaido University, Sapporo, Japan*

(Summary of Master's thesis written under
direction of Dr. S. MIURA)

Two hundred and seventy strains of *Escherichia coli* were isolated from the internal organs (heart, liver, spleen, kidneys, and lungs) of 1198 chickens; 217 from the unabsorbed yolk of the same chickens; and 185 from the droppings of 153 normal day old chickens. Although a few internal organs and unabsorbed yolks were taken from birds about 8 weeks of age; the majority were taken from normal day old chickens. The results of the present experiments may be summarized as follows:

1. Using 42 types of anti *E. coli* O serums (39 known and 3 unknown), 71.5% strains isolated from the internal organs, 79.3% from the yolk, and 74.6% from the feces were typable.
2. There were 25 different O groups of *E. coli* isolated from the internal organs, 22 from unabsorbed yolks, and 17 from droppings.
3. Of 672 strains studied, about 60% belonged to one of the following O groups; O 60 (20.7%), 8-18 (10.3%), Y 813 (9.2%), O 53 (6.7%), O 1 (5.1%), O 2 (2.7%), O 21 (2.4%), and N 128 (2.1%). Serotype O 78, which is common in poultry respiratory diseases in England, was not found in the present studies.
4. Five strains of O 28 ac were isolated from the unabsorbed yolks and 2 strains from the internal organs. The yolks yielded 2 strains of O 125 and the internal organs 2 of the same type.
5. From 24 chickens yielding *E. coli* from all tissues examined (heart, liver, spleen, kidney, lung and unabsorbed yolk), 22 yielded strains which were typed within one or two O groups.
6. Of the 672 strains tested for sensitivity to antibiotics in vitro, all were sensitive to chloramphenicol, 92% to streptomycin, and 67% to tetracycline. In general, the *E. coli* recovered from the internal organs were more resistant to streptomycin and tetracycline than those isolated from the droppings.