



Title	STUDIES ON PURIFICATION OF CLOSTRIDIUM BOTULINUM TYPE C TOXIN
Author(s)	SYUTO, Bunei
Citation	Japanese Journal of Veterinary Research, 13(2), 63-63
Issue Date	1965-06
Doc URL	<a href="http://hdl.handle.net/2115/1820">http://hdl.handle.net/2115/1820</a>
Type	bulletin (article)
File Information	KJ00002369146.pdf



[Instructions for use](#)

**STUDIES ON PURIFICATION OF  
*CLOSTRIDIUM BOTULINUM* TYPE C TOXIN**

Bunei SYUTO

*Department of Biochemistry  
Faculty of Veterinary Medicine  
Hokkaido University, Sapporo, Japan*

(Summary of Master's thesis written under direction of Dr. T. HAGA)

Procedures for development of a medium for the production of *Clostridium botulinum* type C toxin and purification of the toxin produced in this medium were described.

It was observed that the medium consisting of 4% peptone, 2% Bacto-casitone, 2% corn steep liquor, 1% yeast extract, 1% glucose, 0.5% sodium chloride and 0.05% sodium thioglycollate was most favorable for the toxin production among the media examined.

The culture filtrate in which the organisms were cultivated for 168 hours at 33°C contained  $1 \times 10^6$  mouse intraperitoneal MLD per ml.

The toxin was purified by adsorption and elution from Zn-gel, fractionation with ammonium sulfate, and gel filtration with Sephadex G-50, followed by chromatography on DEAE cellulose with stepwise and gradient elution by elevating the sodium chloride concentration in phosphate buffer solution.

The purified toxin was homogeneous electrophoretically with the specific activity of  $42.15 \times 10^6$  mouse intraperitoneal MLD per E (extinction at 275 m $\mu$  of ultraviolet ray).