



Title	INCIDENCE OF PATHOGENIC ESCHERICHIA COLI FROM NORMAL FOOD MATERIALS AND FECAL SPECIMENS
Author(s)	YABIKI, Terutake
Citation	Japanese Journal of Veterinary Research, 14(3-4), 135-135
Issue Date	1966-12
Doc URL	http://hdl.handle.net/2115/1857
Type	bulletin (article)
File Information	KJ00003418311.pdf



[Instructions for use](#)

INCIDENCE OF PATHOGENIC *ESCHERICHIA COLI* FROM NORMAL FOOD MATERIALS AND FECAL SPECIMENS

Terutake YABIKI

*Department of Veterinary Public Health
Faculty of Veterinary Medicine
Hokkaido University, Sapporo, Japan*

(Summary of Masters thesis written under direction of Dr. S. HAMADA)

The incidence of pathogenic *Escherichia coli* in 1,128 samples of raw milk, 41 samples of oysters from oyster-beds, 10 samples of squid, and in 326 feces of 100 cows, 111 pigs, 57 fowls, and 58 human beings, was investigated.

1) Twelve hundred and thirty strains of *E. coli* were isolated from the above-mentioned materials. All of these strains, and 195 stock culture strains of *E. coli* originating from oysters and market milk, were examined for pathogenic *E. coli*.

2) Thirty-one of 1,234 strains (smooth types) of *E. coli* were serologically identified as pathogenic *E. coli*. Twelve strains originated from raw milk and 19 strains originated from feces.

3) The serotypes of 12 strains from raw milk are as follows: O 26 (1 strain), O 86 and O 86a (2 strains respectively), O 112^{ac} (3 strains), O 124 (1 strain), O 125 (2 strains), and O 127a (1 strain).

The serotypes of 19 strains from feces are as follows: O 26 (3 strains from cows), O 112^{ac} (1 from fowl, 2 from pigs and 3 from human beings), O 119 (2 from human beings), O 124 (2 from pigs), O 125 (1 from cows), and O 136 (5 from cows).

4) The report that pathogenic *E. coli* was isolated from raw milk is the first one in Japan, as far as I know.