



Title	Hepatosplenic alpha/beta T cell lymphoma presenting with cold agglutinin disease.
Author(s)	Minauchi, Koichiro; Nishio, Mitsufumi; Itoh, Tomoo; Yamamoto, Satoshi; Fujimoto, Katsuya; Sato, Norihiro; Koike, Takao
Citation	Annals of Hematology, 86(2), 155-157 https://doi.org/10.1007/s00277-006-0212-3
Issue Date	2007-02
Doc URL	http://hdl.handle.net/2115/18872
Rights	The original publication is available at www.springerlink.com
Type	article (author version)
File Information	AH86-2.pdf



[Instructions for use](#)

Letter to the Editor

Hepatosplenic alpha/beta T-cell lymphoma presenting with cold agglutinin disease

Koichiro Minauchi¹, Mitsufumi Nishio^{1*}, Tomoo Itoh², Satoshi Yamamoto¹, Katsuya Fujimoto¹, Norihiro Sato³ and Takao Koike¹

1. Department of Medicine II, Hokkaido University Graduated School of Medicine, Sapporo, Japan
2. Department of Surgical Pathology, Hokkaido University Hospital, Sapporo, Japan
3. Blood Transfusion Service, Hokkaido University Hospital, Sapporo, Japan

Correspondence to: Dr. Mitsufumi Nishio, Department of Medicine II, Hokkaido Graduated School of Medicine, N15 W7, Kita-ku, Sapporo 060-8638, Japan

Tel: +81-11-716-1161 (ext 5915) Fax: +81-11-706-7710

E-mail: m-nishio@med.hokudai.ac.jp

1 Table

Dear editor, We read an article in your journal reported by Sallah et al [1] that showed autoimmune hemolytic anemia associated with hepatosplenic $\gamma\delta$ T-cell lymphoma with a great interest. Here, we report a case of a patient who had cold agglutinin disease (CAD) as the initial manifestation of hepatosplenic $\alpha\beta$ T-cell lymphoma and had a regression of hemolytic anemia following the chemotherapy and allogeneic stem cell transplantation.

In January 2005, a 58-year-old Japanese woman was admitted to Hokkaido University Hospital with increasing general fatigue. She had a 6-month history of chronic hemolytic anemia and had been seen by a local physician without any medication. Upon examination, she was very pale and jaundiced, with a tachycardia and pitting ankle edema. Splenomegaly and hepatomegaly were evident. A laboratory work-up showed a hemoglobin level of 6.3 g/dL, a hematocrit of 16.7%, and a reticulocyte count of $50.9 \times 10^9/L$. The white blood cell count was $2.5 \times 10^9/L$ with 9% of atypical lymphocytes with both round and irregular nuclei, clumped chromatin, and a rim of pale blue cytoplasm. The platelet count was $116 \times 10^9/L$. Hemolysis was suspected with an elevated indirect bilirubin concentration (2.2 mg/dL; normal 0-0.3 mg/dL), an elevated lactate dehydrogenase level (318 IU/L; normal 119-229 IU/L) and a decreased level of haptoglobin (2 mg/dL; normal 90-170 mg/dL). The direct antiglobulin test was positive for complement, but negative for IgG. Polyclonal cold agglutinin showing anti-I specificity was strongly positive with a titer of 1:2048 (normal < 1:32). Donath-Landsteiner autoantibody was not detected. Serum protein electrophoresis showed no abnormal monoclonal component. Serologic tests for human T-lymphotropic virus type-I and human immunodeficiency virus were negative. The serum Interleukin-4 (IL-4) level has increased to 14.2 pg/ml (normal <6.0 pg/ml).

Computed tomography confirmed an enlargement of liver and spleen without lymphadenopathy.

A biopsy specimen of iliac bone marrow disclosed severe hypercellular marrow containing an infiltration by atypical small sized lymphoid cells with hyperchromatic nuclei identical to those seen in the peripheral blood. Immunohistochemical staining on the section of bone marrow showed positive findings for CD3, CD5, CD8 and negative findings for CD10, CD23, terminal deoxynucleotidyl transferase (TdT), T-cell intracellular antigen-1 (TIA-1), CD56 and Epstein-Barr viral-encoded RNA-1 (EBER) in atypical lymphocytes. We performed a splenectomy and liver biopsy to confirm the diagnosis of hepatosplenic T-cell lymphoma (HSTCL). Liver histology revealed the sinusoidal infiltration of small atypical lymphocytes. A complete sparing of the liver parenchyma was observed. The spleen was massively enlarged and free from nodules on the cut surfaces. The general architecture was well preserved with marked hyperplasia of the white pulp. Likewise, splenic red pulp was infiltrated with similar lymphocytes. Flow cytometry of these spleen atypical cells revealed positive findings for CD3, CD5, and CD8 and negative for CD4, CD10 and CD56. These malignant cells were positive for $\alpha\beta$ T-cell receptors (TCRs) and negative for $\gamma\delta$ TCRs. Clonal rearrangement of TCRs beta-chain was evident in the Southern blot analysis. The results of our investigations were consistent with a diagnosis of CAD secondary to hepatosplenic $\alpha\beta$ T-cell lymphoma.

She was treated with CHOP chemotherapy [2]. After 3 cycles, her hemoglobin level became sustained without transfusion, and the haptoglobin level reverted to a normal range. After 6 cycles of CHOP, a bone marrow biopsy revealed no residual lymphoma cells and a Southernblot analysis showed a germline of the

TCRs. Her serum IL-4 levels had decreased to 9.9 pg/ml. A complete remission (CR) was thus achieved. The cold agglutinin titer decreased to 1:256 thus suggesting an association between lymphoma and CAD. As HSTCL has a poor outcome, she underwent allogeneic peripheral blood stem cell transplantation with the use of a non-myeloablative conditioning regimen (25 mg/m² fludarabine monophosphate on days -6 to -2, 70 mg/m² melphalan on days -3 and -2) in July 2005 [3]. As the time this report is written, she has been in CR for 14 months.

In HSTCL, anemia has been described in many patients and it seems to be one of the prominent features of this rare type lymphoma. Splenomegaly, infiltration of the bone marrow by the malignant lymphocytes, and the release of various cytokines by the neoplastic cells have all been implicated as possible mechanisms for the cytopoenia in these patients [4]. In addition, hemolysis appears to be another possible mechanism seen in HSTCL. Among the different types of hemolytic anemia, the occurrence of autoimmune hemolytic anemia (AIH) secondary to non-Hodgkin lymphoma is well known [5].

To our knowledge, however, there have been only three cases of AIH secondary to HSTCL reported (Table 1) [1,6,7]. HSTCL is supposed to have a very aggressive clinical course [8]. However, three out of four cases with AIH, including ours, seemed to have some stable period in which only hemolytic anemia, thrombocytopenia or hepatosplenomegaly were evident without fever or general deterioration, which are all signs for the progression of lymphoma. In this stable period, those patients were treated with intravenous immunoglobulin administration or steroids, or they were just observed without being diagnosed as HSTCL (Table 1). Considering the poor outcome of HSTCL, we recommend that a proper diagnosis of

secondary AIH should be made in such patients so that the underlying HSTCL can be properly treated.

The pathogenesis of this AIH is mainly unknown. Although Motta et al proposed that $\gamma\delta$ T cells are attractive candidates as mediators of autoimmune disease in several conditions for both humans and laboratory animals [6], Lai et al already has reported AIH in HSTCL of the $\alpha\beta$ lineage [7]. This finding suggests that mechanisms other than pathological $\gamma\delta$ T cells of HSTCL in AIH should also be considered. Our case reinforced this concept. It is possible that the antibody production is the result of an aberrant immune interaction between the neoplastic T lymphocytes and the B-cell population. The Th-2 type cytokines IL-4 are produced from helper T cells, which usually express $\alpha\beta$ TCRs and stimulate B cells thus causing an expansion of B cells and the production of immunoglobulin. In our case, the levels of IL-4 decreased with the improvement of the hemolysis. IL-4 might therefore play some role in the productions of cold agglutinin in our case. However, more studies are needed to conclude a definite pattern of the serum IL-4 levels in patients who develop CAD with HSTCL $\alpha\beta$ subtype.

In summary, we herein presented a case of HSTCL of $\alpha\beta$ lineage complicated with CAD. We recommend to consider hemolysis as a cause of anemia in this type of lymphoma, or to consider lymphoma whenever one sees a patient with CAD.

Reference

1. Sallah S, Smith SV, Lony LC, et al. (1997) Gamma/delta T-cell hepatosplenic lymphoma: review of the literature, diagnosis by flow cytometry and concomitant autoimmune hemolytic anemia. *Ann Hematol* 74:139
2. McKelvey EM, Gottlieb JA, Wilson HE, et al. (1976) Hydroxyldaunomycin (Adriamycin) combination chemotherapy in malignant lymphoma. *Cancer* 38:1484-1493
3. Malladi RK, Peniket AJ, Norton AE, et al. (2004) Favourable outcome for patients with myeloid disorders treated with fludarabine-melphalan reduced-intensity conditioning and allogeneic bone marrow stem cell transplantation without the use of T-lymphocyte-depleting antibodies. *Eur J Haematol* 73:85-92
4. Dommann-Scherrer CC, Kurer SB, Zimmermann DR, et al. (1995) Occult hepatosplenic T-gamma delta lymphoma. Value of genotypic analysis in the differential diagnosis. *Virchows Arch* 426:629-634
5. Genty I, Michel M, Hermine O, et al. (2002) Characteristics of autoimmune hemolytic anemia in adults: retrospective analysis of 83 cases. *Rev Med Interne* 23:901-909
6. Motta G, Vianello F, Menin C, et al. (2002) Hepatosplenic gammadelta T-cell lymphoma presenting with immune-mediated thrombocytopenia and hemolytic anemia (Evans' syndrome). *Am J Hematol* 69:272-276
7. Lai R, Larratt LM, Etches W, et al. (2000) Hepatosplenic T-cell lymphoma of alphabeta lineage in a 16-year-old boy presenting with hemolytic anemia and thrombocytopenia. *Am J Surg Pathol* 24:459-463
8. Weidmann E. (2000) Hepatosplenic T cell lymphoma. A review on 45 cases since the first report describing the disease as a distinct lymphoma entity in 1990. *Leukemia* 14:991-997

Table 1 Reported cases of HSTCL with autoimmune hemolytic anemia

	Age	Sex	Subtype	D.Coombs	I.Coombs	CA	Initial diagnosis	B symptoms	Initial treatment	References
1	42	M	$\gamma\delta$	+	-	+	HSTCL	-	CHOP	Sallah et al[1]
2	61	M	$\gamma\delta$	+	+	NA	HSTCL	+	Steroids/CPA	Motta et al[6]
3	16	M	$\alpha\beta$	+	NA	NA	Evan's syndrome	-	IVIg, steroids	Lai et al[7]
4	56	F	$\alpha\beta$	+	-	+	CAD	-	Observation	Present case

M: Male, F: Female, D.Coombs: direct Coombs, I.Coombs: indirect Coombs, NA: Not available
 CA: Cold agglutinin, HSTCL: hepatosplenic T-cell lymphoma, CAD: Cold agglutinin disease
 CHOP: cyclophosphamide, adriamycin, vincristine, predonine, CPA: cyclophosphamide
 IVIG: intravenous immunoglobulin