



Title	DISTRIBUTION OF ANTIBODIES IN DOGS AGAINST ADENO-ASSOCIATED SATELLITE VIRUS ASSOCIATED WITH INFECTIOUS CANINE HEPATITIS VIRUS AND SEROLOGICAL TYPING OF THE SATELLITE VIRUS
Author(s)	ONUMA, Misao
Citation	Japanese Journal of Veterinary Research, 19(1-2), 40-40
Issue Date	1971-06
Doc URL	<a href="http://hdl.handle.net/2115/1975">http://hdl.handle.net/2115/1975</a>
Type	bulletin (article)
File Information	KJ00003418330.pdf



[Instructions for use](#)

**DISTRIBUTION OF ANTIBODIES IN DOGS AGAINST  
ADENO-ASSOCIATED SATELLITE VIRUS ASSOCIATED WITH  
INFECTIOUS CANINE HEPATITIS VIRUS AND  
SEROLOGICAL TYPING OF THE SATELLITE VIRUS**

Misao ONUMA

*Department of Hygiene and Microbiology  
Faculty of Veterinary Medicine  
Hokkaido University, Sapporo, Japan*

An adeno-associated satellite virus (ASV) associated with infectious canine hepatitis virus (ICHV) was found in this laboratory in 1968. This strain of ASV designated as M is the only ASV associated with ICHV. In the present paper, distribution of antibodies against ASV strain M is presented by the complement fixation (CF) test.

1) An attempt was first made to increase the yield of strain M. For this purpose, strain Matsuda with a high infectivity titer of ICHV was obtained by three times applying a limiting dilution experiment and was mixed with the original strain Matsuda, containing strain M, and inoculated into a dog kidney cell culture. The resultant viral material contained more ASV particles than the original strain Matsuda.

2) About 10<sup>-1</sup> of strain M material thus obtained was finally concentrated and purified by banding three times in isopycnic ultracentrifugation in CsCl gradient. The purified strain M did not react with anti-ICHV dog serum but reacted with the antisera against strain Matsuda containing strain M. The anti-strain M serum was prepared by inoculating the purified strain M into guinea pigs.

3) Distribution of antibodies to the strain M in dogs was then examined. Eighty-four dog sera collected in Sapporo were titrated, beginning at 1:2 dilution, by CF test (microtiter technique) using 4 units of the antigen of strain M. Eight out of 38 new-born puppies (less than 10 days-old) were positive, none out of 7 young puppies (about a month-old), and 14 out of 39 adult dogs were positive. A total of 26.2% of the dogs possessed antibodies against strain M. Antibodies against ICHV were found in 17 out of 38 new-born dogs, in none out of 7 puppies, and in 16 out of 39 adult dogs. ICHV antibody was found in 72.7% of the dog sera possessing the strain M antibody. The antibody titer against strain M was usually lower than that of ICHV. From these findings, it is presumed that natural infection of strain M among dogs is possible and, in such cases, multiplication of the ASV in dogs will require coinfection of ICHV.