



Title	ON THE RELATIONSHIP BETWEEN THE EXCITATION OF NON-ADRENERGIC INHIBITORY NERVE AND THE RELEASE OF ADENINE NUCLEOTIDES
Author(s)	KOHDA, Tadayuki
Citation	Japanese Journal of Veterinary Research, 24(3-4), 110-110
Issue Date	1976-10
Doc URL	http://hdl.handle.net/2115/2093
Type	bulletin (article)
File Information	KJ00003407793.pdf



[Instructions for use](#)

**ON THE RELATIONSHIP BETWEEN THE
EXCITATION OF NON-ADRENERGIC INHIBITORY NERVE
AND THE RELEASE OF ADENINE NUCLEOTIDES**

Tadayuki KOHDA

*Department of Veterinary Pharmacology
Faculty of Veterinary Medicine
Hokkaido University, Sapporo, Japan*

An attempt was made to clarify whether ATP or a related adenine nucleotide is the transmitter substance of the non-adrenergic inhibitory nerve. Isolated preparations of guinea-pig stomach-vagus nerves were perfused with a nutrient media via the coeliac artery. A quantitative measurement of adenine nucleotides (ATP, ADP and AMP) in the perfusate following vagal stimulation was made using the firefly luminescence method.

1) Vagal stimulation invariably caused a significant ($P < 0.05$) increase in the adenine nucleotides release associated with gastric contraction, and some correlation between the release and the frequency of the stimulation could be found. After the addition of atropine (5×10^{-7} g/ml) to the media, only a small increase in the nucleotides was found with gastric relaxation, and there was no relationship between the release and the frequency of the stimulation.

2) Under perfusion with a solution containing hemicholinium-3 (2×10^{-5} g/ml) and hexamethonium (1×10^{-6} g/ml), vagal stimulation elicited neither contractions nor an increase of ATP, while ACh (10^{-5} g/ml) infusion caused both effects.

3) When the stomach was perfused with a hypertonic solution, the contractile response to ACh and vagal stimulation was abolished or greatly reduced. At this time, any increase in the adenine nucleotides could not be detected.

4) From these results, the most part of the increase in the adenine nucleotides can not be attributed to a nervous origin, and therefore, the hypothesis that ATP or a related compound is the transmitter substance of the non-adrenergic inhibitory nerves can not be supported.