



Title	STUDIES ON THE RELATIONSHIP BETWEEN THE SERUM GAMMA GLOBULIN LEVELS OF NEONATAL PIGLETS AND THEIR MORTALITY OR GROWTH DURING THE FIRST TWO MONTHS OF AGE : AN EVALUATION FOR THE AMMONIUM SULPHATE TURBIDIMETRIC METHOD
Author(s)	YAGUCHI, Hideyuki
Citation	Japanese Journal of Veterinary Research, 27(1-2), 36-36
Issue Date	1979-04
Doc URL	http://hdl.handle.net/2115/2173
Type	bulletin (article)
File Information	KJ00003407895.pdf



[Instructions for use](#)

**STUDIES ON THE RELATIONSHIP BETWEEN THE SERUM GAMMA
GLOBULIN LEVELS OF NEONATAL PIGLETS AND THEIR
MORTALITY OR GROWTH DURING THE FIRST
TWO MONTHS OF AGE: AN EVALUATION
FOR THE AMMONIUM SULPHATE
TURBIDIMETRIC METHOD**

Hideyuki YAGUCHI

*Department of Veterinary Internal Medicine
Faculty of Veterinary Medicine
Hokkaido University, Sapporo 060, Japan*

In order to examine the relationship between the serum gamma globulin levels of neonatal piglets and their mortality or growth during the first 2 months of age, serum total protein concentrations, electrophoretic fraction concentrations, and turbidity readings (Kunkel's units) by the ammonium sulphate turbidity readings (Kunkel's units) by the ammonium sulphate turbidimetric method were investigated for 266 piglets in 27 litters at 12~18 hours after birth.

The results were as follows ;

1) Wide variations in serum total proteins (from 2.7 to 8.6 g/100 ml), serum protein profiles, and Kunkel's units (from 0 to 44 units) were found in individual piglets.

2) A significant correlation ($r=0.9663$, $p<0.001$) was found between serum gamma globulin concentrations and Kunkel's units, indicating that the ammonium sulphate turbidimetric method is useful for the estimation of serum gamma globulin levels.

3) Kunkel's units of piglets were generally influenced by the birth order and birth weight within the same litter. Particularly, it was noted that low weight piglets at birth in the large litter size tended to give low Kunkel's units. It is considered that such piglets ingested little or no colostrum due to the competition among littermates at suckling.

4) The rate of loss (death and stunt) during the first 2 months of age was high in the piglets with low Kunkel's units and in low weight piglets at birth.

In conclusion, one of the most important causes of loss in baby pigs is perhaps that low weight piglets at birth tend to be hypogammaglobulinaemia during the neonatal period.