



Title	EXPERIMENTAL STUDIES ON BLOOD TRANSFUSION INTO DOGS BLOOD GROUPS AND CLINICAL AND HEMATOLOGICAL FINDINGS ON TRANSFUSION USING ERYTHROCYTES
Author(s)	YAMAMOTO, Keiko
Citation	Japanese Journal of Veterinary Research, 28(1-2), 52-52
Issue Date	1980-05-31
Doc URL	http://hdl.handle.net/2115/2197
Type	bulletin (article)
File Information	KJ00003407922.pdf



[Instructions for use](#)

collected in Biei district near Mt. Tokachi in the central area. The antibody titer was 1:10 to Negishi, but was not tested to other flaviviruses.

**EXPERIMENTAL STUDIES ON BLOOD TRANSFUSION INTO
DOGS BLOOD GROUPS
AND
CLINICAL AND HEMATOLOGICAL FINDINGS ON
TRANSFUSION USING ERYTHROCYTES**

Keiko YAMAMOTO

*Department of Veterinary Surgery
Faculty of Veterinary Medicine
Hokkaido University, Sapporo 060, Japan*

The blood groups in dogs were studied to make a basic investigation of the clinical blood transfusion in dogs. Immune antibodies were produced in dogs and goats by repeated infusions of canine erythrocytes. The blood groups of one hundred thirty-six random mongrel dogs caught in Sapporo were examined.

Furthermore, for the purpose of observing the clinical and hematological findings produced in dogs by the transfusion of incompatible erythrocytes, a 45% suspension of canine compatible or incompatible erythrocytes in physiological saline solution was transfused into dogs twice at 10-day intervals. The amount of cell suspension transfused was equivalent to 15 percent of the total blood volume.

The following results were obtained :

1) Four types of antibodies obtained from the dogs were tentatively designated as anti-S₁, S₂, S₃, and S₄. One type of antibody obtained from a goat was tentatively designated as anti-Sg.

2) The incidence of antigens S₁ through Sg among the 136 dogs was as follows: antigen-S₁ 72.1% ; S₂ 41.2% ; S₃ 22.8% ; S₄ 69.9% ; and Sg 48.5%.

3) The incidence of combinations of blood group antigens was as follows: 12.5% of all of the examined dogs had none of these 5 blood typing antigens; 17.0% had only one antigen; 17.6% had two antigens; 19.1% had three antigens; 24.2% had four antigens; and 9.6% had all of these antigens.

4) Twenty kinds of combinations were observed in these 136 dogs. The highest frequency was 19.8% observed in the combination S₁ S₂ S₄ Sg, and the lowest frequency was 0.7% observed in the combination S₁ Sg, S₁ S₂, S₃ S₄ Sg, and S₁ S₃ S₄ Sg.

5) After the first transfusion, there was no difference between the compatible and incompatible erythrocytes transfusion in the clinical findings. After the second transfusion, however, one fifth of the dogs transfused with the incompatible erythrocytes showed clinical findings of intensive shock.