



Title	ISOLATION OF NON-PILIATED CLONE OF CORYNEBACTERIUM RENALE STRAIN 115
Author(s)	OKAMOTO, Ryoichi; TAKAI, Shinji; YANAGAWA, Ryo; SATO, Hisaaki
Citation	Japanese Journal of Veterinary Research, 29(3-4), 79-82
Issue Date	1981-12-05
DOI	10.14943/jjvr.29.3-4.79
Doc URL	http://hdl.handle.net/2115/2239
Type	bulletin (article)
File Information	KJ00003436417.pdf



[Instructions for use](#)

**ISOLATION OF NON-PILIATED CLONE OF
CORYNEBACTERIUM RENALE
STRAIN 115**

Ryoichi OKAMOTO, Shinji TAKAI, Ryo YANAGAWA
and Hisaaki SATO

*Department of Hygiene and Microbiology
Faculty of Veterinary Medicine
Hokkaido University, Sapporo 060, Japan*

(Received for publication, September 18, 1981)

A non-piliated (P^-) clone was isolated from 368 colonies of a densely piliated (P^+) clone of *Corynebacterium renale* strain 115 by means of a very weak agglutination with anti-pili serum. Lack of pili in P^- was confirmed by immunodiffusion and electron microscopy.

INTRODUCTION

Corynebacterium renale was found to be the first Gram-positive bacteria which possessed pili^{9,10}. The pili of *C. renale* have been studied in the authors' laboratory morphologically^{9,10}, chemically^{4,5}, and immunologically^{4,5,9}. Adhesion properties of the piliated corynebacteria have been reported^{1,2,3}. Clones of *C. renale* possessing numerous and no pili have been thought to be essential for studies of the function of the pili in the bacteria. The present communication deals with isolation of a non-piliated (P^-) clone from a densely piliated (P^+) clone of *C. renale* strain 115.

MATERIALS AND METHODS

Strain *C. renale* strain 115 was used. This strain possessed pili¹¹, agglutinated trypsinized sheep erythrocytes¹¹, and adhered to tissue cultured cells⁹ and the mucous membrane of the urinary bladder of mice⁹.

Medium and cultivation The organisms were cultivated on nutrient agar (pH 7.2) at 37°C for 24 hours.

Antiserum to *C. renale* Antisera against *C. renale* were prepared as reported⁸.

Anti-pili serum Anti-pili serum was prepared as previously reported¹¹. Anti-strain 115 serum was absorbed with the organisms of *C. renale* ATCC 19412, which possessed only a few pili, and the somatic antigen identical with strain 115. The cells of ATCC 19412 collected from the culture on the agar medium were mixed and incubated with 1 ml of anti strain 115 serum at 37°C for 2 hours. The mixture was centrifuged, and the supernatant was filtered through a membrane filter (Millipore Corp., 450 nm pore size). The procedure was repeated until the absorption was completed, which was confirmed by immunodiffusion.

Slide agglutination test using anti-pili serum One drop of the anti-pili serum was mixed on a slide glass with a drop of saline suspended with bacteria. Normal rabbit serum was used instead of the anti-pili serum for control. Agglutination was recorded as positive soon after the mixing.

Purification of pili Pili of P⁺ was partially purified according to the method described previously⁹⁾.

Electron microscopy Examination by electron microscopy was done as reported⁹⁾.

Immunodiffusion Antigens for immunodiffusion were extracted from organisms of *C. renale* with acid⁶⁾; bacteria were suspended in 1 ml of saline containing N/20 HCl, bathed in boiling water for 15 min, cooled, and after being sedimented by centrifugation, the supernatant was used as the antigen after being neutralized with 1 N NaOH.

RESULTS

Seven arbitrarily selected colonies of *C. renale* strain 115 were found without exception to be strongly agglutinated by the anti-pili serum. And electron microscopy revealed that they possessed pili. One of the arbitrarily selected clones which cloned twice was again examined; all 15 colonies of this clone were agglutinated by the anti-pili serum and possessed pili. One of the colonies (s-13-1-5) was designated as P⁺; its electron micrograph is shown in figure 1.

The colonies of P⁺ were examined by slide agglutination using anti-pili serum to determine if any were not agglutinated by the antiserum. Of the 368 colonies examined, only one was very weakly agglutinated by the anti-pili serum, and it appeared only after a prolonged period of time after mixing. Electron microscopically, the organisms of this colony did not possess pili (fig. 2). This clone (s-13-1-5-P⁻-16) was designated as P⁻.

There was no difference between the colonial features of P⁺ and P⁻; both colonies were smooth, slightly yellowish, glistening and moist.

In immunodiffusion using anti-P⁺ serum, P⁺ produced 3 lines while P⁻ formed 2 lines, which were common to those formed by P⁺. P⁻ lacked the third line produced by the P⁺ adjacent to the antigen well (fig. 3). This line corresponded to that produced by purified pili (fig. 4). ATCC 19412, which possessed only a few pili on a very limited number of organisms, also lacked the line produced by the pili.

DISCUSSION

Since their discovery in 1968¹⁰⁾, the pili of *C. renale* have been studied morphologically^{9,10)}, chemically^{4,5)}, and immunologically.^{4,5,9)} Adhesion of *C. renale* to trypsinized sheep erythrocytes, tissue cultured cells and the mucous membrane of the urinary bladder of mice mediated by the pili was disclosed by inhibition of the adhesion by anti-pili serum.^{1,2,3)} These experiments were done, however, using densely and slightly piliated

strains of *C. renale*. If we could have used piliated and non-piliated clones derived from the same strain, the results would probably have been clearer.

The P⁻ clone of *C. renale* strain 115, which was isolated by means of very weak agglutination with anti-pili serum and verified by immunodiffusion and electron microscopy in the present study, will be a useful tool for studies on the biological function of the pili of *C. renale*. As a matter of fact, our recent study of pH dependent adhesion of piliated *C. renale*ⁿ was completed with the aid of P⁻.

REFERENCES

- 1) HONDA, E. & YANAGAWA, R. (1974): Agglutination of trypsinized sheep erythrocytes by the pili of *Corynebacterium renale* *Infect. Immun.*, **10**, 1426-1432
- 2) HONDA, E. & YANAGAWA, R. (1975): Attachment of *Corynebacterium renale* to tissue culture cells by the pili *Am. J. Vet. Res.*, **36**, 1663-1666
- 3) HONDA, E. & YANAGAWA, R. (1978): Pili-mediated attachment of *Corynebacterium renale* to mucous membrane of urinary bladder of mice *Ibid.*, **39**, 155-158
- 4) KUMAZAWA, N. & YANAGAWA, R. (1972): Chemical properties of the pili of *Corynebacterium renale* *Infect. Immun.*, **5**, 27-30
- 5) KUMAZAWA, N. & YANAGAWA, R. (1973): Comparison of the chemical and immunological properties of the pili of three types of *Corynebacterium renale* *Jpn. J. Microbiol.*, **17**, 13-19
- 6) LANCEFIELD, R. C. (1928): The antigenic complex of *Streptococcus haemolyticus* I. Demonstration of a type-specific substance in extracts of *Streptococcus haemolyticus* *J. Exp. Med.*, **47**, 91-103
- 7) TAKAI, S., YANAGAWA, R. & KITAMURA, Y. (1980): pH-dependent adhesion of piliated *Corynebacterium renale* to bovine urinary bladder epithelial cells *Infect. Immun.*, **28**, 669-674
- 8) YANAGAWA, R., BASRI, H. & OTSUKI, K. (1967): Three types of *Corynebacterium renale* classified by precipitin reaction in gels *Jpn. J. Vet. Res.*, **15**, 111-120
- 9) YANAGAWA, R. & OTSUKI, K. (1970): Some properties of the pili of *Corynebacterium renale* *J. Bacteriol.*, **101**, 1063-1069
- 10) YANAGAWA, R., OTSUKI, K. & TOKUI, T. (1968): Electron microscopy of fine structure of *Corynebacterium renale* with special reference to pili *Jpn. J. Vet. Res.* **16**, 31-38

EXPLANATION OF PLATE

- Fig. 1 Electron micrograph of P⁺ (×20000)
- Fig. 2 Electron micrograph of P⁻ (×20000)
- Fig. 3 Immunodiffusion of anti-P⁺ serum (central well) with the antigens of P⁺ and P⁻
- Fig. 4 Immunodiffusion of anti-P⁺ serum (central well) with the antigens of P⁺, P⁻, ATCC 19412 and purified pili of P⁺

