

| Title | DIATOM COMMUNITIES IN WESTERN ALEUTIAN WATERS ON THE BASIS OF NET SAMPLES COLLECTED IN MAY-JUNE 1953 |
|------------------|---|
| Author(s) | MOTODA, Sigeru; KAWARADA, Yutaka |
| Citation | 北海道大學水産學部研究彙報, 6(3), 191-200 |
| Issue Date | 1955-11 |
| Doc URL | http://hdl.handle.net/2115/22926 |
| Туре | bulletin (article) |
| File Information | 6(3)_P191-200.pdf |



DIATOM COMMUNITIES IN WESTERN ALEUTIAN WATERS ON THE BASIS OF NET SAMPLES COLLECTED IN MAY-JUNE 1953*

Sigeru MOTODA and Yutaka KAWARADA

Faculty of FisheriesHakodate MarineHokkaido UniversityObservatory

The present studies are based on materials collected by the Training Ship "Oshoro Maru" of the Faculty of Fisheries, Hokkaido University, during her training and research cruise to the western Aleutian waters from May to June 1953. The main purpose of the studies is to get the figure of distribution of diatom communities in upper layers of these areas which are certainly bounded by particularities of the water system.

The plankton net used in samplings measures 20 cm in mouth diameter and 68 cm in length along the side; it is made of fine bolting silk, XX 16, i. e., 157 meshes a linear inch. Samplings were made by 50 metre approximately vertical hauls.

The positions and dates of the samplings were quite the same as those of zooplankton samplings (fig. 1 and appendix; cf. Anraku, 1954). The number of stations, however, is different from those given in the hydrographic studies by Mishima & Nishizawa (1955), so that in referring to their hydrographic data care is necessary to read degrees of latitude and longitude of the stations, but not the number of the stations.

In the areas investigated the diatom flora is comparatively rich in species including the following nine leading species:

| Chaetoceros | atlanticus | Corethron hystrix |
|-------------|------------|------------------------------------|
| Ch. | convolutus | Denticula sp. |
| Ch. | debilis | Nitzschia seriata |
| Ch. | decipiens | Rhizosolenia hebetata f. semispina |
| Ch. | radicans | |

In addition to these species, *Coscinodiscus oculus iridis*, *Thalassiosira decipiens*, *Th. nordenskiöldii* and *Thalassiothrix longissima* are more or less abundant through the entire area, and *Nitzschia closterium* is also sometimes rather abundant though in restricted location. A few species of dinoflagellates and tintinnodes are present too but in very small quantities.

Although the quantitative values of the present samples are limited because of difficulty of exact estimation of filtration coefficient of the net, counting of the cell numbers of diatoms is made to get an approximate figure of density of population

^{*} おしよろ丸北洋調査報告 No. 3 (1953年度)

which, together with the composition of the population, might have bearing upon the hydrography of the areas.

Illustrated in figures 2-6 respectively is the quantitative distribution of diatoms as given by the cell numbers of total communities as well as by those in percentage of four major groups separately, i. e., *Chaetoceros-Hyalochaete*, *Chaetoceros-Phaeoceros*, *Nitzschia seriata* and *Rhizosolenia hebetata* f. semispina.

From the characteristics of the diatom communities in respect to their density as well as to their composition, the areas investigated may be divided into six regions: they are symbolized by A, B, C, D, E and F in figure 1 and table 1.

| | Regions | А | В | С | D | Е | F | |
|-----------------------------------|--|-------|-------------------|-------|-------------------|-------------------|-------------------|--|
| Tota | l number of diatom cells per m ³ per a station | 3×107 | 3×10 ⁴ | 6×104 | 3×10 ⁵ | 8×10 ⁴ | 4×10 ⁵ | |
| Percentage of six major groups | Chaetoceros – Hyalochaete | 82 | 41 | 0 | 7 | 17 | 2 | |
| | Chaetoceros—Phaeoceros | 6 | 15 | 69 | 67 | 8 | 60 | |
| | Corethron hystrix | + | + | 13 | 2 | 7 | 8 | |
| | Denticula sp. | 1 | 6 | 16 | 5 | 2 | 4 | |
| | Nitzschia seriata | 2 | 12 | + | 4 | 50 | 4 | |
| | Rhizosolenia hebetata f. semispina | 2 | 24 | + | 11 | 13 | 19 | |
| | Diatoms miscellaneous | 7 | 2 | 2 | 4 | 3 | 3 | |

Table 1. Abridged table showing characteristics of diatom communities in six regions

A Region

The A region covering St. 1, 4-8, 10 and 20 around the Near Islands and Rat Islands is densely populated with diatoms which are present to the number of more than one million cells per 1 m³ of water. *Chaetoceros-Hyalochaete* group, mainly composed of *Chaetoceros constrictus*, *Ch. compressus*, *Ch. debilis*, *Ch. decipiens*, *Ch. radicans* and *Ch. teres*, are most prevalent. Every station in this region shows the occurrence of *Hyalochaete* in the bulk more than 60 per cent of the total numbers of diatoms; especially on the course from St. 4 to St. 8 this group is remarkably predominant, forming more than 80 per cent of the total. Other diatoms, such as *Chaetoceros convolutus*, *Nitzschia closterium*, *Nitz. seriata*, *Rhizosolenia hebetata* f. *semispina*, *Thalassiosira nordenskiöldii*, *Thalassiothrix longissima*, etc., are also found to be prevalent though to a less extent.

It is supposed from the fact of the richness in diatom population as well as of the nearly monotonous presence of fine species of Subgenus *Hyalochaete* that the waters in **A** region are much influenced by neritic elements. Low transparency and green colour of water have been reported in this region (Taguchi & Hirose, 1954).

It has been found that a warm water mass of low salinity moves westward along the southern side of the Aleutian Chain from far to the east and turns into the Bering

1955) Motoda & Kawarada: Diatom Communities in Aleutian Waters

Sea passing through the channels between the Aleutian Islands (Barnes & Thompson, 1938; Sverdrup *et al.*, 1942; Watanabe, 1954; Mishima & Nishizawa, 1955). The communities found in A region resemble to certain extent those of the neritic waters in the Bering Sea on the coast of Alaska Peninsula which fact has been reported by Aikawa (1938).

B Region

In the **B** region covering St. 11 and 12 the total population of diatoms is very small, less than 1000000 cells per 1 m³ of water. The leading forms are *Hyalochaete*, among which *Chaetoceros decipiens* prevails, as in **A** region but in less percentage. On the other hand, *Phaeoceros* including *Chaetoceros convolutus*, *Nitzschia seriata* and *Rhizosolenia hebetata* f. *semispina*, become relatively abundant. Thus the **B** region is characterized by the paucity of total diatom population, the occurrence of *Phaeoceros* and relative decrease in number of *Hyalochaete*. These facts will indicate that warm water of low salinity from **A** region here contacts and mingles with the cold oceanic water from **D** region; that is the **B** region is an intermediate mixing area between **A** and **D** regions.

C Region

The C region covers St. 2 and 4. Although the amount of total diatoms is very small as in **B** region, the simplicity of communities and the predominance of *Phaeoceros* clearly show that water of this region differs from neritic water. *Phaeoceros* including *Chaetoceros atlanticus*, *Ch. concavicornis*, *Ch. convolutus*, etc., of which the predominant species is *Ch. convolutus*, occupy more than 55 per cent of total number of diatoms, and in addition, typical cold water species, *Corethron hystrix* and *Denticula* sp. are present, while *Hyalochaete* is entirely absent. These facts indicate that the water in this region is purely oceanic. The water in this region has been reported to be very transparent, more than 20 metres of Secchi disc reading (Taguchi & Hirose, 1954). Although Taguchi & Hirose (1954) and also Mishima & Nishizawa (1955) have judged this region to be the continuation of Kuroshiwo extension, the diatom communities observed here suggest that the surface water of this region is more like to that of cold oceanic waters of **D** region.

D Region

The **D** region covering St. 13, 15, 19 and 21, contains diatom cells varying from 7500 to 800000 per 1 m^3 of water, showing thus a more dense population than in **B** or **C** region. The communities are characterized by the predominance (60 per cent of total diatoms) of *Phaeoceros* mainly comprising *Chaetoceros atlanticus* and *Ch. convolutus*.

-193-

Together with these species, Corethron hystrix, Denticula sp. and Rhizosolenia hebetata f. semispina are also abundantly found here. These communities quite resemble those in C region far to the south-east, both reflecting the cold oceanic nature of the water. It has been suggested from the observations on the drift course of ice bergs that there might exist a surface current from the northern Bering Sea to near the western Aleutian Chain through the northeastern offing of Kamchatka (Hokkaido-Cho, 1906).

It is known that diatom communities in wide areas of the Bering Sea are represented by *Phaeo* plankton associations (Aikawa, 1932, 1933, 1935, 1936a, 1936b, 1938, 1940; Kanno, 1936). The similarity of communities between **D** region and those in a large part of the Bering Sea and the suggestion of a surface current above mentioned will indicate that this region has a water mass very similar to that of the northern reaches of the Bering Sea. So far as the surface water is concerned the water mass in this region seems from the present data to be of different origin from that suggested by the results of the hydrographic studies of Mishima & Nishizawa (1955).

E Region

The E region covers St. 17, 18 and 23. The count of diatoms is relatively low, being only about 100000 cells per 1 m³ of water. *Nitzschia seriata* is prevalent and *Chaetoceros debilis* is present in considerable number, while *Phaeoceros* remarkably decrease in occurrence. These communities are typically represented at St. 17 and 23, indicating the neritic nature of the water to certain extent, while at St. 18 certain effects of cold oceanic water are proved by the presence of *Nitzschia seriata* and *Chaetoceros convolutus* in about equal number and also by the abundant occurrence of *Rhizosolenia hebetata* f. *semispina*.

Although both A and E regions are shown to be of neritic nature in respect to diatom communities, there is a wide difference between the two in the composition of communities, indicating that the water comes from different origins. The presence of *Nitzschia seriata* in considerable number in E region suggests that the water in this region is influenced by the coastal waters near Kamchatka, because this species has been reported to be the most prevalent one in the neritic waters on both the east and the west coasts of Kamchatka (Aikawa, *loc. cit.*; Tsuruta & Chiba, 1954).

F Region

The F region covers St. 16 and 24. The water is populated with diatoms in considerable density. The most predominant species is *Chaetoceros convolutus*. *Corethron hystrix* and *Rhizosolenia hebetata* f. *semispina* are also fairly numerous. Curiously *Nitzschia seriata* is hardly observed, numbering only about 4 per cent of the total diatoms. Thus the communities in this region show a close resemblance with those of

-194--

Motoda & Kawarada: Diatom Communities in Aleutian Waters

1955)

C and D region reflecting the presence of cold oceanic water, but not with those of E region. The limitation of areas investigated leaves unsolved the origin of the water in F region.

There are certain facts of disaccord in diatom associations in the present areas between the results of the preceding investigators and the present data. That is, (1) *Thalassiothrix, Coscinodiscus* and *Thalassiosira* have been reported to be the most representative forms in the Aleutian waters, but in the present investigations they appear only in small quantities; (2) *Chaetoceros atlanticus* is known to be the most dominant and wide spread species in these areas, but it is abundantly collected only at one station in the present cruise; (3) on the other hand, *Hyalochaete* group are widely found in the present cruise, while the previous investigators have reported that the distribution of this species is certainly localized.

Summary

- 1. The diatoms are comparatively abundant, as a whole, in the upper layers of the areas investigated, occurring about one hundred million cells per 1 m³ of water in the maximum, about ten million in average and about several thousand in the minimum.
- 2. The leading species are Chaetoceros atlanticus, Ch. convolutus, Ch. debilis, Ch. decipiens, Ch. radicans, Corethron hystrix, Denticula sp., Nitzschia seriata and Rhizosolenia hebetata f. semispina.
- 3. From the characteristics of the diatom communities six regions are distinguishable in the investigated areas (fig. 1).
- 4. The A region is characterized by *Chaeto*-plankton associations, demonstrating the neritic nature of the water.
- 5. The B region presents mixed communities of *Chaeto*-plankton, *Phaeo*-plankton, *Rhizosolenia hebetata* f. *semispina* and *Nitzschia seriata*, indicating that the water is mixed with cold oceanic water to a certain extent.
- 6. The C region is also represented by *Phaeo*-plankton associations, indicating the purely cold and oceanic nature of the water.
- 7. The D region is also represented by *Phaeo*-plankton associations, indicating cold oceanic water like that in the C region.
- 8. The E region is mainly occupied by *Nitzschia seriata*, having similar communities to those of the neritic water off Kamchatka that have been reported by the preceding investigators.
- 9. The F region is indicated to be oceanic by diatom communities.

The authors are gratefully obliged to Dr. Y. Takenouti of the Hakodate Marine

-195-

Observatory for his kind criticism of the oceanographic considerations in the present studies and to Mr. S. Nishizawa of the University for his kindness in giving many valuable advices during the studies. Cordial thanks are due to Captain S. Mishima and his crew aboard the "Oshoro Maru" for their courtesy in providing facilities for samplings and also to Mr. N. Ogawa, a student of the University at that time, for his help in the laborious work of making the samplings at sea.

Literature cited

Aikawa, H. (1932). On the summer plankton in the waters of the west Aleutian Islands in 1928. Bull. Jap. Soc. Sci. Fish. 1 (2), 70-74. (in Japanese).

(1933). On the planktology of Okhotsk Sea in autumn. Ibid. 2 (4), 175 182. (in Japanese).

(1935). On the quantitative analysis of the plankton association in the adjacent seas of Japan III. Jour. Imp. Fish. Exp. Sta. (6), 131-172. (in Japanese).

(1936a). The planktological properties of the principal sea areas surrounding Japan. Bull. Jap. Soc. Sci. Fish. 5 (1), 33-41. (in Japanese).

(1936b). On the diatom communities in the waters surrounding Japan. Rec. Oceanogr. Wks. Jap. 3 (1), 1-159.

(1938). On the quantitative analysis of the plankton associations in the adjacent seas of Japan V. Jour. Imp. Fish. Exp. Sta. (9), 67-86. (in Japanese).

(1940). (On the plankton associations in the Bering Sea and the Okhotsk Sea]. Kaiyô-Gyogyô 5 (1), 20 31. (in Japanese).

- Anraku, M. (1954). Gymnoplea copepoda collected in Aleutian waters in 1953. Bull. Fac. Fish., Hokkaido Univ. 5 (2), 123-136.
- Barnes, C. A. & Thompson, T. G. (1938). Physical and chemical investigations in Bering Sea and portions of the North Pacific Ocean. Univ. Wash., Publ. Oceanogr. 3 (2), 35-79.
- Hokkaido-Cho Daisambu (1906). (Report of the investigations of drifting ice-bergs in Hokkaido). (in Japanese).
- Kanno, R. (1935). The distribution of plankton during summer in the South Okhotsk Sea and on the coast of Kamchatka. Jour. Fish., Hakodate Coll. Fish. (38), 22-32. (in Japanese).
- Mishima, S. & Nishizawa, S. (1955). Report on hydrographic investigations in Aleutian waters and the southern Bering Sea in the early summer of 1953 and 1954. Bull. Fac. Fish., Hokkaido Univ. 6 (2), 84-124.
- Sverdrup, H. U., Johnson, M. W. & Fleming, R. H. (1942). The oceans: their physics, chemistry and general biology. 1087p. N. Y. Prentice-Hall Inc.
- Taguchi, K. & Hirose, Y. (1954). Surface current and oceanographical condition in the northern Pacific salmon fishing ground. Bull. Jap. Soc. Sci. Fish. 20 (7), 576-580. (in Japanese).
- Tsuruta, A. & Chiba, T. (1954). On the distribution of plankton at the fishing ground of salmons in the North Pacific, 1952. Jour. Shimonoseki Coll. Fish. 3 (3), 39-45. (in Japanese).
- Watanabe, N. (1954). A report on oceanographical investigations in the salmon fishing grounds of the North Pacific, 1952 and 1953. Special Publ. Tokai Reg. Fish Res. Lab. (3), 1-5. (Mimeogr.).

-196-



Fig. 1. Approximate sampling positions (1-8, 10-13, 15-21, 23-24) and demarcation of regions (A-F) which are set according to the characteristics of diatom communities



Fig. 2. Isotherms and isohalines at the surface (Mishima & Nishizawa, 1955)

-197-

1955)



Fig. 3. Horizontal distribution of total numbers of diatoms



Fig. 4. Occurrence of *Chaetoceros-Hyalochaete* in percentage of total diatoms in each haul



Fig. 5. Occurrence of *Chaetoceros-Phaeoceros* in percentage of total diatoms in each haul



Fig. 6. Occurrence of *Rhizosolenia hebetata* f. semispina in percentage of total diatoms in each haul

1955)

Bull. Fac. Fish., Hokkaido Univ.



Fig. 7. Occurrence of Nitzschia seriata in percentage of total diatoms in each haul

Appendix: Full data of diatom count.

| | | | | | • | | | | | | | | والمراجع والمتكاف والمتشاط ومرارك ومراجعات | | المحمد الم | - | | | | | |
|---|------------------|---------------------------------------|-----------------------------|------------|-----------------|-------------|----------------|-------------|-----------|--|-----------|----------|--|-----------|--|---------------------------------|----------|-------------|-------------|-----------|-----------|
| A Station | 1 | 2 | 3 | 4 | 5 | 6 | 7 | 8 | 10 | 11 | 12 | 13 | 15 | 16 | 17 | 18 | 19 | 20 | 21 | 23 | 24 |
| P. Logation Lat. | 49°22′ N | 48°57′ N | 50°00' N | 50°58' N | 52°36' N | 53°22′ N | 54°00′ N | 53°50.5' N | 53°42′ N | 52°15' N | 51°53′ N | 49°33′ N | 51°28′ N | 50°55.5'N | 50°52′ N | 49°50.8' N | 49°24' N | 50°51.5′ N | 49°48′ N | 50°03′ N | 49°55' N |
| D Location Long. | 178°18′ E | 178°49′W | 178°44′W | 178°40.5'W | 178°46′W | 179°27′E | 178°57′ E | 175°24.5′ E | 172°09′ E | 172°07′E | 170°23'E, | 167°01′E | 166°24′ E | 161°23′E | 163°18′ E | 166°46.2′E | 172°00′E | 172°00′ E | 170°22′ E | 163°32′E | 160°15′ E |
| C Date | V-15 | V—15 | V16 | V-16 | V-17 | V—17 | ∇ −18 | V—19 | V20 | V20 | V—21 | V24 | V-27 | VI—1 | VI-1 | VI2 | VI-5 | VI-6 | VI-10 | VI16 | VI—17 |
| D Time | 1:20 | 14:00 | 2:00 | 10:30 | 3:00 | 14:45 | 8:30 | 10:30 | 8:00 | 20:30 | 19:00 | 14:30 | 17:00 | 8:00 | 20:00 | 20:00 | 16:00 | 13:00 | 19:30 | 19:00 | 19:00 |
| E Hauled distans (m) . | 0-50 | 0-50 | 0-50 | 0-50 | 0-50 | 0~50 | 0~50 | 0-50 | 0-50 | 0-50 | 0-50 | 0-50 | 0-50 | 0-50 | 0~50 | 0-50 | 0-50 | 0-50 | 0-50 | 0 50 | 0-50 |
| F Settling volume of total | 0.9 | 4.0 | 3.5 | 20.8 H | 52.5 | 15.4 | 11.3 | 12.5 | 0.8 | 0.3 3 | | 1.3 | 4.7 | 1.2 | 0.6 | | 2.9 | 8.4 | 5.0 | 0.3 | 3.2 |
| sample (cc/m ³) | | | | 1000000 | | | | | | | | 1.0 | | , | | | 10700 | 1000000 | | 1 01 500 | 100000 |
| G Number of diatom cells/m ³ | 141750 | 39800 | 73500 | 42800000 3 | 123540000 | 21210000 | 2200000 | 21890000 | 238000 | 41850 | 10500 | 7550 | 125600 | 17700 | 37250 | 28750 | 12700 | 18830000 | 777000 | 161700 | 1038000 |
| | No. % | No. % | No. % | No % | No. % | No. % | No. % | No. % | No. % | No. % | No. % | No. % | No. % | No. % | No. % | No. % | No. % | No. % | No. % | No. % | No. % |
| Diatomeae | 100 | 100 | 100 | 1 | 100 | 100 | 100 | 100 | | | | 100 | 700 | 100 | 100 | 100 | 100 | 100 | 100 | 100 | 100 |
| · | 100 | 100 | 100 | | 100 | 100 | 100 | 100 | 100 | 100 | 100 | 100 | 100 | 100 | 100 | 100 | 100 | 100 | 100 | 100 | 100 |
| Amphora sp. | | | | + | + | 2.1 | | | | | | - | | | | | | | | | |
| Asteromphalus sp. | | | | | + | · + | + | + | | + | | | | | 200 0.5 | 150 0.5 | 2 | 35000 0.2 | | | |
| Biddulphia aurita | | | | | | | + | - | | | | | | | | | | | | | |
| Chaetoceros total | 105350 74 | 22000 55 | 61800 84 | 40600000 9 | 5 119740000 97 | 20070000 95 | 19400000 88 | 20420000 93 | 187600 79 | 21980 53 | 6300 60 | 4650 62 | 90800 72 | 9500 54 | 16000 43 | 6150 21 | 9500 75 | 15960000 85 | 678000 87 | 17400 11 | 720000 69 |
| Hyalochaete total | 88900 63 | • • • • • • • • • • • • • • • • • • • | Manage New Local and a star | 37400000 8 | 37 118360000 96 | 17910000 84 | 19100000 87 | 18880000 86 | 174300 73 | 16340 39 | 4500 43 | 50 0.7 | 4400 3.5 | 300 1.7 | 15750 42 | an an ann an an Arabana an Arab | | 11690000 62 | 126000 16 | 11400 7.0 | 18000 1.7 |
| Charles and a second transmission | | | 1 | 900000 - | 1 10010000 0 1 | | | | 9500 7 5 | | 1 | | | | | <u> </u> | 1 | | | | |
| Chaletoceros compressus | | | • | 220000 p. | 7 - | -1 | 200000 0 0 | | 5500 1.5 | | | | | | | | | | 10000 0.0 | | |
| C. constructus | 40950 05 | | | 1400000 3. | 1 T | 2640000 10 | 20000 0.9 | 4790000 00 | 700 0.3 | | | | | | 1-0-0 10 | | | 505000 00 | 10000 2.3 | 11100 0 0 | 10000 1 1 |
| C. <i>aeonis</i> | 49550 55 | | | 14400000 3 | 4 25190000 20 | 2040000 12 | 14300000 65 | 4720000 22 | 6300 2.6 | 10-00 00 | 1000 | | | | 15750 42 | | | 5950000 32 | 15000 1.9 | 11100 6.9 | 18000 1.7 |
| C. aecipiens | 33950 24 | T | | | 1040000 0.8 | 1440000 68 | 1600000 7.3 | 13180000 60 | 147000 62 | 13520 32 | 4200 40 | + | 1600 1.3 | 300 1.7 | | | | 980000 5.2 | 36000 4.6 | 300 0.2 | |
| C. aecipiens 1. singularis | | | | | 1 | 60000 0.3 | | | 2100 0.9 | 280 0.7 | | | | | | | | 140000 0.7 | | | |
| C. radicans | 2450 1.7 | | | 17000000 4 | 0 79350000 64 | 480000 2.3 | 1900000 8.6 | 70000 0.3 | 6300 2.6 | 2540 6.1 | | | 2800 2.2 | | | | | 3395000 18 | 57000 7.3 | | |
| C. seiracanthus | 3150 2.2 | | | 800000 1. | 9 + | 120000 0.6 | 600000 2.7 | 910000 4.2 | | | | | | | | | х. — х. | 1225000 6.5 | | | |
| C. socialis | | | • | | 350000 0.3 | | • | | | | | | | | | | | | | | |
| C. teres | | | | | 690000 0.6 | 150000 0.7 | + | | 5600 2.4 | | 300 2.9 | | | | | | | | | | |
| Hyalochaete miscs. | | | | 1400000 3. | 3 1730000 1.4 | 60000 0.3 | 500000 2.3 | | 2800 1.2 | | | 50 0.7 | | | | | | | | | |
| Phaseseres total | 16450 19 | 99000 55 | C1000 94 | 2200000 77 | F 1290000 1 1 | 9160000 10 | 200000 1 4 | 1=40000 7 1 | 19900 5 6 | FC 40 - 70 | 1000 10 | 4000 01 | 90100 00 | 0000 50 | 050 0 7 | 6150 91 | 0500 775 | 4970000 99 | FE9000 71 | 6000 9 7 | 700000 69 |
| Chastoseros atlanticus | 10450 12 | 22000 55 | 01000 04 | 200000 7. | 5 1360000 1.1 | 450000 2 1 | 50000 1.4 _ | 1540000 7.1 | 1000 0.0 | 5640 15 | 1600 17 | 4000 01 | 66400 69 | 9200 52 | 250 0.7 | 0100 21 | 9500 75 | 4270000 23 | 202000 71 | 000 0.0 | 702000 00 |
| Contentions & andax | | 300 020 | | 20000 0. | D | 40000 2.1 | | 550000 1.0 | 700 0.3 | 7 | т | T | 400 0.3 | 100 0.0 | • | | | 550000 1.9 | 202000 30 | 500 0.0 | 15000 1.4 |
| C. atlanticus 1. audus | | | | | | | | | | | | | | | | | | | 6000 0.8 | | |
| C. boreaiis | T | 400 7 0 | | | | | | + | | 280 0.7 | | | | | | | | 0.1=000 = 0 | | | |
| C. concavicornis | 2800 2.0 | 400 1.0 | | + | | | | 420000 1.9 | | 990 2.4 | | + | | 200 1.1 | | | <u> </u> | 945000 5.0 | | 300 0.2 | 21000 2.0 |
| C. convolutus | 13650 9.7 | 21300 54 | 61800 84 | 3000000 7. | 0 1380000 1.1 | 1710000 8.1 | 300000 1.4 | 770000 3.5 | 12600 5.3 | 4370 10 | 1800 17 | 4600 61 | 86000 68 | 8900 50 | 250 0.7 | 6150 21 | 9500 75 | 2975000 16 | 264000 34 | 4800 3.0 | 666000 64 |
| C. densus | | • | | | | | • + | | | | | | | | | | | | | | |
| Corethron hystrix | 2100 1.5 | 8900 22 | 2400 3.3 | 3 + . | + | + | + | + | 4 | + | 100 1.0 | 200 2.7 | 2800 2.2 | 2300 13 | 1950 5.2 | 4100 14 | 1500 12 | + | + | 5100 3.2 | 27000 2.6 |
| Coscinodiscus concinus | | | | + | | + | | | | | | | | | 100 0.3 | | | | | | |
| C. excentricus | | | | | + | + | + | + | | | | | | | | 150 0.5 | | | + | | • |
| C giggs | [| 1 | 1 | 1 | | | | | | 740 0 9 | | | | | | | | | | | |
| C. grgus | | 4 | | | | | 4 | | ~ | 140 0.3 | | 100 1 9 | | 900 1 1 | 50 0 1 | _ | | | | | |
| C. mu ginatus | | | | | | • | • | • | 700 0 0 | 140 0.3 | 100 1 0 | 100 1.3 | 400 0.9 | 200 1.1 | 50 0.1 | 150.0 # | =0.0.4 | | | | |
| Denticula sp | 7000 1 0 | 9200 21 | 9100 11 | 100000 0 | o + | - | 400000 1 8 | -1- | 700 0.3 | 4000 10 | 100 1.0 | 000.19 | 400 0.5 | | | 600 2 1 | 200 1 6 | 1050000 5 6 | 20000 2.0 | 9100 5 0 | 94000 9 7 |
| Denticula sp. | 7000 4.5 | 0200 21 | 0100 11 | 40000 0. | .9 1 | | 40000 1.0 | т | 5600 2.4 | 4230 10 | 100 1.0 | 950 13 | | | | 000 2.1 | 200 1.0 | 1050000 5.0 | 3000 3.9 | 8100 5-0 | 84000 8.1 |
| | | | | | | | - | | | | | | | | | | | | | | |
| Eucampia zoodiacus | 700 0.5 | 5 | | • | | | | | | | | • | | | | | | | | | |
| Navicula spp. | | | | | 690000 0.6 | | 200000 0.9 | 140000 0.6 | | р. — — — — — — — — — — — — — — — — — — — | | | | 400 2.3 | 50 0.1 | | | | 3000 0.4 | 300 0.2 | 3000 0.3 |
| Nitzschia closterium | 21000 15 | 5 | | 200000 0. | .5 ,+ | | | | | | | | | | | | | 35000 0.2 | | | |
| N. seriata | | 200 0.5 | 600 0.8 | 8 + | + | 810000 3.8 | + | 490000 2.3 | 16100 6.8 | 6490 15 | 900 8.6 | 550 7.3 | 4800 3.8 | 700 4.0 | 18700 50 | 6000 21 | 150 1.2 | 1155000 6.1 | 6000 0.8 | 128100 79 | 51000 4.9 |
| Rhizosolenia alata f. inermis | + | 200 0.5 | | + | 690000 0.6 | 210000 1.0 | 100000 0.5 | 70000 0.3 | 3500 1.5 | 280 0.7 | | • | 1200 1.0 | 100 0.6 | 150 0.4 | 100 0.3 | | 350000 1.9 | 36000 4.6 | 600 0.4 | 12000 1.2 |
| R. fragilissima | 1 | 1 | 1 | | 1040000 0 5 | | • | [| 1 | | | | | | | | | | | | |
| R. hebetata f hiemalis | | • | | | 1010000 010 | 4 | + | + | | | | | | | | 100 0 9 | | 4 | | | + |
| R. hebetata f comistina | 1000 2 | 5 100 0 9 | 600.05 | 8 4 | + | + | 200000 0 0 | 4 | 21000 0 0 | 2020 10 | 2000 29 | 850 11 | . 94000 10 | 4900 94 | | 11000 99 | 1950 0 9 | 105000 0 4 | 9 18000 9 9 | 000 0 6 | 141000 14 |
| Stophanopuris michonica | | | | | 4 | <u>+</u> | - | - | 21000 0.0 | - 0000 13 | 2300 20 | 11 060 | - 24000 13 | 7600 64 | | 11000 30 | 1400 3.0 | 10000 9.0 | 10000 2.0 | 300 0.0 | 141000 14 |
| Thalassionoma nitrechinidae | | | | + | + | • | | • | | - | | | | | | | | . | | | |
| | 1 | 1 5. 1 | 1 | <u> </u> | * | 1 | 1 | 1 | | | |] | | | | | | | | | |
| Thalassiosira decipiens | | | | + | + | 60000 0.3 | 100000 0,5 | + | | | | 100 1.3 | | | | | | | | | |
| T. nordenskiöldii | 8 6 5 6 | | | 1600000 3. | .7 1380000 1.1 | + | 1500000 6.8 | 630000 2.8 | • | | 100 1.0 | | | | | | | | | | |
| Thalassiothrix longissima | 700 0.: | 5 200 0.5 | 5 | + | + | 60000 0.3 | 100000 0.5 | 140000 0.6 | 3500 1.5 | 560 1.3 | 3 | 100 1.3 | 1600 1.3 | 300 1.7 | 50 0.1 | 250 0.9 | 50 0.4 | 140000 0.7 | 6000 0.8 | 1200 0.7 | + |