



Title	REPORT FROM THE “ OSHORO MARU ” ON OCEANOGRAPHIC AND BIOLOGICAL INVESTIGATIONS IN THE BERING SEA AND NORTHERN NORTH PACIFIC IN THE SUMMER OF 1955 : . Diatom Standing Crops and the Major Constituents of the Populations as Observed by Net Sampling
Author(s)	KAROHJI, Kohei
Citation	北海道大學水産學部研究彙報, 8(4), 243-252
Issue Date	1958-02
Doc URL	<a href="http://hdl.handle.net/2115/23010">http://hdl.handle.net/2115/23010</a>
Type	bulletin (article)
File Information	8(4)_P243-252.pdf



[Instructions for use](#)

# REPORT FROM THE "OSHO RO MARU" ON OCEANOGRAPHIC AND BIOLOGICAL INVESTIGATIONS IN THE BERING SEA AND NORTHERN NORTH PACIFIC IN THE SUMMER OF 1955

## IV. Diatom Standing Crops and the Major Constituents of the Populations as Observed by Net Sampling<sup>1)</sup>

Kohei KAROHJI<sup>2)</sup>

*Faculty of Fisheries, Hokkaido University*

### I. Introduction

On the 1955 cruise of the training ship "Oshoro Maru" to the Bering Sea sampling of plankton by vertical haul for upper 50 meter zone with a fine mesh net (0.112 mm in mesh aperture) was carried out covering almost all parts of the Bering Sea except north of 60 degrees north latitude, though stations of sampling were distributed rather sparsely (Fig. 1). Detailed descriptions on gear, methods, sampling stations and other




Fig. 1. Chart showing the location of the stations occupied by  
"Oshoro Maru" during June and July, 1955

records are given in the introductory notes of Motoda & Fujii (1956). The data of rough estimation on the displacement volume, weight and pigment content of the samples in these collections are presented in *Data Record of Oceanographic Observations and Exploratory Fishing*, No. 1, 1957 (pp. 88-92).

In the present studies the author intends to make more exact quantitative estimation of diatom standing crops by counting the cell number, and also to report the characteristic

1) おしよろ丸北洋調査報告 No.8 (昭和30年度)

2) 唐牛公平, 北海道大学水産学部浮游生物学教室, 函館市港町

species which occupy the main constituents of populations.

The author is gratefully indebted to Prof. S. Motoda, Asst. Prof. T. Kawamura and Mr. M. Anraku for their painstaking guidance throughout the present studies. He expresses his sincere gratitude to Mr. Y. Kawarada, Hakodate Marine Observatory, for his kindness not only in making available his unpublished data but in giving valuable criticism on the present paper. He also sincerely thanks Captain T. Fujii and his crew members as well as the research staff aboard the "Oshoro Maru" on that cruise for their work in sampling at sea. Cordial thanks are also due to Asst. Prof. H. Koto for his generosity in offering his unpublished data on the hydrographic analysis of this cruise.

## II. Standing Crops of Total Diatom Populations

As the net used was not equipped with a flow meter at its mouth, the exact quantity of water filtered by its cloth is unknown. All of the data on cell number presented in the present report are those which are multiplied by four on the basis of assumption that such type of the net would allow as little as 25 per cent of the water column to pass through the bolting cloth (Motoda *et al.*, 1957). Moreover, the increase of hauling distance due to the drift of the ship caused by wind may have added to the height of water column. This error was corrected on the basis of the actual records of revolutions of current meter which was attached near the net (see *Data Record*, No. 1, p. 89).




Fig. 2. Abundance of total diatom populations

The full data of the present studies are given in Table 2, among which abundance of total populations without regard to the species composition is illustrated in Fig. 2.

Throughout the whole areas investigated the quantity of diatoms vary from 100000 to 80000000 cells per 1 m<sup>3</sup> of water. Motoda & Kawarada (1955) gave an account of the quantity of diatoms on the basis of similar net samples in the western Aleutian waters taken on the cruise of May-June, 1953. Their data are not corrected either for errors due to filtration coefficient of the net nor for increase of hauling distance in rough weather. For comparison of size of populations between the 1953 and 1955 cruises the value 0.25 is tentatively adopted for filtration coefficient also to the data of 1953 cruise. Thus, the comparison of data from the two cruises shows that there were generally larger diatom populations in 1953 than in 1955.

	1953 cruise (Western Aleutian waters)	1955 cruise (Western Bering Sea: St. Os 1-Os 27)	1955 cruise (All stations)
Largest	494,160,000	79,600,000	79,600,000
Least	30,000	74,800	74,800
Mean	44,384,000	9,870,600	8,456,000

There are found rich populations of diatoms near Attu Island (St. Os 6 & Os 7), east of Bower's Bank (St. Os 20 & Os 23). Also diatoms are rich at St. Os 15, Os 27, Os 34 and at the south-easternmost station out of the Bering Sea (St. Os 37).

Locality	Cell number / m <sup>3</sup>
Near Attu Island (St. Os 6 & Os 7)	14,056,000-79,600,000
East of Bower's Bank (St. Os 20 & Os 23)	19,700,000-20,880,000
St. Os 15	55,088,000
St. Os 27	31,920,000
St. Os 34	6,956,000
St. Os 37	33,324,000

Previous report also shows diatom-rich water in the east of Bower's Bank, and considerably too, near Attu Island (Motoda & Kawarada, 1955).

Quantitative data on the diatoms in the Aleutian waters have been presented by Aikawa (1932, 1935). His data are based upon the samples collected with Kitahara's fine mesh net, and no correction for the error regarding filtration coefficient is made. He reported cell number per 1 m<sup>3</sup> of water as follows:

Year		Largest	Mean
1928	Over	3,000,000	935,000
1933	"	1,000,000	803,000

Although the comparison is very rough in exactness, the above data indicate far smaller populations than the present data. According to the dip-water samplings of Allen (1929,

1930), quantities of diatoms at the surface in Alaskan and Aleutian waters are as follows:

Locality and year	Cell number per m <sup>3</sup> in maximum catch among the stations
Alaskan waters (1923-24)	over 100,000,000
Mary I., Revillagigedo Channel, Alaska (1929)	40,000,000

Cupp (1937) recorded 28560000 cells per m<sup>3</sup> on average for five years (1927, 1929-32) at Scotch Cap Light, Unimak Island. In her summer collection the diatom cells were fairly abundant compared with the present data on St. Os 34 and Os 35 adjacent to Unimak Island.

### III. Species Composition of Diatom Populations and its Regional Characteristics

The diatom species which most predominantly appear in the present samples are as follows:

<i>Chaetoceros convolutus</i>	<i>Ch. concavicornis</i>
<i>Ch. debilis</i>	<i>Ch. didymus</i>
<i>Ch. seiracanthus</i>	<i>Ch. furcellatus</i>
<i>Ch. constrictus</i>	<i>Nitzschia seriata</i>
<i>Rhizosolenia hebetata</i> f. <i>semispina</i>	<i>Fragilaria</i> spp.
<i>Rh. heb.</i> f. <i>hiemalis</i>	<i>Thalassiothrix longissima</i>
<i>Denticula</i> sp.	

Table 1. Relative abundance of important diatoms in each region  
(In percent of cell number)

Main constituent	Region Station No.	1	2	3	4	5	6	7	8	9	10	11	12	13
		6-9, 14,15	3,4, 10,11	12,13	1	4,5	16-19	21,22 24-26	20,23	27	30-32	28,29	33,34	35-37
<i>Chaetoceros-Phaeoceros</i>		3	1	5	2	45	9	6	9	6	57	35	3	2
<i>Chaetoceros-Hyalochaete</i>		13	2	31	83	4	26	2	6	40	1	25	82	9
<i>Denticula</i> sp.		+	9	2	1	15	17	3	2	+	1	2	3	1
<i>Melosira sulcata</i>											9		+	
<i>Nitzschia seriata</i>		84	18	14		28	21	4	4	47	9	25	2	30
<i>Rhizosolenia hebetata</i> f. <i>semispina</i>			+				3	2	68	+	+	1	4	58
<i>Rhiz. heb.</i> f. <i>hiemalis</i>		+	1	1			4	18	+		+	8	+	
<i>Thalassiothrix longissima</i>		+	62	33	5	4	19	63	10	1	11	4		
Other Diatoms		+	7	14	9	4	2	2	1	6	12	+	+	

Fig. 3. Isotherms and isohalines at 25 meter depth (H. Koto)

A map of the Western Pacific Ocean region, showing the Philippines, Indonesia, and parts of the surrounding landmasses. The map is divided into 13 numbered regions, each with a different hatching pattern. The regions are numbered 1 through 13. The map includes latitude and longitude markings: 160E, 170E, 180, 170W, 160W on the top; and 50, 55, 60 on the right. A dashed line runs from the top left towards the bottom right, passing through regions 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, and 13. Region 1 is located in the central Philippines. Region 2 is to the west of Region 1. Region 3 is to the north of Region 1. Region 4 is to the southwest of Region 1. Region 5 is to the south of Region 1. Region 6 is to the southeast of Region 1. Region 7 is to the east of Region 1. Region 8 is to the south of Region 7. Region 9 is to the north of Region 7. Region 10 is to the east of Region 7. Region 11 is to the north of Region 10. Region 12 is to the south of Region 10. Region 13 is to the southeast of Region 12.

Fig. 4. Division of area according to the characteristics of diatom populations

Region 1 (St. Os 6–Os 9, Os 14 & Os 15): In the area covering St. Os 6, Os 7 and Os 15, *Nitzschia seriata* is abundant, occupying more than three-fourths of total

diatoms, and it is followed by *Chaetoceros-Hyalochaete*, mainly composed of *Ch. debilis*, *Ch. seiracanthus*, *Ch. decipiens*, *Ch. compressus* and *Ch. teres*, altogether which are less than 20% of total diatoms. It is indicated that the population generally includes neritic elements, except in the northern part (St. Os 8, Os 9 & Os 14) where such cold oceanic forms as *Chaetoceros convolutus*, *Thalassiothrix longissima* and *Denticula* sp. are present.

Region 2 (St. Os 2, Os 3, Os 10 & Os 11): In the western Bering Sea off the east coast of Kamchatka, cold-water species such as *Thalassiothrix longissima*, *Fragilaria* spp., *Denticula* sp. and *Nitzschia seriata* are widely distributed. The first species supplies more than a half of total diatoms. The dominance of these species may indicate the prevalence of cold water that flows southward along the coast of Kamchatka (Sverdrup *et al.*, 1942).

Region 3 (St. Os 12 & Os 13): Also *Thalassiothrix longissima* is dominant. *Hyalochaete* including *Chaetoceros debilis*, *Ch. compressus*, *Ch. decipiens*, *Ch. seiracanthus* and *Ch. furcellatus* is also numerous. This subgenus may indicate neritic influence in the southern offing of Cape Olyutorskii.

Region 4 (St. Os 1): In the offing to the southeast of Kamchatka, predominance of *Chaetoceros furcellatus* is observed, as abundant as about four-fifths of total diatoms. *Chaetoceros debilis*, *Ch. atlanticus*, *Fragilaria* spp. and *Thalassiothrix longissima* are found too, but very few in number. Predominance of such neritic forms has been also reported by previous workers in the area to the south of Kamchatka (Aikawa, 1938, 1940; Motoda & Kawarada, 1955; Marumo, 1956).

Region 5 (St. Os 4 & Os 5): Differing from the neighbouring stations, St. Os 4 and Os 5 are populated by somewhat peculiar composition, i. e., subgenus *Phaeoceros* (45%) containing *Chaetoceros convolutus*, *Ch. concavicornis* and *Ch. atlanticus*, and *Nitzschia seriata* (28%). In small quantities, *Thalassiothrix longissima*, *Corethron hystrix* and *Denticula* sp. are present. These diatom species reflect the cold oceanic nature of water in this area, agreeing with the results of previous investigations (Tsuruta & Chiba, 1954; Motoda & Kawarada, 1955). It is noted that the cold oceanic flora in this region is different in composition from that off east coast of Kamchatka (Region 2).

Region 6 (St. Os 16- Os 19): Along the northern side of Aleutian Islands, from Near Islands to Rat Islands, there are mixed associations, composed of *Hyalochaete* (26%), *Phaeoceros* (9%) and other oceanic forms. They are *Chaetoceros debilis*, *Ch. seiracanthus*, *Ch. atlanticus*, *Ch. convolutus*, *Denticula* sp. (17%), *Nitzschia seriata* (21%) and *Thalassiothrix longissima* (19%), but at St. Os 17 the last two species are considerably dominant. Predominance of *Hyalochaete* was reported by Motoda & Kawarada (1955) from the waters of western Aleutian Islands, but the present data show

that considerable portions of populations in the same area are occupied by oceanic forms.

Region 7 (St. Os 21, Os 22, Os 24 - Os 26): Region 7 covers both sides of Bower's Bank and extends toward the north. *Thalassiothrix longissima* is a leading species, more than 60%, and next comes *Rhizosolenia hebetata* f. *hiemalis*. But at St. Os 26 the latter species is exceptionally numerous. *Chaetoceros atlanticus*, *Nitzschia seriata*, *Denticula* sp. and *Ch. debilis* occur in certain number at St. Os 25 and Os 26. The above dominant forms reflect the cold oceanic nature of the water.

Region 8 (St. Os 20 & Os 23): In the south of the east side of Bower's Bank, rich diatom populations are found, as many as more than 21 millions. The populations are characterized by the dominance of *Rhizosolenia hebetata* f. *semispina*, *Thalassiothrix longissima*, *Nitzschia seriata* and several species of *Hyalochaete*. It is shown that similar diatom floras are to be found between here and the south of Alaska Peninsula (St. Os 37); e. g., *Rhizosolenia hebetata* f. *semispina* is found to constitute a major portion of populations in both regions.

Region 9 (St. Os 27): This station is located close to the edge of the shallow shelf of the eastern Bering Sea. Diatoms are remarkably rich, and represented by *Nitzschia seriata* (47%) and *Hyalochaete* (40%) including *Chaetoceros debilis*, *Ch. seiracanthus*, *Ch. subsecundus*, etc. *Chaetoceros concavicornis*, *Fragilaria* spp. and *Thalassiosira decipiens* are also present in small number. These associations are generally resemblant to those in Region 1. The transparency of water measures only 6 meters, and the temperature falls from 6.7 °C at 25 meter depth to 2.4°C at 50 meter depth.

Region 10 (St. Os 30-Os 32): In Bristol Bay large portions of diatom populations are occupied by *Chaetoceros convolutus* and *Ch. concavicornis*, accompanying with *Coscinodiscus* spp., *Nitzschia seriata* and *Melosira sulcata*. Contrary to the expectation, neritic associations are not observed. General poverty of diatoms and predominance of *Phaeoceros* indicate that the central portion of Bristol Bay is much affected by cold oceanic water.

Region 11 (St. Os 28 & Os 29): This region is located on the shallow shelf of the eastern Bering Sea between Regions 9 and 10. There are present mixed associations of *Hyalochaete* (25%), mainly composed of *Chaetoceros debilis*, and *Phaeoceros* (35%), represented by *Ch. convolutus*, together with *Nitzschia seriata* (25%). Accordingly, this region may be the transitional area between Regions 9 and 10.

Region 12 (St. Os 33 & Os 34): Northwest coast of Alaska Peninsula is occupied by *Chaetoceros didymus*, *Ch. constrictus*, *Ch. debilis*, *Ch. radicans*, etc. These diatoms reflect the presence of warm water in the eastern Bering Sea. In poor number, *Rhizosolenia hebetata* f. *semispina*, *Melosira sulcata* and *Corethron hystrix* are present. It is noted that such a warm-water species as *Chaetoceros didymus* is



dominant in the Bering Sea.

Region 13 (St. Os 35 - Os 37): On the Pacific side of the Alaska Peninsula, *Nitzschia seriata* and *Rhizosolenia hebetata* f. *semispina* are prevalent, especially rich in number and dominant in composition at St. Os 37. At St. Os 35 such *Hyalochaete* species as those found in Region 12 are numerous present, giving proof of neritic influence. Toward the south, *Chaetoceros concavicornis* and *Denticula* sp. occur in certain number, evidencing an indication of oceanic influence. *Rhizosolenia hebetata* f. *semispina* is also found which has been observed as a representative diatom on the east side of Bower's Bank.

There are found some inconstancies in the distributional feature of diatoms between the results of the present observations and the previous reports. In the present cruise the greater part of the Bering Sea is found to be dominated by such forms as *Thalassiothrix longissima*, *Nitzschia seriata*, *Rhizosolenia hebetata* f. *semispina* and *Rh. heb.* f. *hiemalis*. Such wide distribution of *Nitzschia seriata* has not been reported by previous worker (Aikawa, 1936, 1938, 1940). Also the dominant occurrence of *Chaetoceros didymus* which is known as a south temperate species, in the Bering sea is noted. The distribution of *Phaeo*-plankton is fairly localized in the present cruise. *Corethron hystris* and *Coscinodiscus* spp., the representatives of cold oceanic diatoms, are very rare compared with the previous records (Kanno, 1935; Yanagisawa, 1942; Aikawa, 1936, 1938, 1940; Marumo, 1956).

The surface diatom data by water dipping method on the same cruise as the present studies are provided by Mr. Y. Kawarada, Hakodate Marine Observatory. The diatom cell numbers are generally larger in his data than in the present. He obtained the maximum occurrence of diatoms amounting to over  $10^6$  cells per liter (i. e.,  $10^9$  cells per  $m^3$ ) at St. Os 27, and also found abundance of diatoms in Regions 1, 3, 12, 13 and 6. The present studies show the maximum occurrence in Region 1 amounting to over  $10^7$  per  $m^3$  and also abundant occurrence in Regions 8, 9 and 13. It must be kept in mind that his data are concerned with the surface dipping, while the present data are obtained by 0-50 meter hauls.

There are general agreements in distribution of dominant forms between the above vertical and the surface data. However, there are found remarkable differences of major constituents of diatoms between the two methods of sampling in the central portions of the Bering Sea (Regions 7 and 8). In Regions 7 and 8, *Thalassiothrix longissima*, *Rhizosolenia hebetata* f. *semispina* and *Rh. heb.* f. *hiemalis* are dominant in the upper 50 meters while *Denticula* sp. and *Hyalochaete* are exclusively dominant at the surface. There are also some inconstancies between the above two samplings in other several localities: *Hyalochaete* is exclusively dominant at the surface in Regions 6, 7 and 9, whilst mixed associations of *Hyalochaete*, *Thalassiothrix longissima*, *Nitzschia seriata*

and *Phaeoceros* are found in the vertical haul.

In conclusion, the area investigated may be divided on the basis of certain peculiarities of diatom populations as follows: neritic areas are (1) off the east coast of Kamchatka, (2) Alaskan coast, (3) central Aleutian waters, and (4) a west portion of shallow shelf of the eastern Bering Sea. However, (1) and (2) are distinguished from (3) and (4), by the different representatives of diatom species. Oceanic areas are (5) almost all parts of the western Bering Sea and (6) portions of the eastern Bering Sea. In this demarcation, there is a disagreement between the present data and the results of observations by Johnson (1953) and Vinogradov (1956), e. g., the area (6), contrary to their cases, is indicated as an oceanic area by the present data. It has been reported that the water highly influenced by neritic elements from Alaskan coast prevails in the eastern Bering Sea and this water flows northward along the Alaskan coast (Barnes & Thompson, 1938; Goodman *et al.*, 1942; Sverdrup *et al.*, 1942). However, the diatom distribution observed on the present cruise does not confirm the above statements in the point that the eastern Bering Sea is indicated to be oceanic by the dominant presence of cold oceanic populations.

#### IV. Summary

Observations of the diatom populations in the Bering Sea in the summer of 1955 were made. The density of diatoms as a whole varies between 100000 and 80000000 cells per 1 m<sup>3</sup> of water.

Cold oceanic populations represented by *Thalassiothrix longissima*, *Rhizosolenia hebetata* f. *hiemalis*, *Fragilaria* spp. and *Denticula* sp. are widely distributed in the western Bering Sea (Regions 2 & 7) (Fig. 4). In Region 1 *Nitzschia seriata* is dominant, accompanying with subgenus *Hyalochaete*, which represents somewhat neritic nature.

In the eastern Bering Sea and southwestern part of the western Bering Sea an oceanic group, *Phaeoceros*, is distributed in certain restricted locations (Regions 5 & 10).

Neritic populations represented by *Hyalochaete* occur to the southeast of Kamchatka, in western Bering Sea and to the east of the shallow shelf of the eastern Bering Sea (Regions 4, 12 & 9).

Mixed associations of subgenus *Hyalochaete* and other oceanic forms are found in Aleutian waters (St. Os 16 - Os 19), to the south of Cape Olyutorskii and in the eastern Bering sea, suggesting the mixing of oceanic and neritic waters (Regions 3, 6 & 11).

*Rhizosolenia hebetata* f. *semisfina* and *Nitzschia seriata* are abundantly distributed in the offshore area of Pacific side of Alaska Peninsula, and the former species is also found in the east of Bower's Bank (Region 8 & 13).

## Literature cited

- Aikawa, H. (1932). On the summer plankton in the waters of the west Aleutian Islands in 1928. *Bull. Jap. Soc. Sci. Fish.* 1 (2), 70-74. (in Japanese).
- (1935). On the quantitative analysis of the plankton associations in adjacent seas of Japan III. *Jour. Imp. Fish. Exp. Sta.* (6), 131-172. (in Japanese).
- (1936). The planktological properties of the principal sea areas surrounding Japan. *Bull. Jap. Soc. Sci. Fish.* 5 (1), 33-41. (in Japanese).
- (1938). On the quantitative analysis of the plankton associations in the adjacent seas of Japan V. *Jour. Imp. Fish. Exp. Sta.* (9), 67-86. (in Japanese).
- (1940). On the plankton associations in the Bering Sea and the Okhotsk Sea. *Kaiyō-Gyogyō* 5 (1), 20-31. (in Japanese).
- Allen, E. W. (1929). Surface catches of marine diatoms and dinoflagellates made by U. S. S. "Pioneer" in Alaskan waters in 1924. *Bull. Scripps Inst. Oceanogr. Tech. Ser.* 2 (2), 139-153.
- (1930). Quantitative studies of surface catches of marine diatoms and dinoflagellates taken in Alaskan waters by the International Fisheries Commission in the fall and winter of 1927-1928 and 1929. *Ibid.* 2 (10), 289-399.
- Anonymous (1957). *Data record of oceanographic observations and exploratory fishing* (1). 247p. Fac. Fish., Hokkaido Univ.
- Barnes, C. A. & Thompson, T. G. (1938). Physical and chemical investigations in Bering Sea and portions of the North Pacific Ocean. *Univ. Wash. Publ. Oceanogr.* 3 (2), 1-243.
- Cupp, E. E. (1937). Seasonal distribution and occurrence of marine diatoms and dinoflagellates at Scotch Cap, Alaska. *Bull. Scripps Inst. Oceanogr. Tech. Ser.* 4 (3), 71-100.
- Goodman, J. R., Lincoln, J. H., Thompson, T. G. & Zeusler, F. A. (1942). Physical and chemical investigations in Bering Sea, Bering Straits and Chukchi Sea during the summer of 1937 and 1938. *Univ. Wash. Publ. Oceanogr.* 3 (4), 105-169.
- Johnson, M. W. (1953). Studies on plankton of the Bering Sea and Chukchi Sea and adjacent areas. *Proc. Seventh Pac. Sci. Congr.* 4, 480-500.
- Kanno, R. (1935). The distribution of plankton during summer in the south Okhotsk Sea and on the coast of Kamchatka. *Jour. Fish., Hakodate Coll. Fish.* (38), 22-32. (in Japanese).
- Marumo, R. (1956). Diatom communities in Bering Sea and its neighbouring waters in the summer of 1954. *Oceanogr. Mag.* 8 (1), 69-73.
- Motoda, S. & Fujii, T. (1956). Report from the "Oshoro Maru" on oceanographic and biological investigations in the Bering Sea and northern North Pacific in the summer of 1955 I. *Bull. Fac. Fish., Hokkaido Univ.* 6 (4), 280-294.
- & Kawarada, Y. (1955). Diatom communities in western Aleutian waters on the basis of net samples collected in May-June, 1953. *Ibid.* 6 (3), 191-200.
- Sverdrup, H. U., Johnson, M. W. & Fleming, R. H. (1942). *The Oceans, their physics, chemistry and general biology*. 1087 p. Prentice-Hall Inc., N. Y.
- Tsuruta, A. & Chiba, T. (1954). On the distribution of plankton at the fishing ground of salmon in the North Pacific, 1952. *Jour. Shimonoseki Coll. Fish.* 3 (3), 39-45. (in Japanese).
- Vinogradov, M. E. (1956). Distribution of zooplankton in the western regions of Bering Sea. *Contr. Soviet Hydrobiol. Soc. Acad. Sci., U. S. S. R.* 7, 173-203. (in Russian).
- Yanagisawa, T. (1942). Plankton in the vicinity of Kurile Islands. *Jour. Oceanogr., Kobe. Mar. Obs.* 13 (3), 730-738. (in Japanese).

Table 2. Full Data of Diatom Observations (Cell number per cubic meter)

Station No.	1	2	3	4	5	6	7	8	9	10	11	12	13	14	15	16	17	18	19	20	21	22	23	24	25	26	27	27'	28	29	30	31	32	33	34	35	36	37		
Position	N 51-43 E 160-30	N 52-52 E 161-50	N 53-00 E 164-00	N 53-01 E 166-00	N 53-00 E 168-00	N 53-02 E 170-00	N 53-02 E 172-00	N 54-30 E 170-33	N 55-58 E 169-11	N 57-41 E 167-59	N 58-29 E 166-32	N 59-00 E 170-00	N 58-30 E 173-15	N 56-54 E 173-17	N 55-10 E 173-17	N 53-25 E 173-38	N 53-10 E 175-04	N 52-50 E 176-36	N 52-28 E 178-40	N 52-30 W 178-54	N 53-14 E 178-40	N 53-54 E 178-40	N 53-03 W 178-52	N 53-50 W 178-52	N 55-34 W 178-48	N 57-00 W 178-58	N 58-49 W 178-43	N 58-49 W 178-43	N 57-16 W 171-50	N 57-27 W 170-11	N 57-26 W 167-05	N 57-23 W 163-58	N 57-16 W 161-40	N 56-04 W 163-15	N 54-46 W 165-19	N 53-40 W 163-31	N 52-58 W 162-19	N 52-15 W 161-09		
Transparency of water (m)	15.00	17.20	23.28	14.08	16.17	14.16	10.11	21.24	15.55	15.33	15.00	10.03	13.22	9.00	10.45	10.18	10.20	9.40	12.51		18.30		10.27	15.08	17.30	12.25	5.14	5.14	15.36	12.08	9.20	9.15	10.00	12.18	12.24		21.19	20.18		
Species																																								
<i>Achnanthes longipes</i>														6000 (1.6)																										
<i>Asteromphalus</i> sp.									1200 (0.9)																															
<i>Biddulphia aurita</i>																																								
<i>Chaetoceros</i> total	366000 (85)	26000 (2.8)	332000 (19.92)	236000 (50.8)	188000 (42.5)	1520000 (10.8)	19360000 (24.3)	16600 (22.2)	31600 (23.4)	19600 (3.7)	14400 (2.2)	692000 (36.6)	170000 (33.8)	28000 (7.6)	8160000 (15.4)	452000 (31.8)	96000 (16.8)	788000 (36.8)	1360000 (38.2)	2640000 (13.4)	280000 (9.8)	332000 (25.4)	3720000 (18.02)	156000 (5.8)	80000 (6.1)		21000000 (40.2)	8288000 (72.2)	256000 (47.1)	528000 (69.4)	728000 (69)	3200 (1.8)	4000 (1.1)	143000 (40.8)	242400 (70.3)	6320000 (91.2)	48000 (34.8)	152000 (12.9)	3484000 (10.5)	
<i>Phaeoceros</i> total	10000 (2.3)	4000 (0.43)	312000 (18.72)	230000 (49.5)	158000 (34.7)	1040000 (7.4)	2720000 (3.5)	6600 (8.8)	10800 (8)	7600 (1.4)	4800 (0.7)	72000 (3.8)	26000 (5.1)	8000 (2.2)	120000 (0.2)	180000 (12.6)	58000 (10.2)	124000 (5.8)	360000 (10.1)	1880000 (9.5)	240000 (8.4)	212000 (16.2)	1960000 (9.4)	96000 (3.2)	80000 (6.1)		2840000 (5.4)	768000 (6.7)	132000 (24.3)	320000 (42)	728000 (69)	24200 (14.9)	143000 (40.8)	12400 (3.6)	184000 (2.6)	8000 (5.8)	120000 (10.2)	380000 (1.1)		
<i>Chaetoceros atlanticus</i>	10000 (2.3)	4000 (0.43)	12000 (0.72)	10000 (2.1)	10000 (2.3)	1040000 (7.4)	320000 (0.5)	1800 (2.42)	1200 (0.9)	4800 (0.7)	4800 (0.7)	12000 (0.6)	14000 (2.7)	48000 (0.09)	16000 (0.7)	180000 (12.6)	44000 (7.7)	16000 (0.8)	240000 (6.7)	1480000 (7.5)	240000 (8.4)	212000 (16.2)	560000 (2.7)	96000 (3.2)	80000 (6.1)		160000 (0.3)	48000 (0.5)	8000 (1.5)	320000 (42)	728000 (69)	24200 (14.9)	143000 (40.8)	12400 (3.6)	184000 (2.6)	8000 (5.8)	120000 (10.2)	380000 (1.1)		
<i>Ch. convolutus</i>			300000 (18)	200000 (43.1)	120000 (28)	1040000 (7.4)	2400000 (3)	4800 (6.4)	9600 (7.1)	2800 (0.5)	60000 (3.2)	12000 (2.4)	8000 (2.2)	72000 (0.14)	14000 (2.5)	108000 (5.0)	120000 (3.4)	400000 (2.0)																						
<i>Ch. concavicornis</i>				20000 (4.3)	18000 (4.2)	480000 (3.4)	16640000 (20.8)	10000 (13.4)	19600 (14.5)	12000 (2.3)	9600 (1.5)	620000 (32.8)	134000 (28.1)	20000 (5.4)	8040000 (15.2)	232000 (19.2)	38000 (6.6)	664000 (31.0)	1000000 (28.1)	760000 (3.9)	40000 (1.4)	120000 (9.1)	1800000 (8.6)	60000 (2.0)																
<i>Hyalochaete</i> total	356000 (82.7)	22000 (2.4)	20000 (1.2)	6000 (1.3)	30000 (7.5)	480000 (3.4)	16640000 (20.8)	10000 (13.4)	19600 (14.5)	12000 (2.3)	9600 (1.5)	620000 (32.8)	134000 (28.1)	20000 (5.4)	8040000 (15.2)	232000 (19.2)	38000 (6.6)	664000 (31.0)	1000000 (28.1)	760000 (3.9)	40000 (1.4)	120000 (9.1)	1800000 (8.6)	60000 (2.0)																
<i>Ch. decipiens</i>					8000 (1.8)		160000 (0.2)	1200 (2.4)	9600 (7.1)			180000 (9.5)	34000 (6.6)	14000 (3.8)	120000 (0.2)	30000 (2.1)	4000 (0.7)																							
<i>Ch. compressus</i>												60000 (3.2)	20000 (3.9)	6000 (1.6)	120000 (0.2)	8000 (0.5)																								
<i>Ch. didymus</i>																																								
<i>Ch. constrictus</i>																																								
<i>Ch. subsecundus</i>												80000 (4.2)	4000 (0.8)																											
<i>Ch. seiracanthus</i>							480000 (0.6)					40000 (2.2)			5280000 (9.9)	40000 (2.8)	10000 (1.8)	240000 (11)	240000 (6.7)	120000 (0.6)	28000 (0.9)																			
<i>Ch. debilis</i>	16000 (3.7)	22000 (2.4)		6000 (1.3)	3600 (0.8)		16000000 (20)	5200 (7)	9600 (7.1)			100000 (5.3)	50000 (9.8)		2400000 (4.5)	100000 (7)	20000 (3.5)	380000 (17.7)	520000 (14.6)	200000 (1.0)	40000 (1.4)	120000 (9.1)	1600000 (7.7)	32000 (1.1)																
<i>Ch. radicans</i>																																								
<i>Ch. furcellatus</i>	340000 (79)											20000 (1.05)	32000 (6.3)																											
<i>Ch. teres</i>					14000 (3)	320000 (2.3)		3000 (4)																																
<i>Hyalochaete</i> spp.			20000 (1.2)		4400 (1.5)	160000 (1.1)		8800 (6.5)				140000 (7.4)	4000 (0.8)		120000 (0.2)	42000 (3.6)	12000 (0.8)																							
<i>Corethron hystrix</i>	6000 (1.4)		8000 (0.5)	24000 (6.1)	10000 (2.5)			1200 (0.9)		4000 (0.8)			4000 (0.8)			24000 (1.7)																</								