



Title	PHOTOSYNTHESIS OF A NATURAL PHYTOPLANKTON POPULATION MAINLY COMPOSED OF A COLD DIATOM, THALASSIOSIRA HYALINA, IN HAKODATE HARBOR, MARCH 1962
Author(s)	MOTODA, Sigeru; KAWAMURA, Teruyoshi; SUZUKI, Tsuneyoshi; MINODA, Takashi
Citation	北海道大學水産學部研究彙報, 14(3), 127-130
Issue Date	1963-11
Doc URL	<a href="http://hdl.handle.net/2115/23184">http://hdl.handle.net/2115/23184</a>
Type	bulletin (article)
File Information	14(3)_P127-130.pdf



[Instructions for use](#)

PHOTOSYNTHESIS OF A NATURAL PHYTOPLANKTON POPULATION  
MAINLY COMPOSED OF A COLD DIATOM, *THALASSIOSIRA*  
*HYALINA*, IN HAKODATE HARBOR, MARCH 1962\*

Sigeru MOTODA, Teruyoshi KAWAMURA, Tsuneyoshi SUZUKI  
and Takashi MINODA

*Faculty of Fisheries, Hokkaido University*

**Observation**

In shallow water of Hakodate Harbor major part of phytoplankton population was composed of a cold water species of diatoms, *Thalassiosira hyalina*, in early spring 1962 (Table 1). Photosynthesis and chlorophyll *a* of this population were observed on board the "Oshoro Maru" lying at anchor in the harbor on March 7, 1962.

The water was very turbid, showing only 2 meters of Secchi disc reading. Underwater light intensities were observed by the third writer using a submarine photometer, Murayama Denki B type covered with neutral filter glass, at 10.00–11.00 a.m. on that day. The incident light was largely extinct in the water, becoming only 50 per cent of solar radiation at 0.3 m (Fig. 1). Solar radiation on the deck was 46 Klux at noon, 34 Klux at 1.00 p.m., 29 Klux at 2.00 p.m., 14 Klux at 3.00 p.m., 8.7 Klux at 4.00 p.m. and 0.5 Klux at 5.30 p.m. It was expected that the shallow water in the harbor would be well mixed because the water was subjected to the wind. Temperature gradient was very small, 5.0°C being recorded at the surface and 5.2°C at 5.5 m depth. Chlorophyll *a* as measured photometrically was 0.34 mg/m<sup>3</sup> at the surface, 4.0 mg/m<sup>3</sup> at 1.5 m depth and 0.37 mg/m<sup>3</sup> at 3.5 m

Table 1. Composition of phytoplankton population in Hakodate  
Harbor on March 7, 1962

Component	Cells per m <sup>3</sup>			Percentage		
	0 m	1.5 m	3.5 m	0 m	1.5 m	3.5 m
<i>Thalassiosira hyalina</i>	240×10 <sup>4</sup>	600×10 <sup>4</sup>	1840×10 <sup>4</sup>	80	70	85
<i>Chaetoceros debilis</i>	30×10 <sup>4</sup>	+	+	10		
<i>Fragilaria</i> sp.	30×10 <sup>4</sup>			10		
<i>Nitzschia seriata</i>	+	240×10 <sup>4</sup>			30	
<i>Donkinia</i> sp.			80×10 <sup>4</sup>			4
<i>Navicula</i> sp.			240×10 <sup>4</sup>			11

\* AEC Contract (04-3)-15 through Dr. Maxwell S. Doty

depth, indicating the uniformity of the water, although some differences in phytoplankton composition with depth were indicated (Table 1).

Water was sampled from the surface (100 % light), 1.5 m (20 % light), 3.5 m (5 % light) and 5.5 m (1 % light) depth with a Van Dorn Sampler. Each sample was transferred into two light bottles and one dark bottle, all 100 cc in capacity, to which 1 cc of  $^{14}\text{C}$  isotope ( $4.99 \times 10^4$  cpm) was added. Bottles were suspended




Fig. 1. Underwater light intensities and relative photosynthesis in Hakodate Harbor on March 7, 1962

in the sea from the ship at respective depths of sampling during about 6 hours from meridian passage of the sun (11.48 a.m.) to sunset (5.34 p.m.). Relative photosynthesis was highest at 1.5 m among samples from 0 m, 3.5 m and 5.5 m. At the surface it was as low as 70 % of that (12.3 mgC/m<sup>3</sup>/day) at 1.5 m. It became abruptly almost negligible between 1.5 m and 3.5 m (Fig. 1).

### Discussion

It will be permissible to say that the species, *Thalassiosira hyalina*, represented the population present in the water investigated at that time. The geographical distribution of this species has been studied by Smayda (1958). It is arctic form inhabiting mainly all the major polar currents. It has been recorded from the high latitudes of the Atlantic Ocean, and also from the Japan Sea at temperature below 10°C, excepting one record of occurrence obtained in Pusan in November when the sea temperature was 15.6°C. The species inhabiting high latitudes are thought to have characters of shade forms to some extent. Generally speaking, Ik is low in shade species inhabiting the water of low temperature. Winter population in Danish waters is considered to be shade forms showing 2.5–3.6 Klux (Steemann Nielsen and Hansen, 1956, 1961), and shade populations of Lake Fukami were reported as having 1–2 Klux of Ik (Ichimura, 1960 a, b). Although no tank experiment was carried out in the present observation, if average solar radiation on the deck (18 Klux) is taken for 1.00 p.m. (35 Klux)–3.30 p.m. (10 Klux), Ik of the population is estimated about 2.5 Klux, very similar value to those of shade forms obtained by the previous workers.

Carbon uptake per one mg of chlorophyll *a* is calculated 2.0 mgC at the surface, 2.4 mgC at 1.5 m and 0.6 mgC at 3.5 m. Steemann Nielsen and Hansen (1959) obtained a value as low as about 1 mgC/mgChl.-*a* of shade phytoplankton in Danish waters in winter. Generally speaking, the assimilation number is lower in winter population than in summer population, and lower in shade population in deep layer than in sun population near the surface. However, the assimilation number obtained by Ichimura and Saijo (1959) in the Kuroshio area is rather low (below 2 mgC/mgChl.-*a*).

Total daily production in the present case is estimated about 33.0 mgC/m<sup>2</sup>/day. This amount of production is much less compared with that on the Pacific coast of central Japan (300–500 mgC/m<sup>2</sup>/day), in the Kuroshio region (70–150 mgC/m<sup>2</sup>/day) (Ichimura and Saijo, 1959) and in the northwestern Pacific (200–300 mgC/m<sup>2</sup>/day) (Saijo and Ichimura, 1960) all measured in summer.

In conclusion, *Thalassiosira hyalina* which supplies the major part of the plankton population in Hakodate Harbor in early spring is probably of shade

form in photosynthetic activity; Ik is roughly estimated 2.5 Klux.

The writers are obliged to Captain T. Fujii of the "Oshoro Maru" who provided facilities for the experiments. Supply of filters and isotopes as well as the counting of radioactivity of the filters by Dr. Maxwell S. Doty, Botany Department, University of Hawaii, are greatly appreciated.

#### Literature cited

- Ichimura, S. & Saijo, Y. (1959). Chlorophyll content and primary production of the Kuroshio off the southern midcoast of Japan. *Bot. Mag. (Tokyo)*, **72** (851), 193-202.
- Ichimura, S. (1960 a). Diurnal fluctuation of chlorophyll content in lake water. *Ibid.* **73** (864), 217-224.
- (1960 b). Photosynthetic pattern of natural phytoplankton relating to light intensity. *Ibid.* **73** (869-870), 458-467.
- The Faculty of Fisheries, Hokkaido University (1961). Data on C-14 primary productivity measurements on the "Oshoro Maru" cruises to and from Manila in November-December 1959 and to and from Seattle in June-August 1960. *Data Rec. Oceanogr. Obs. and Expl. Fish., Fac. Fish., Hokkaido Univ.* (5), 33-39; 142-165.
- Saijo, Y. & Ichimura, S. (1960). Primary production in the northwestern Pacific Ocean. *Journ. Oceanogr. Soc. Japan*, **16** (3), 139-145.
- Smayda, T. T. (1958). Biogeographical studies of marine phytoplankton. *Oikos*, **9** (2), 158-191.
- Steemann Nielsen, E. & Hansen, Vagn Kr. (1959). Light adaptation in marine phytoplankton populations and its interrelation with temperature. *Physiol. Plant.* **12**, 353-370.
- & ——— (1961). Influence of surface illumination on plankton photosynthesis in Danish waters (50°N) throughout the year. *Ibid.* **14**, 595-613.