



Title	INTRACEREBROVENTRICULAR ADMINISTRATION OF CATECHOLAMINES INHIBITS THE PANCREATIC SECRETORY RESPONSES INDUCED BY INTRACEREBROVENTRICULAR ADMINISTRATION OF THYROTROPIN RELEASING HORMONE IN ANAESTHETIZED RATS
Author(s)	EGUCHI, Junichi
Citation	Japanese Journal of Veterinary Research, 33(1-2), 68-68
Issue Date	1985-04-30
Doc URL	<a href="http://hdl.handle.net/2115/2323">http://hdl.handle.net/2115/2323</a>
Type	bulletin (article)
File Information	KJ00002374295.pdf



[Instructions for use](#)

INTRACEREBROVENTRICULAR ADMINISTRATION OF CATECHOLAMINES  
INHIBITS THE PANCREATIC SECRETORY RESPONSES INDUCED BY  
INTRACEREBROVENTRICULAR ADMINISTRATION OF THYROTROPIN  
RELEASING HORMONE IN ANAESTHETIZED RATS.

Junichi EGUCHI

*Department of Veterinary Physiology,  
Faculty of Veterinary Medicine,  
Hokkaido University, Sapporo 060, Japan*

Pancreatic secretory responses such as protein output and juice flow were induced by intracerebroventricular (i. c. v.) injection of TRH (50 pmol / 100g B. W.) in anaesthetized rats. Preceding the TRH-induced responses, catecholamines (adrenaline, noradrenaline and dopamine) were injected into the cerebral ventricle. The TRH-induced pancreatic responses were inhibited in a dose-dependent manner by i. c. v. pre-injections of adrenaline ranging from 0.1 to 2.5 nmol / 100g B. W.

I. c. v. injection of 0.1 nmol / 100g B. W. noradrenaline had little, if any, effect on the TRH-induced pancreatic secretion. I. c. v. injection of 0.5 or 1 nmol / 100g B. W. noradrenaline tended to reduce the TRH-induced pancreatic responses, which, however, were not significantly smaller than the respective responses to TRH alone. I. c. v. injection of 2.5 nmol / 100g B. W. noradrenaline significantly inhibited the TRH-induced pancreatic secretory responses. The heart rate was unchanged when a maximal dose (2.5 nmol / 100g B. W) of adrenaline or noradrenaline and 50 pmol / 100g B. W. of TRH were administered intracerebroventricularly.

The TRH-induced pancreatic responses were inhibited in a dose-dependent manner by i. c. v. pre-injections of dopamine ranging from 5 to 50 nmol / 100g B. W. The intravenous (i. v.) administration of a high dose (500 nmol / 100g B. W.) of dopamine significantly inhibited TRH-induced pancreatic secretion, but the i. v. administration of 50 nmol / 100g B. W. of dopamine did not cause any effect on the secretory responses to the electrical stimulation of the vagus nerve.

These results suggest that i. c. v. pre-injection of catecholamine inhibits the TRH-induced pancreatic secretion in the central nervous system. The inhibitory effects of adrenaline and noradrenaline were about 20 times as potent as those of dopamine.