



Title	DIFFERENCE OF Ca^{2+} and Na^+ DEPENDENCY AMONG SECRETORY RESPONSES TO VARIOUS CCK_8 CONCENTRATIONS IN RAT EXOCRINE PANCREAS
Author(s)	KAWAHARA, Junichi
Citation	Japanese Journal of Veterinary Research, 33(1-2), 74-74
Issue Date	1985-04-30
Doc URL	http://hdl.handle.net/2115/2329
Type	bulletin (article)
File Information	KJ00002374301.pdf



[Instructions for use](#)

DIFFERENCE OF Ca^{2+} and Na^+ DEPENDENCY AMONG SECRETORY
RESPONSES TO VARIOUS CCK_8 CONCENTRATIONS IN RAT EXOCRINE
PANCREAS.

Junichi KAWAHARA

*Department of Veterinary Physiology,
Faculty of Veterinary Medicine,
Hokkaido University, Sapporo 060, Japan.*

The roles of extracellular Na^+ ($[\text{Na}^+]_o$) and Ca^{2+} ($[\text{Ca}^{2+}]_o$) were examined in pancreatic responses (protein output and juice flow) induced by CCK_8 at physiological (5 pM), maximal (50 pM) and a supramaximal (1 nM) concentrations in the isolated perfused rat pancreas.

The pancreatic responses induced by 5 pM CCK_8 were completely abolished by removing either $[\text{Na}^+]_o$ or $[\text{Ca}^{2+}]_o$. The pancreatic responses induced by 50 pM CCK_8 were decreased, but still remained by removing either $[\text{Na}^+]_o$ or $[\text{Ca}^{2+}]_o$. The pancreatic responses induced by 1 nM CCK_8 were smaller than those induced by 50 pM CCK_8 . The responses were not completely abolished by removing $[\text{Ca}^{2+}]_o$, and were not affected by removing $[\text{Na}^+]_o$. The responses, however, were completely abolished by removing both $[\text{Na}^+]_o$ and $[\text{Ca}^{2+}]_o$.

Based on these and other results, the following conclusions were made :

- (1) The pancreatic secretory responses induced by CCK_8 at the physiological concentration are initiated by $[\text{Na}^+]_o$ -dependent Ca^{2+} influx (e. g. Na^+ and Ca^{2+} cotransport).
- (2) The pancreatic secretory responses induced by CCK_8 at the maximal concentration consist of two components : $[\text{Na}^+]_o$ -independent Ca^{2+} influx and $[\text{Na}^+]_o$ -dependent Ca^{2+} release from intracellular Ca stores.
- (3) The pancreatic secretory responses induced by CCK_8 at the supramaximal concentration also contain the above two components. The response may coincide with $\text{Na}^+/\text{Ca}^{2+}$ countertransport, and diminution of the responses may be due to an extreme increase in $[\text{Ca}^{2+}]_i$.