



Title	ISOLATION OF NEW SPECIES OF CORYNEBACTERIA FROM URINARY TRACT OF PIGS AND PRODUCTION OF MONOCLONAL ANTIBODIES AGAINST BOVINE CORYNEBACTERTAL PILI
Author(s)	KUDO, Yukiko
Citation	Japanese Journal of Veterinary Research, 33(1-2), 82-82
Issue Date	1985-04-30
Doc URL	http://hdl.handle.net/2115/2337
Type	bulletin (article)
File Information	KJ00002374309.pdf



[Instructions for use](#)

ISOLATION OF NEW SPECIES OF CORYNEBACTERIA FROM URINARY TRACT
OF PIGS AND PRODUCTION OF MONOCLONAL ANTIBODIES AGAINST BOVINE
CORYNEBACTERIAL PILI

Yukiko KUDO

*Department of Hygiene and Microbiology
Faculty of Veterinary Medicine
Hokkaido University, Sapporo 060, Japan*

Urease-positive corynebacteria were isolated from the prepuce of 50 out of 77 (64.9%) hogs and from the urinary bladder and urethra of 3 out of 10 (30%) sows. Ninety-three isolates were divided into 4 groups, A, B, C and ungrouped, based on their morphological, cultural and biochemical characteristics. Group A strains were the highest in frequency (53.2% of hogs), followed by strains of group B (5.2%), group C (2.6%) and ungrouped (10.4%). Corynebacteria of group A and ungrouped were also isolated from sows. A numerical taxonomic study of the 94 features of the strains showed that there was at least 1 phenon, which consisted of group A strains. The remaining strains did not form distinct phenons. Phenon 1 (group A) seemed to be a new species of genus *Corynebacterium*, however, further study is necessary to define the taxonomic position.

Four and one monoclonal antibodies to the pili of *C. renale* 115 and *C. pilosum* 92 were produced. These monoclonal antibodies bound to the purified pili of homologous strain in enzyme-linked immunosorbent assay (ELISA) and agglutinated piliated bacteria (P^+), but not non-piliated bacteria (P^-), of each homologous strain. Anti-*C. pilosum* 92 pili monoclonal antibody in the form of IgG and Fab inhibited the adherence of *C. pilosum* 92 P^+ bacteria to the isolated bovine vulva epithelial cells. Of the monoclonal antibodies against *C. renale* 115 pili, one inhibited the adherence of *C. renale* 115 P^+ bacteria to the bovine vulva epithelial cells, while the other did not, indicating the possible presence of different antigenic determinants on the pilus.