



Title	Stability Constant of Copper with Fulvic Acid
Author(s)	MATSUNAGA, Katsuhiko; NEGISHI, Masami; FUKASE, Sigeru
Citation	北海道大學水産學部研究彙報, 30(4), 301-305
Issue Date	1979-11
Doc URL	http://hdl.handle.net/2115/23699
Type	bulletin (article)
File Information	30(4)_P301-305.pdf



[Instructions for use](#)

Stability Constant of Copper with Fulvic Acid

Katsuhiko MATSUNAGA*, Masami NEGISHI* and Sigeru FUKASE*

Abstract

Fulvic acid, which is extracted from bottom sediments in both the Kunebetsu River and Shukunohe River, was titrated by the addition of copper ion and the uncomplexed copper ion was measured with a copper ion-selective electrode. From equation (4), the stability constant of copper with fulvic acid was calculated and then the value of $\log \beta=5.3$ was obtained.

Introduction

Though the chemical species of heavy metals which were utilized or taken up by phytoplankton are not clearly known, we found that a fulvic acid-Fe complex in Ohnuma Lake was utilized by phytoplankton¹⁾. Thus, it is very important to study the chemical species of heavy metals in natural waters.

Zirino and Yamamoto²⁾, and Long and Angino³⁾ estimated thermodynamically the chemical species of several metals by using several inorganic anions as ligand. However, natural waters contain not only the inorganic ligands but also organic ligands. Thus, it is insufficient to take into account only the inorganic anions as ligands. In recent, Mantoura et al.⁴⁾ calculated the chemical species of heavy metal ions on the assumption of the presence of organic ligands such as humic material. However, the stability constant of copper with humic material which they used for the calculation was higher than already reported values.^{5,6)}

The purpose of this paper is to measure the stability constant of copper with fulvic acid, which is the main humic materials in river.^{7,8)} By measuring the stability constant, the chemical species of copper in rivers or lakes in a suburb of Hakodate would be thermodynamically calculated and then chemical species which are utilized by phytoplankton would be known.

Experiment

Reagents

1 M NaClO₄: 12.2 g of NaClO₄ was dissolved in 100 ml of distilled water. Cu²⁺ standard solution (10⁻¹, 10⁻², 10⁻³, 10⁻⁴ M): 2.49 g of CuSO₄·5H₂O was dissolved in 100 ml of distilled water and 10⁻², 10⁻³, and 10⁻⁴ M standard solutions were prepared by diluting the 10⁻¹ M Cu²⁺ solution.

All chemical reagents used were of analytical grade.

* Laboratory of Marine Chemistry, Faculty of Fisheries, Hokkaido University
(北海道大学水産学部海洋化学講座)

Apparatus

The potentials were measured with a Toa Denpa HM-15A digital pH meter by using an Orion copper ion-selective electrode (ISE).
pH was measured with a Hitachi-Horiba F-5 pH meter.

Sample used

Bottom sediments both in the Kunebetsu River in October 1977 and in the Shukunohe River which flows in Ohnuma Lake in June 1978 were taken.

Extraction of fulvic acid from the bottom sediments: 200 ml of 1 N NaOH was added to 50 g of the sediment and the mixed solution was shaken for 2 hr. The undissolved solid was separated by a centrifuge and then the upper layer was adjusted to pH to 1 with the addition of HCl. The solution was again centrifugalized to remove a precipitation which was humic acid. The upper layer was filtered through an HA type Millipore filter with a 0.45 μm pore size. The filtrate obtained was used as a sample of fulvic acid.

Procedures

A solution, which was adjusted ion intensity to 0.6 with NaClO_4 and the concentration of fulvic acid to about 10^{-3} M, was adjusted pH to 6.0 with NaOH at 25°C. After an aliquot of a standard Cu^{2+} solution was added to the solution, the solution was mixed with a stirrer. The uncomplexed concentration of Cu^{2+} was measured by ISE.

Calculations

The response of ISE is given by

$$E = E_0 + \frac{RT}{2F} \ln[\text{Cu}] = E_0 + \rho \log [\text{Cu}] \quad (1)$$

where $[\text{Cu}]$ is the concentration of the hydrated ion Cu^{2+} in the solution, E and E_0 are the potentials of Cu^{2+} and the standard potential of Cu^{2+} respectively. R , T , F and ρ are gas constant, absolute temperature, Farady constant and $2.3 RT/2F$ respectively.

The degree of complexation of Cu^{2+} , α , is defined as

$$\alpha = [\text{Cu}]_T / [\text{Cu}] \quad (2)$$

where $[\text{Cu}]_T$ and $[\text{Cu}]$ are the total concentration of Cu^{2+} and the hydrated Cu^{2+} respectively.

By combining equations (1) and (2)

$$\alpha = [\text{Cu}]_T \times 10^{(E_0 - E)/\rho} \quad (3)$$

It is assumed that fulvic acid is approximately dissociated at pH 6.0 because of the dissociation constant of $pK_a=4.2$.⁹⁾ Thus, the calculation of the stability constant of Cu^{2+} with fulvic acid was slightly modified by the Buffle method.⁶⁾

$$Y = \left(\frac{[\text{L}]_T}{[\text{Cu}]_T} \right) \left(\frac{\alpha}{\alpha - 1} \right) = M_W + \left(\frac{M_W}{\beta} \right) \left(\frac{\alpha}{[\text{Cu}]_T} \right) \quad (4)$$

where $[L]_T$, M_w and β are the total concentration of fulvic acid, the molecular weight of fulvic acid and the stability constant of Cu^{2+} with fulvic acid respectively.

A plot of the left hand term of equation (4) vs. $\alpha/[\text{Cu}]_T$ should give a straight line and from the slope and the intercept on the ordinate, M_w and β can be evaluated.

Results and discussion

Plots of E vs. $[\text{Cu}]_T$ for the titration of fulvic acid with Cu^{2+} at pH 6.0 in the Kunebetsu River are shown in Fig. 1. Solid and dotted lines represent the plots in the absence of fulvic acid and in the presence of fulvic acid respectively. The difference of both lines shows the formation of a fulvic acid-Cu complex. From equation (4), plots of Y vs. $\alpha/[\text{Cu}]_T$ are shown in Fig. 2. The straight line was calculated by the method of least squares and the equation, $Y=515+2.6 \times 10^{-3} \times \alpha/[\text{Cu}]_T$ was obtained. From the slope and intercept, $M_w=515$ and $\log \beta=5.3$ were obtained. From the results, Cu^{2+} forms 1:1 complex with fulvic acid.

The stability constant of Cu^{2+} with fulvic acid which was extracted from bottom sediment in the Shukunohe River was also the same results as that in the Kunebetsu River. The value obtained $\log \beta=5.3$ is assumed to be reasonable one because already reported values are $\log \beta=5.67-5.95^5)$, $4.8-5.0.^6)$ By using the value, chemical species of copper in river or lake waters in a suburb of Hakodate were thermodynamically calculated.¹⁰⁾




Fig. 1. Variation of potential titrated with copper.
Solid line: in absence of fulvic acid; ●: in presence of fulvic acid




Fig. 2. Relation between Y and $1/[\text{Cu}]$ according to eq. (4).

Acknowledgements

We thank Profs. H. Yoshida and M. Taga, Faculty of Science, Hokkaido University, for lending us a copper ion-selective electrode.

References

- 1) Matsunaga, K., K. Igarashi and S. Fukase (1979). Study on a organic-Fe complex in lake water: paper presented at the annual meeting of the Oceanographical Society of Japan, Tokyo, April, p. 198.
- 2) Zirino, A. and S. Yamamoto (1972). A pH-dependent model from the chemical speciation of copper, zinc, cadmium and lead in seawater: *Limnol. Oceanogr.*, **17**, 661-671.
- 3) Long, D.T. and E.E. Angino (1977). Chemical speciation of Cd, Cu, Pb and Zn in mixed fresh water, seawater and brine solutions: *Geochim. Cosmochim. Acta*, **41**, 1183-1191.
- 4) Mantoura, R.F.C., A. Dickson and J.P. Riley (1978). The complexation of metals with humic materials in natural waters: *Est. Coastal Mar. Sci.*, **6**, 387-408.
- 5) Shuman, M.S. and J.L. Cromer (1979). Copper association with aquatic fulvic and humic acids, estimation of conditional formation constants with a titrimetric anodic stripping voltammetry procedure: *Environ. Sci. Tech.*, **13**, 543-545.
- 6) Buffle, J., F.L. Greter and W. Haerdi (1977). Measurement of complexation properties of humic and fulvic acids in natural waters with lead and copper ion-selective electrode: *Anal. Chem.* **49**, 216-222.
- 7) Kemp, A.L.W. and H.K.T. Wong (1974). Molecular weight distribution of humic substances from lakes Ontario and Erie sediments: *Chem. Geol.*, **14**, 15-22.

- 8) Reuter, J.H. and E.M. Perdue (1977). Importance of heavy metal-organic matter interactions in natural waters: *Geochim. Cosmochim. Acta*, **41**, 325-334.
- 9) Wilson, D.E. (1977). Effects of polymeric charge variations on the proton-metal ion equilibrium of humic materials: *Limnol. Oceanogr.*, **22**, 281-288.
- 10) Matsunaga, K., M. Negishi, S. Fukase and K. Hasebe (1979). Chemical species of copper in river and lake waters: submitted to *Bull. Chem. Soc. Japan*.