



Title	Mitosis in <i>Archaeomysis kokuboi</i> (Mysidae, Crustacea)
Author(s)	YABU, Hiroshi; KAMIHIRA, Yukiyoshi
Citation	北海道大學水産學部研究彙報, 32(2), 136-137
Issue Date	1981-06
Doc URL	http://hdl.handle.net/2115/23749
Type	bulletin (article)
File Information	32(2)_P136-137.pdf




[Instructions for use](#)

Mitosis in *Archaeomysis kokuboi* (Mysidae, Crustacea)

Hiroshi YABU* and Yukiyoishi KAMIHIRA**

Ecology of a mysid, *Archaeomysis kokuboi*¹⁾ occurring on the shore around Hakodate, Hokkaido, have been studied by the junior author²⁾. In our recent cytological study on this species, we succeeded to count approximate number of chromosomes.

The materials were collected on the intertidal zone at the open sandy shore of Nesaki, Hakodate in January, 1981. They were fixed immediately with acetic alcohol (1:3) after catch by a core sampler (basal area 30 cm², with 10 cm height) and separated from the sand through mesh of 0.5 mm square opening. The staining was done with the simple squash method recommended by Wittmann³⁾. The individuals studied were 11-15 mm in total length in both sexes, and most of them were immature.



Figs. 1-7. Nuclear divisions in *Archaeomysis kokuboi* Ii
1-3. Metaphase in male somatic cells 4. Metaphase in female somatic cell
5. Early anaphase in male somatic cell 6. Mid-anaphase in male somatic cell
7. Metaphase in spermatogonium
Magnification: 1-4, $\times 1,450$; 5, $\times 1,800$; 6, $\times 1,350$; 7, $\times 1,200$.

* Laboratory of Marine Botany, Faculty of Fisheries, Hokkaido University.
(北海道大学水産学部水産植物学講座)

** Research fellow: Laboratory of Planktology, Faculty of Fisheries, Hokkaido University.
(北海道大学水産学部浮遊生物学講座)

The dividing nuclei were found in thoracic appendages and various internal organs. The chromosomes in both sexes showed to have $2n \approx 40$ (Figs. 1-3 & 7) including one or two slightly longer chromosomes with occasionally formed chromosome bridge or trail at anaphase (Figs. 5 & 6). All metaphase chromosomes in somatic cells were metacentric, assuming small dumbbell- or V-shape. At the polar view of metaphase, the chromosome alignment forms a ring (Fig. 3). The diploid chromosomes in spermatogonia are much larger than in somatic cells and are discernible to differ in size (Fig. 7). No centrosome was detected in any metaphase or anaphase.

References

- 1) Ii, N. (1964). Fauna Japonica Mysidae (Crustacea). *Biogeograph. Soc. Japan. National Science Museum*, Tokyo, 1-610.
- 2) Kamihira, Y. (1979). Ecological studies of macrofauna on a sandy beach of Hakodate, Japan. II. On the distribution of peracarids and the factors influencing their distribution. *Bull. Fac. Fish. Hokkaido Univ.* **30**(2), 133-143.
- 3) Wittmann, W. (1965). Aceto-iron-haematoxylin-chloral hydrate for chromosome staining. *Stain Tech.* **40**, 161-164.