



Title	IMMUNOCYTOCHEMICAL AND ULTRASTRUCTURAL STUDIES ON THE ENDOCRINE CELLS IN CHICK THYMUS
Author(s)	KAWAI, Koji
Citation	Japanese Journal of Veterinary Research, 41(1), 26-26
Issue Date	1993-05-27
Doc URL	<a href="http://hdl.handle.net/2115/2418">http://hdl.handle.net/2115/2418</a>
Type	bulletin (article)
File Information	KJ00002377631.pdf



[Instructions for use](#)

IMMUNOCYTOCHEMICAL AND ULTRASTRUCTURAL STUDIES  
ON THE ENDOCRINE CELLS IN CHICK THYMUS

Koji KAWAI

*Department of Veterinary Anatomy,  
Faculty of Veterinary Medicine,  
Hokkaido University, Sapporo 060, Japan*

Endocrine cells in chick thymus were studied immunocytochemically and ultrastructurally.

Serotonin-, dopamine-, gastrin/CCK-, CCK8-, neurotensin-, peptide YY-, glucagon-, somatostatin-, insulin-, calcitonin-, growth hormone-, and SP-1/chromogranin-immunoreactive cells were detected in the chick thymus by the peroxidase-antiperoxidase technique. These immunoreactive cells, except for the calcitonin cells, were restrictively observed in the medulla and particularly predominated in the juxtacortical medulla. They were polymorphous and were often ovoid in shape. Calcitonin-immunoreactive cells were observed extensively in the deep cortex and medulla, and the cells in the cortex were reticular, while those in the medulla were oval in shape.

Ultrastructurally, the endocrine cells were identified by the presence of numerous secretory granules and by possessing an electron-lucent nucleus. These cells were classified into eleven types by their granule sizes and internal structures. Immunoelectron microscopically, gastrin/CCK-, somatostatin-, serotonin-, calcitonin-, glucagon-, and neurotensin-immunoreactive cells were demonstrated with the protein A-colloidal gold technique.

The developmental profile of serotonin-immunoreactive cells was estimated to examine the possible function of thymic endocrine cells on the maturation of intrathymic lymphocytes. The relative number of serotonin-immunoreactive cells significantly increased from hatching day to one week of age, whereas it then decreased from young chicks to adults. These results suggest that the intrathymic endocrine cells may share some associations with the functional maturation of thymocytes on the neonatal establishment of cell-mediated immunity and humoral immunity.