



Title	ANOMALOUS SALT EFFECT ON THE RATE OF HYDROLYSIS OF CHLOROFORM. : An Additional Remark
Author(s)	TANABE, Kozo
Citation	JOURNAL OF THE RESEARCH INSTITUTE FOR CATALYSIS HOKKAIDO UNIVERSITY, 8(1), 12-13
Issue Date	1960-07
Doc URL	http://hdl.handle.net/2115/24712
Type	bulletin (article)
File Information	8(1)_P12-13.pdf



[Instructions for use](#)

ANOMALOUS SALT EFFECT ON THE RATE OF HYDROLYSIS OF CHLOROFORM.

An Additional Remark



By

Kozo TANABE^{*}

(Received February 10, 1960)

In the previous paper we reported that salt effect on the rate of hydrolysis of chloroform is anomalously large; namely that, in aqueous solution around pH 4, 0.01 N sodium sulfate decreases the rate about six-fold and 0.2 N sodium sulfate does sixteen-fold at least¹⁾. The pronounced salt effect has been accounted for by assuming an salt effect on elementary step (2) particularly when HA is H₂O in our reported mechanism^{1,2)}.

More detailed discussion is given here on the nature of the salt effect. The mechanism of hydrolysis of chloroform proposed by HORIUTI *et al.*²⁾ is quoted briefly.



where HA and B denote Brönsted acid and base respectively and Cl₂C[×]ClH an isomer of chloroform. In acidic region, step (2) controls the hydrolysis, step (1) being in equilibrium. We shall now show that the pronounced salt effect leads necessarily to that on the activity of the critical complex of step (2). The observed rate of hydrolysis is, according to the mechanism, identical with the forward rate \bar{v} of rate-determining step (2), which is expressed as $\bar{v} = k_2 a^{\text{CCl}_3^-} a^{\text{H}^+}$, where $a^{\text{CCl}_3^-}$ or a^{H^+} is the absolute activity of trichloromethyl anion CCl₃⁻ or hydrogen ion H⁺. Since step (1) is in equilibrium, $a^{\text{CCl}_3^-}$ is fixed at the ratio $a^{\text{CHCl}_3}/a^{\text{H}^+}$ of the absolute activity of chloroform a^{CHCl_3} to a^{H^+} . Therefore, $a^{\text{CCl}_3^-} a^{\text{H}^+}$ is kept constant independent of neutral salt concentration

^{*} Research Institute for Catalysis, Hokkaido University, Sapporo.

Anomalous Salt Effect on the Rate of Hydrolysis of Chloroform

at constant pH, since a^{CHCl_3} is fixed at that of practically pure chloroform phase present under the experimental condition. The neutral salt effect comes hence into play only through the rate constant k_2 or the activity coefficient f^* of the critical complex according to the equation^{*)} $k_2 = k_{2,0}/f^*$, where $k_{2,0}$ is the value of k_2 at infinite dilution of components of variable concentration, when $f^* = 1$. The f^* is thus varied by an amount which is hardly accounted for by the well-known neutral salt effect dealt with by Brönsted.

References

- 1) K. TANABE and Y. WATANABE, this Journal **7**, 79 (1959).
- 2) J. HORIUTI, K. TANABE and K. TANAKA, *ibid.* **3**, 147 (1955).
- 3) J. HORIUTI, *ibid.* **1**, 8 (1948).

*) For a step l in a homogeneous fluid, the forward rate \bar{v}_l for unit volume is given as³⁾,

$$\bar{v}_l = \kappa \frac{kT}{h} \cdot \frac{Q^{\delta^*}}{p^{\delta_l^I}},$$

where Q^{δ^*} is the partition function of the critical complex³⁾ δ^* and the $p^{\delta_l^I}$ the BOLTZMANN factor of the chemical potential $\mu^{\delta_l^I}$ of the initial complex δ_l^I . Since $p^{\delta_l^I}$ is the reciprocal absolute activity according to a relation $RT \log a^{\delta_l^I} = \mu^{\delta_l^I}$, we have the following equations.

$$k_l = \kappa \frac{kT}{h} Q^{\delta^*} \quad \text{or} \quad k_l = k_{l,0}/f^*,$$

where

$$k_{l,0} = \kappa \frac{kT}{h} Q_0^{\delta^*} \quad \text{and} \quad f^* = Q_0^{\delta^*}/Q^{\delta^*}$$

is the statistical mechanical expression for the activity coefficient of the critical complex δ^* of the step and $Q_0^{\delta^*}$ the particular value of Q^{δ^*} at infinite dilution of components of variable concentration.