



Title	EFFECTS OF OXYGEN TENSION AND GLUCOSE CONCENTRATIONS ON IN VITRO FERTILIZATION OF BOVINE OOCYTES
Author(s)	ARASHIMA, Chie
Citation	Japanese Journal of Veterinary Research, 43(1), 54-54
Issue Date	1995-06-15
Doc URL	<a href="http://hdl.handle.net/2115/2499">http://hdl.handle.net/2115/2499</a>
Type	bulletin (article)
File Information	KJ00002398162.pdf



[Instructions for use](#)

EFFECTS OF OXYGEN TENSION AND GLUCOSE CONCENTRATIONS  
ON *IN VITRO* FERTILIZATION OF BOVINE OOCYTES

Chie ARASHIMA

*Department of Theriogenology  
Faculty of Veterinary Medicine,  
Hokkaido University, Sapporo 060, Japan*

To determine the effects of oxygen tension and concentrations of glucose in the insemination medium in the process of *in vitro* fertilization of bovine oocytes, fertilization rates (Exp. 1), sperm motility (Exp. 2) and the development of embryos after *in vitro* fertilization (Exp. 3) were examined. *In vitro* matured bovine oocytes were co-incubated with frozen-thawed sperm ( $5 \times 10^6$  cell/ml) in modified Brackett and Oliphant isotonic medium (mBO) supplemented with 2.5 mM theophylline and 0, 1.39 or 13.9 mM glucose under 5 or 20% O<sub>2</sub>. In experiment 1, insemination in 5% O<sub>2</sub> improved total and normal fertilization rates compared with 20% O<sub>2</sub> ( $p < 0.05$ ). Total and normal fertilization rates of oocytes fertilized in mBO supplemented with 1.39 or 13.9 mM glucose were higher than those fertilized in mBO without glucose ( $p < 0.05$ ). In experiment 2, there was no significant difference in the sperm motility between 5 and 20% O<sub>2</sub> after incubation for 1 hr. After incubation for 3 and 6 hrs, the percentage of motile sperm in 20% O<sub>2</sub> was higher than in 5% O<sub>2</sub> ( $p < 0.05$ ). As the concentration of glucose increased, the percentage of motile sperm was higher in either 1-, 3- or 6-hr incubation periods. That is, total and normal fertilization rates were higher under 5% O<sub>2</sub> regardless of the lower sperm motility. In experiment 3, presumptive zygotes were cultured in modified synthetic oviduct fluid medium (mSOF) supplemented with 20 kinds of amino acids, polyvinylalcohol (1 mg/ml), gentamicin sulfate (50  $\mu$ g/ml) and insulin (10  $\mu$ g/ml), without glucose and bovine serum albumin. Cleavage rates were determined after 2 days of *in vitro* culture and all of the blastocysts were obtained after 6 days of *in vitro* culture. The percentage and total cell numbers of blastocysts were not different among the conditions in which the oocytes were inseminated under 5 or 20% O<sub>2</sub> and 1.39 or 13.9 mM glucose. However, the cleavage rate in 5% O<sub>2</sub> tended to be higher than that in 20% O<sub>2</sub> ( $p = 0.067$ ). In conclusion, it was shown that the addition of glucose to the modified BO medium used for *in vitro* fertilization and incubation in 5% O<sub>2</sub> can improve fertilization rates.