



Title	STANDARD MOLAL REAL FREE ENERGIES OF SOLVATION OF MONOATOMIC IONS AND ABSOLUTE ELECTRODE POTENTIALS IN FUSED SALTS
Author(s)	NOTOYA, Reiko; MATSUDA, Akiya
Citation	JOURNAL OF THE RESEARCH INSTITUTE FOR CATALYSIS HOKKAIDO UNIVERSITY, 28(1), 1-14
Issue Date	1980-08
Doc URL	http://hdl.handle.net/2115/25070
Type	bulletin (article)
File Information	28(1)_P1-14.pdf



[Instructions for use](#)

STANDARD MOLAL REAL FREE ENERGIES OF SOLVATION OF MONOATOMIC IONS AND ABSOLUTE ELECTRODE POTENTIALS IN FUSED SALTS

By

Reiko NOTOYA* and Akiya MATSUDA*

(Received May 14, 1980)

Abstract

The standard molal real free energies of solvation of monoatomic ions $-\alpha_i^0$'s and the absolute electrode potentials φ_i^0 's of the associated redox systems in fused salts have been estimated by an empirical method proposed previously for 40 ionic species from quadrivalent cations to monovalent anions.

It has been found that α_i^0 's in fused salts are proportional to those in water with proportionality constants close to unity.

1. Introduction

The standard free energy of formation ΔF_i° of a gaseous monoatomic ion $M_i^{z_i}$ from its element M_i may be expressed at given temperature as the sum of the standard molal real free energy of solvation of the ion $-\alpha_i^0$ in a given solvent and the absolute electrode potential φ_i^0 of the redox system $M_i^{z_i}/M_i$ as deduced in a previous work¹

$$\Delta F_i^{\circ} = -\alpha_i^0 + z_i F \varphi_i^0, \quad (1)$$

where F is the Faraday and φ_i^0 is referred to the standard state of the electron in gaseous state at the given temperature. When the value of $-\alpha_i^0$ is given, the value of φ_i^0 may be estimated on the basis of Eq. (1) using the thermodynamical data for ΔF_i° .

In the present work the values of $-\alpha_i^0$ and φ_i^0 will be estimated for a series of monoatomic ions in fused salts by the empirical method developed in a previous work¹ on the basis of the standard electrode potentials of the associated redox systems.

* The Research Institute for Catalysis, Hokkaido University, Sapporo, 060 Japan.

R. NOTOYA and A. MATSUDA

2. The basic concept of the empirical method

The usual standard electrode potential $E_{ij}^o(T)$ of the redox system $M_i^{z_i}/M_j$ in a fused salt referred to an appropriate reference electrode $M_j^{z_j}/M_j$ at a given temperature T and that in aqueous solution $E_{ij,aq}^o(298)$ at $25^\circ C$ are given by the difference of the absolute potentials of the respective electrodes

$$E_{ij}^o(T) = \varphi_i^o(T) - \varphi_j^o(T), \quad (2a)$$

$$E_{ij,aq}^o(298) = \varphi_{i,aq}^o(298) - \varphi_{j,aq}^o(298) \quad (2b)$$

Using Eqs. (1)-(2 b), we have an expression for the relative value of the free energy of solvation of the ion $M_i^{z_i}$ to that of the reference ion $M_j^{z_j}$ in a fused salt as²⁾

$$-\frac{1}{z_i} \alpha_i^o(T) + \frac{1}{z_j} \alpha_j^o(T) = a_{ij} + b_{ij} \quad (3)$$

where

$$a_{ij} = -\frac{1}{z_i} \alpha_{i,aq}^o(298) + \frac{1}{z_j} \alpha_{j,aq}^o(298) - F \Delta E_{ij}^o \quad (4)$$

$$b_{ij} = \frac{1}{z_i} \left\{ \Delta F_i^o(T) - \Delta F_i^o(298) \right\} - \frac{1}{z_j} \left\{ \Delta F_j^o(T) - \Delta F_j^o(298) \right\} \quad (5)$$

$$\Delta E_{ij}^o = E_{ij}^o(T) - E_{ij,aq}^o(298). \quad (6)$$

It can be seen from Eqs. (3)-(6) that the relative value of $-\frac{1}{z_i} \alpha_i^o(T) + \frac{1}{z_j} \alpha_j^o(T)$ in a fused salt is a quantity which can be evaluated, since a_{ij} is given by the relative free energies of hydration at $25^\circ C$ and the standard electrode potential referred to that in water which are all known quantities and b_{ij} is given by the free energies of formation of gaseous ions at T and $25^\circ C$ which can be estimated from the thermodynamical data.

If we assume here an ideal ion which should exhibit no interaction with any solvent, *i.e.*, $\alpha_i^o(T) = \alpha_{i,aq}^o(298) = 0$, then the value of $a_{ij} + b_{ij}$ for such an ideal ion gives the free energy of solvation of the reference ion $\frac{1}{z_j} \alpha_j^o(T)$, as readily seen from Eq. (3). When the value of $\frac{1}{z_j} \alpha_j^o(T)$ is given, it may be possible to estimate the value of $-\alpha_i^o(T)$ from Eq. (3) and the value of $\varphi_i^o(T)$ from Eq. (1).

However, such an ideal ion should exist, if any, as a limiting approach of the ions in real systems, for instance, as the limiting value of $a_{ij} + b_{ij}$

Free Energies of Solvation of Ions and Absolute Potentials in Fused salts

TABLE I. Standard free energies of formation of gaseous monoatomic ions from elements $\frac{1}{z_1} \Delta F_1^\circ(T)$ in electron volts

Temperature Electrode	450°C °K	475°C °K	150°C 423.15°K	177°C 450.15°K	250°C 523.15°K	625°C 898.15°K	161.85°C 435°K	221.85°C 495°K	700°C 973.15°K	550°C 823.15°K	25°C 298.15°K
	723.15 °K	748.15 °K	423.15°K	450.15 °K	523.15 °K	898.15 °K	435°K	495°K	973.15 °K	823.15 °K	298.15 °K
H ⁺ /H ₂	15.38	15.35	15.63	15.61	15.55	15.21	15.62	15.57	15.14	15.28	15.72
Li ⁺ /Li	6.13	6.10	6.52	6.49	6.39	5.89	6.51	6.43	5.79	5.99	6.67
Na ⁺ /Na	5.46	5.43	5.82	5.79	5.70	5.23	5.81	5.73	5.14	5.33	5.97
Cu ⁺ /Cu	10.16	10.12	10.68**	10.61	10.49	9.86	10.63	10.54	9.74	9.99	10.85
Cu ²⁺ /Cu	15.10	15.08	15.44**	15.40	15.32	14.90	15.41	15.35	14.82	14.99	15.58
Ag ⁺ /Ag	9.51	9.46	10.01**	9.95	9.83	9.21	9.97	9.87	9.09	9.34	10.19
Be ²⁺ /Be	14.83	14.80	15.14	15.11	15.04	14.64	15.12	15.07	14.56	14.72	15.25
Mg ²⁺ /Mg	11.57	11.55	11.85	11.83	11.76	11.40	11.84	11.79	11.33	11.47	11.96
Zn ²⁺ /Zn	13.78	13.75	14.06	14.04	13.97	13.61	14.05	14.00	13.54	13.68	14.18
Cd ²⁺ /Cd	12.98	12.96	13.25	13.23	13.16	12.81	13.24	13.19	12.74	12.89	13.37
Al ³⁺ /Al	18.39	18.37	18.62	18.60	18.54	18.24	18.61	18.56	18.18	18.30	18.70
Ga ³⁺ /Ga	19.61	19.59	19.82**	19.80	19.75	19.47	19.81	19.77	19.42	19.53	19.89
In ³⁺ /In	17.87	17.86	18.08	18.06	18.01	17.74	18.07	18.03	17.68	17.80	18.16
Tl ⁺ /Tl	7.04	7.01	7.47	7.43	7.32	6.80	7.45	7.36	6.69	6.90	7.64
Sc ³⁺ /Sc	15.59	15.57	15.82	15.80	15.75	15.45	15.81	15.77	15.39	15.51	15.90
Y ³⁺ /Y	14.05	14.03	14.28	14.26	14.21	13.91	14.27	14.23	13.85	13.97	14.36
La ³⁺ /La	12.89	12.87	13.11	13.09	13.04	12.75	13.10	13.06	12.69	12.81	13.19
Ce ³⁺ /Ce	13.84	13.82	14.07	14.05	14.06	13.69	14.06	14.02	13.63	13.76	14.16
Nd ³⁺ /Nd	13.66	13.64	13.89	13.87	13.82	13.51	13.88	13.84	13.45	13.57	13.98
Gd ³⁺ /Gd	14.09	14.07	14.32	14.30	14.24	13.95	14.31	14.26	13.89	14.01	14.40
Th ⁴⁺ /Th	18.95	18.93	19.15	19.13	19.09	18.83	19.14	19.10	18.77	18.88	19.22
U ³⁺ /U	14.23	14.20	14.48	14.45	14.39	14.07	14.47	14.42	14.00	14.14	14.57
Sn ²⁺ /Sn	12.01	11.99	12.27	12.25	12.18	11.84	12.26	12.21	11.78	11.92	12.38
Pb ²⁺ /Pb	11.68	11.66	11.95	11.92	11.86	11.52	11.94	11.89	11.45	11.59	12.07
Ti ²⁺ /Ti	12.01	11.98	12.33	12.30	12.22	11.81	12.31	12.25	11.73	11.90	12.45
Zr ⁴⁺ /Zr	20.76	20.74	20.96	20.94	20.89	20.63	20.95	20.91	20.57	20.68	21.03
Hf ⁴⁺ /Hf	19.71	19.69	19.91	19.89	19.84	19.58	19.90	19.86	19.52	19.63	19.98
V ²⁺ /V	12.39	12.36	12.71	12.68	12.60	12.19	12.69	12.63	12.10	12.28	12.83
Cr ²⁺ /Cr	12.77	12.75	13.01***	13.05	12.98	12.58	13.07	13.01	12.50	12.66	13.20
Cr ³⁺ /Cr	18.38	18.35	18.63	18.61	18.55	18.22	18.63	18.58	18.15	18.29	18.73
Cl ⁻ /½Cl ₂	2.763	2.76	2.73	2.74	2.75	2.768	2.735	2.743	2.768	2.767	2.65
Br ⁻ /½Br ₂	2.742	2.74	2.72	2.72	2.73	2.745	2.72	2.726	2.744	2.744	2.66
I ⁻ /½I ₂	2.618	2.62	2.58	2.59	2.61	2.617	2.586	2.606	2.614	2.618	2.51
Mn ²⁺ /Mn	12.87	12.84	13.19	13.16	13.08	12.67	13.18	13.11	12.59	12.76	13.31
Fe ²⁺ /Fe	13.36	13.33	13.69	13.66	13.58	13.16	13.68	13.61	13.07	13.24	13.82
Fe ³⁺ /Fe	19.00	18.98	19.25	19.23	19.17	18.84	19.25	19.20	18.77	18.91	19.37
Co ²⁺ /Co	14.17	13.88	14.23	14.48	14.40	13.97	14.49	14.43	13.88	14.06	14.63
Rh ³⁺ /Rh	20.22	20.20	20.46	20.44	20.38	20.06	20.45	20.41	20.00	20.13	20.55
Ni ²⁺ /Ni	14.46	14.43	14.78	14.76	14.68	14.25	14.77	14.71	14.17	14.34	14.91
Pd ²⁺ /Pd	15.21	15.18	15.54	15.52	15.44	15.01	15.53	15.47	14.92	15.10	15.67
Pt ²⁺ /Pt	15.62	15.59	15.59	15.92	15.84	15.41	15.94	15.87	15.32	15.50	16.08

R. NOTOYA and A. MATSUDA

extrapolated to $-\frac{1}{z_i} \alpha_{i,\text{aq}}^0(298) = 0$ when $a_{ij} + b_{ij}$ is plotted against $-\frac{1}{z_i} \alpha_{i,\text{aq}}^0$ (298) for a series of real ions.

3. Graphic presentation of $a_{ij} + b_{ij}$ vs. $-\frac{1}{z_i} \alpha_{i,\text{aq}}^0(298)$

The values of a_{ij} for a series of monoatomic ions in fused salts were estimated in a previous work². The values of b_{ij} for these ions are estimated in the present work using the values of ΔF_i^o 's at a temperature T and 298°K. The thermodynamical data used for the calculation of ΔF_i^o are those in the tables of JANAF³, Samsonov⁴, Ryabin *et al*⁵, CRC Handbook of Weast and Astle⁶ and Matsuda⁷. The standard electrode potentials in




Fig. 1. $\frac{1}{z_i} \alpha_i^0(T)$ vs. $\frac{1}{z_i} \alpha_{i,\text{aq}}^0(298)$ in LiCl-KCl at 450°C.

Free Energies of Solvation of Ions and Absolute Potentials in Fused salts




Fig. 2. $\frac{1}{z_i} \alpha_i^0(T)$ vs. $\frac{1}{z_i} \alpha_{i,\text{aq}}^0(298)$ in $\text{MgCl}_2\text{-NaCl-KCl}$ at 475°C .

molal scale in fused salts are quoted from the table of Plambeck⁸ and those in water from Dobos⁹. The values of $\Delta F_i^\circ/z_i$ estimated for monoatomic ions are listed in Table I, where the double asterisks ** in the column of 423°K means the value of ΔF_i° at 408°K and *** that at 491°K which are quite close to that at 423°K within the accuracy of the present treatment.

The values of $a_{ij} + b_{ij}$ estimated in this way for a number of monoatomic ions in fused salts are plotted against the values of $-\frac{1}{z_i} \alpha_{i,\text{aq}}^0(298)$ in Figs. (1)-(10). In these Figures the values of $a_{ij} + b_{ij}$ for an ion is measured from that of the reference ion which is chosen arbitrarily on the vertical axis.

R. NOTOYA and A. MATSUDA




Fig. 3. $\frac{1}{z_i} \alpha_i^0(T)$ vs. $\frac{1}{z_i} \alpha_{i,aq}^0(298)$ in $\text{AlCl}_3\text{-NaCl-KCl}$ at 150°C.

4. Estimation of $-\frac{1}{z_j} \alpha_j^0(T)$ and $\varphi_j^0(T)$

It can be seen from these Figures that $a_{ij} + b_{ij}$ changes linearly with $-\frac{1}{z_i} \alpha_{i,aq}^0(298)$ as

$$a_{ij} + b_{ij} = a_0 + \beta x \quad (7)$$

where x denotes $-\frac{1}{z_i} \alpha_{i,aq}^0(298)$, β is a constant characteristic to the solvent. a_0 is the value of $a_{ij} + b_{ij}$ which is obtained from the intersection of this line with the vertical axis, and a_0 may reveal $\frac{1}{z_j} \alpha_j^0(T)$ for the reference

Free Energies of Solvation of Ions and Absolute Potentials in Fused salts




Fig. 4. $\frac{1}{z_i} \alpha_i^0(T)$ vs. $\frac{1}{z_i} \alpha_{i,aq}^0(298)$ in $\text{LiNO}_3\text{-KNO}_3$ at 177°C .




Fig. 5. $\frac{1}{z_i} \alpha_i^0(T)$ vs. $\frac{1}{z_i} \alpha_{i,aq}^0(298)$ in $\text{NaNO}_3\text{-KNO}_3$ at 250°C .

R. NOTOYA and A. MATSUDA




Fig. 6. $\frac{1}{z_i} \alpha_i^o(T)$ vs. $\frac{1}{z_i} \alpha_{i,aq}^o(298)$ in $\text{Li}_2\text{SO}_4-\text{K}_2\text{SO}_4$ at 625°C .

ion according to Eqs. (3) and (7), since $-\frac{1}{z_i} \alpha_i^o(T)$ may be assumed to be zero at $x=0$.

In this way we have a proportionality relation between $-\frac{1}{z_i} \alpha_i^o(T)$ and x from the linearity between $a_{ij}+b_{ij}$ and x on the basis of Eqs. (7) and (3),

$$-\frac{1}{z_i} \alpha_i^o(T) = \beta x, \quad (8)$$

which is also shown by the same line of $a_{ij}+b_{ij}$ vs. x with the origin on the vertical axis displaced by a_0 , as shown in these Figures.

The values of $-\alpha_j^o(T)$ for the reference ion, the absolute potential $\varphi_j^o(T)$ for the reference electrode and the proportionality constant β are listed in Table I for a series of fused salts. It is surprising that β is quite

Free Energies of Solvation of Ions and Absolute Potentials in Fused salts




Fig. 7. $\frac{1}{z_i} \alpha_i^0(T)$ vs. $\frac{1}{z_i} \alpha_{i,aq}^0(298)$ in NaSCN-KSCN at 161.85°C.

TABLE II. Standard molal real free energies of solvation of reference ions, absolute potentials of reference electrodes, and proportionality constant β in fused salts, standard deviation $\pm\sigma$ of $-\frac{1}{z_i} \alpha_i^0(T)$ from Eq. (8).

Fused Salt	Temp. (K)	Reference Ion	$-\frac{1}{z_j} \alpha_j^0(T)$ (eV)	$\varphi_j^0(T)$ (v)	β	$\pm\sigma$ (eV)
LiCl-KCl	723	Ag ⁺	5.08	4.43	0.955	0.23
MgCl ₂ -NaCl-KCl	748	Ag ⁺	4.77	4.69	0.932	0.23
AlCl ₃ -NaCl-KCl	423	Ag ⁺	5.39	4.62	0.951	0.30
LiNO ₃ -KNO ₃	450	Ag ⁺	4.65	5.30	0.974	0.11
NaNO ₃ -KNO ₃	523	Ag ⁺	4.88	4.95	0.946	0.13
Li ₂ SO ₄ -K ₂ SO ₄	898	Ag ⁺	5.71	3.50	1.087	0.30
NaSCN-KSCN	435	Ag ⁺	4.83	5.14	0.922	0.33
CH ₃ COOLi-CH ₃ COONa-	495	Zn ²⁺	10.77	3.23	1.026	0.05
CH ₃ COOK						
NaPO ₃ -KPO ₃	973	Ag ⁺	4.88	4.12	0.990	0.03
Li ₂ CO ₃ -Na ₂ CO ₃	823	Ag ⁺	5.06	4.29	1.019	0.02

R. NOTOYA and A. MATSUDA




Fig. 8. $\frac{1}{z_i} \alpha_i^o(T)$ vs. $\frac{1}{z_i} \alpha_{i,aq}^o(298)$ in $\text{CH}_3\text{COOLi}-\text{CH}_3\text{COONa}-\text{CH}_3\text{COOK}$ at 221.85°C .

close to unity in any fused salt available as in the case of organic and inorganic non-aqueous solvents reported in the previous work⁹. The standard deviation $\pm\sigma$ of $-\frac{1}{z_i} \alpha_i^o(T)$ from the proportionality relation Eq. (8) are found in fused salts to be less than 0.33 ev from Table II.

5. Estimation of $-\frac{1}{z_i} \alpha_i^o(T)$ and $\varphi_i^o(T)$

The values of $-\frac{1}{z_i} \alpha_i^o(T)$ of the ions given in Figs. (1)-(10) are listed in Table III. The values of $-\frac{1}{z_i} \alpha_i^o(T)$ of the ions for which we have no experimental data of the standard electrode potential of the associated redox system can be calculated from Eq. (8) using the value of β obtained from the Figures. The values of $-\frac{1}{z_i} \alpha_i^o(T)$ calculated in this way are

Free Energies of Solvation of Ions and Absolute Potentials in Fused salts




Fig. 9. $\frac{1}{z_i} \alpha_i^0(T)$ vs. $\frac{1}{z_i} \alpha_{i,aq}^0(298)$ in $\text{NaPO}_3\text{-KPO}_3$ at 700°C .




Fig. 10. $\frac{1}{z_i} \alpha_i^0(T)$ vs. $\frac{1}{z_i} \alpha_{i,aq}^0(298)$ in $\text{Li}_2\text{CO}_3\text{-Na}_2\text{CO}_3$ at 550°C .

R. NOTOYA and A. MATSUDA

TABLE III Standard molal real free energies of solvation
of monoatomic ions in fused salts $-\frac{1}{z_1} \alpha_1^o(T)$
in electron volts[#]

Temper- ature $M_i z_i$	450°C	475°C	150°C	177°C	250°C	625°C	161.85 °C	221.85 °C	700°C	550°C	25°C
	723.15 °K	748.15 °K	423.15 °K	450.15 °K	523.15 °K	898.15 °K	435°K	495°K	973.15 °K	823.15 °K	298.15 °K
H ⁺	10.79*	10.54*	10.75*	11.01*	10.69*	12.29*	10.41*	11.60*	11.19*	11.52*	11.30
Li ⁺	4.56	4.94*	5.04*	5.16*	5.01*	5.76*	4.88*	5.44*	5.25*	5.40*	5.30
Na ⁺	3.53	3.97*	4.05*	4.15*	4.03*	4.63*	3.93*	4.37*	4.22*	4.34*	4.26
Cu ⁺	5.95	5.61	6.07	5.76*	5.59*	6.57	5.94	6.07*	5.88	6.03*	5.91
Cu ²⁺	10.21	9.96	10.07	10.50*	10.20*	11.29	9.93*	11.06*	10.67	10.99*	10.78
Ag ⁺	5.08	4.77	5.39	4.65	4.88	5.71	4.83	5.09*	4.88	5.06	4.96
Be ²⁺	11.72	11.68*	11.92*	12.21*	11.86*	13.62*	11.55*	12.86*	12.40*	12.77*	12.53
Mg ²⁺	9.00	8.74	9.40*	9.88*	9.35*	10.74*	9.10*	10.14*	9.78*	10.07*	9.88
Zn ²⁺	10.19	9.85	9.92	10.24*	9.84	11.43*	9.63	10.77	10.40*	10.71*	10.51
Cd ²⁺	9.14	8.81	8.95	9.25	8.78	10.16*	8.55	9.58	9.24*	9.52*	9.34
Al ³⁺	15.00	14.80	14.74	15.54*	15.09*	17.34*	14.70*	16.37*	15.79*	16.26*	15.95
Ga ³⁺	15.59	14.91*	15.34	15.58*	15.13*	17.39*	14.74*	16.41*	15.83*	16.30*	15.99
In ³⁺	13.76	13.13*	13.39*	13.72*	13.32*	15.31*	13.20	14.43	13.94*	14.35*	14.08
Tl ⁺	3.34	3.08	3.39*	3.39	3.11	3.87*	2.85	3.61	3.52*	3.63*	3.56
Sc ³⁺	13.00	12.64*	12.90*	13.21*	12.83*	14.74*	12.50*	13.92*	13.42*	13.82*	13.56
Y ³⁺	11.74	11.48*	11.71*	11.99*	11.65*	13.38*	11.34*	12.63*	12.18*	12.55*	12.31
La ³⁺	10.59	10.39*	10.60*	10.85*	10.54*	12.11*	10.27*	11.43*	11.03*	11.36*	11.14
Ce ³⁺	11.60	11.39*	11.62*	11.90*	11.56*	13.29*	11.26*	12.54*	12.10*	12.46*	12.22
Nd ³⁺	11.33	11.19*	11.42*	11.69*	11.35*	13.05*	11.06*	12.32*	11.88*	12.23*	12.00
Gd ³⁺	11.73	11.54*	11.78*	12.06*	11.71*	13.46*	11.41*	12.71*	12.25*	12.62*	12.38
Th ⁴⁺	16.12	15.57*	15.89*	16.27*	15.80*	18.16*	15.39*	17.14*	16.53*	17.02*	16.70
U ³⁺	11.29	10.95	11.37*	11.64*	11.31*	12.99*	11.01*	12.26*	11.83*	12.18*	11.95
Sn ²⁺	7.94	7.68	7.71*	7.89*	7.66*	8.81*	7.46*	8.31*	8.02*	8.26*	8.10
Pb ²⁺	7.63	7.38	7.38*	7.56*	7.48	8.44*	7.15*	8.06	7.68*	7.91*	7.76
Ti ²⁺	8.59	9.12*	9.30*	9.53*	9.25*	10.63*	9.01*	10.04*	9.68*	9.97*	9.78
Zr ⁴⁺	17.42	16.91*	17.26*	17.67*	17.16*	19.72*	16.72*	18.62*	17.96*	18.49*	18.14
Hf ⁴⁺	16.17	16.09*	16.42*	16.81*	16.33*	18.77*	15.91*	17.71*	17.08*	17.60*	17.26
V ²⁺	8.77	8.94*	9.12*	9.34*	9.07*	10.43*	8.84*	9.84*	9.49*	9.78*	9.59
Cr ²⁺	9.04	8.69	9.02	9.43*	9.16*	10.52*	8.92*	9.93*	9.58*	9.87*	9.68
Cr ³⁺	14.36	14.04	14.29*	14.63*	14.21*	16.33*	13.84*	15.42*	14.87*	15.31*	15.02
Cl ⁻	-2.698	-2.750	-3.198	-3.01	-2.90*	-3.34*	-2.83*	-3.15*	-3.04*	-3.13*	-3.07
Br ⁻	-2.547	-2.62*	-2.946	-2.70	-2.68	-3.06*	-2.59*	-2.88*	-2.78*	-2.86*	-2.81
I ⁻	-2.288	-2.31*	-2.36*	-2.34	-2.29	-2.70*	-2.29*	-2.55*	-2.45*	-2.53*	-2.48
Mn ²⁺	9.11	8.73	9.03*	9.24*	8.98*	10.32*	8.75*	9.74*	9.39*	9.67*	9.49
Fe ²⁺	9.38	9.06	9.36*	9.59*	9.31*	10.58	9.07*	10.10*	9.74*	10.03*	9.84
Fe ³⁺	14.61	14.38	14.24*	14.58*	14.16*	16.28*	13.80*	15.36*	14.82*	15.26*	14.97
Co ²⁺	10.01	9.67	9.98*	10.22*	9.93*	11.18	9.23	10.77*	10.38*	10.72	10.49
Rh ³⁺	16.26	14.29*	14.58*	14.93*	14.50*	17.17	14.13*	15.73*	15.17*	15.63*	15.33
Ni ²⁺	10.10	9.77	10.22*	10.46*	10.22	11.68*	9.90*	11.02*	10.63*	10.92	10.74
Pb ²⁺	10.27	9.93	9.76*	9.99*	9.71*	10.97	9.45*	10.53*	10.16*	10.46*	10.26
Pt ²⁺	10.46	10.13	10.08	10.19*	9.90*	11.37*	9.64*	10.74*	10.35*	10.66*	10.46
e ⁻	-1.49*	-1.45*	-1.48*	-1.52*	-1.48*	-1.70*	-1.44*	-1.60*	-1.54*	-1.59*	-1.56

Free Energies of Solvation of Ions and Absolute Potentials in Fused salts

TABLE IV Absolute electrode potentials $\varphi_i^o(T)$
in fused salts[#]

Electrode	Temperature °K	450°C	475°C	150°C	177°C	250°C	625°C	161.85 °C	221.85 °C	700°C	550°C	25°C
		723.15 °K	748.15 °K	423.15 °K	450.15 °K	523.15 °K	898.15 °K	435°K	495°K	973.15 °K	823.15 °K	298.15 °K
H ⁺ /H ₂	4.59*	4.81*	4.88*	4.60*	4.86*	2.92*	5.21*	3.97*	3.96*	3.76*	4.42	
Li ⁺ /Li	1.57	1.16*	1.48*	1.33*	1.38*	0.13*	1.63*	0.99*	0.54*	0.59*	1.40	
Na ⁺ /Na	1.93	1.46*	1.77*	1.64*	1.67*	0.60*	1.88*	1.36*	0.92*	0.99*	1.71	
Cu ⁺ /Cu	4.21	4.51	4.61	4.85*	4.90*	3.29	4.69	4.47*	3.85	3.96*	4.94	
Cu ²⁺ /Cu	4.89	5.12	5.37	4.90*	5.12*	3.61	5.48*	4.29*	4.15	4.00*	4.77	
Ag ⁺ /Ag	4.43	4.69	4.62	5.30	4.95	3.50	5.14	4.78*	4.12	4.29	5.22	
Be ²⁺ /Be	3.12	3.12*	3.22*	2.90*	3.18*	1.02*	3.57*	2.21*	2.16*	1.95*	2.72	
Mg ²⁺ /Mg	2.57	2.81	2.45*	1.95*	2.41*	0.66*	2.74*	1.65*	1.55*	1.40*	2.08	
Zn ²⁺ /Zn	3.59	3.90	4.14	3.80*	4.13	2.18*	4.42	3.23	3.14*	2.97*	3.66	
Cd ²⁺ /Cd	3.84	4.15	4.31	3.98	4.38	2.65*	4.69	3.61	3.50*	3.37*	4.02	
Al ³⁺ /Al	3.39	3.57	3.88	3.06*	3.45*	0.90*	3.91*	2.19*	2.39*	2.04*	2.75	
Ga ³⁺ /Ga	4.01	4.68*	4.48	4.22*	4.62*	2.08*	5.07*	3.36*	3.59*	3.23*	3.90	
In ³⁺ /In	4.12	4.73*	4.69*	4.34*	4.69*	2.43*	4.87	3.60	3.74*	3.45*	4.08	
Tl ⁺ /Tl	3.70	3.93	4.08*	4.04	4.12	2.93*	4.60	3.75	3.17*	3.27*	4.08	
Sc ³⁺ /Sc	2.60	2.93*	2.92*	2.59*	2.92*	0.71*	3.31*	1.85*	1.97*	1.69*	2.34	
Y ³⁺ /Y	2.32	2.55*	2.57*	2.27*	2.56*	0.53*	2.93*	1.60*	1.67*	1.42*	2.05	
La ³⁺ /La	2.30	2.48*	2.51*	2.24*	2.50*	0.64*	2.83*	1.63*	1.66*	1.45*	2.05	
Ce ³⁺ /Ce	2.24	2.43*	2.45*	2.15*	2.50*	0.40*	2.80*	1.48*	1.53*	1.30*	1.94	
Nd ³⁺ /Nd	2.33	2.45*	2.47*	2.18*	2.47*	0.46*	2.82*	1.52*	1.57*	1.34*	1.98	
Gd ³⁺ /Gd	2.36	2.53*	2.54*	2.24*	2.53*	0.49*	2.90*	1.55*	1.64*	1.39*	2.02	
Th ⁴⁺ /Th	2.84	3.36*	3.26*	2.86*	3.29*	0.67*	3.75*	1.96*	2.24*	1.86*	2.52	
U ³⁺ /U	2.93	3.25	3.11*	2.81*	3.08*	1.08*	3.46*	2.16*	2.17*	1.96*	2.62	
Sn ²⁺ /Sn	4.07	4.31	4.56*	4.36*	4.52*	3.03*	4.80*	3.90*	3.76*	3.66*	4.28	
Pb ²⁺ /Pb	4.05	4.28	4.57*	4.36*	4.38	3.08*	4.79*	3.83	3.77*	3.68*	4.29	
Ti ²⁺ /Ti	3.42	2.36*	3.03*	2.77*	2.97*	1.18*	3.30*	2.21*	2.05*	1.93*	2.67	
Zr ⁴⁺ /Zr	3.34	3.83*	3.70*	3.27*	3.73*	0.91*	4.23*	2.29*	2.61*	2.19*	2.89	
Hf ⁴⁺ /Hf	3.54	3.60*	3.49*	3.08*	3.51*	0.81*	3.99*	2.15*	2.44*	2.03*	2.72	
V ²⁺ /V	3.62	3.42*	4.59*	3.34*	3.53*	1.76*	3.85*	2.79*	2.61*	2.50*	3.24	
Cr ²⁺ /Cr	3.73	4.06	3.99	3.62*	3.82*	2.06*	4.15*	3.08*	2.92*	2.79*	3.52	
Cr ³⁺ /Cr	4.02	4.31	4.34*	3.98*	4.34*	1.89*	4.79*	3.16*	3.28*	2.98*	3.71	
Cl ⁻ /Cl ₂	5.46	5.51	5.93	5.75	5.65*	6.11*	5.57*	5.89*	5.81*	5.90*	5.78	
Br ⁻ /Br ₂	5.29	5.36*	5.67	5.42	5.42	5.81*	5.31*	5.61*	5.52*	5.60*	5.49	
I ⁻ /I ₂	4.91	4.93*	4.94*	4.93	4.90	5.32*	4.88*	5.16*	5.06*	5.15*	4.95	
Mn ²⁺ /Mn	3.76	4.11	4.16*	3.92*	4.10*	2.35*	4.43*	3.37*	3.20*	3.09*	3.37	
Fe ²⁺ /Fe	3.98	4.27	4.33*	4.07*	4.27*	2.58	4.61*	3.51*	3.33*	3.21*	3.98	
Fe ³⁺ /Fe	4.39	4.60	5.01*	4.65*	5.01*	2.56*	5.45*	3.84*	3.95*	3.65*	4.38	
Co ²⁺ /Co	4.16	4.22	4.25*	4.26*	4.47*	2.79	5.26	3.66*	3.50*	3.34	4.14	
Rh ³⁺ /Rh	3.96	5.91*	5.88*	5.51*	5.88*	2.89	6.32*	4.68*	4.83*	4.50*	5.22	
Ni ²⁺ /Ni	4.36	4.66	4.56*	4.30*	4.46	2.57*	4.87*	3.69*	3.54*	3.42	4.17	
Pd ²⁺ /Pd	4.94	5.25	5.78*	5.53*	5.73*	4.04	6.08*	4.91*	4.76*	4.64*	5.41	
Pt ²⁺ /Pt	5.15	5.46	5.88	5.73*	5.94*	4.04*	6.30*	5.13*	4.97*	4.84*	5.62	
e ⁻ /e _m	1.49*	1.45*	1.48*	1.52*	1.48*	1.70*	1.44*	1.60*	1.54*	1.59*	1.56	

#) The solvent for each column in Table III and IV is the fused salt in Table II specified with the same temperature, except water for the last column.

R. NOTOYA and A. MATSUDA

also listed in Table III with single asterisk *. The values of $-\frac{1}{z_i} \alpha_{i,aq}^{\circ}$ (298) in water are quoted from the previous work⁷.

The values of $\varphi_i^{\circ}(T)$ calculated from the standard potential by Eq. (2 a) using the value of $\varphi_j^{\circ}(T)$ of the reference electrode are listed in Table IV.

The values of $\varphi_i^{\circ}(T)$ calculated by Eq. (1) using the value of $-\frac{1}{z_i} \alpha_i^{\circ}(T)$ and ΔF_i° are also listed in this Table with single asterisk *. The absolute potential in water at 25°C are quoted from the previous work.⁷

It is suggested that the proportionality of the free energies of solvation of monoatomic ions in non-aqueous media and in water may provide a key word for the theoretical approach to the elucidation of the solvation bonding. The numerical values of $-\alpha_i^{\circ}(T)$ and $\varphi_i^{\circ}(T)$ for monoatomic ions in fused salts estimated in the present work may play an essential role in the comparison of the absolute rate of the electron transfer reactions of these ions in different fused salts.

References

- 1) A. Matsuda, J. Res. Inst. Catalysis, Hokkaido Univ., **27**, 101 (1979).
- 2) R. Notoya and A. Matsuda, J. Res. Inst. Catalysis, Hokkaido Univ., **27**, 119 (1979).
- 3) "JANAF Thermochemical Tables", Thermal Research Laboratory, Dow Chemical Co., Project Director D. R. Stull, Reprinted by Horikoshi Research Institute, 1966-1970.
- 4) "Properties of Elements", Part I, "Physical Properties", Edited by G. V. Samsonov, Moscow, Metallurgia, 1976, Translated by Nissotushinsha.
- 5) V. A. Ryabin, M. A. Ostroumov and T. F. Svit, "Thermodynamic Properties of Materials", Handbook, Leningrad, Khimiya, 1977, Translated by Nissotushinsha.
- 6) R. C. Weast and M. J. Astle, "CRC Handbook of Chemistry and Physics", CRC Press, 1978-1979.
- 7) A. Matsuda, J. Res. Inst. Catalysis, Hokkaido Univ., **27**, 167 (1979).
- 8) J. A. Plambeck, Encyclopedia of Electrochemistry of the Elements, Ed. by A.J. Bard, Vol. 10, "Fused Salt Systems", Marcel Dekker Inc., New York and Basel, 1976.
- 9) D. Dobos, Electrochemical Data, Elsevier Sci. Pub. Co., Amsterdam-Oxford-New York, 1975.