



Title	AN EXPERIMENTAL TRIAL OF THE PREVENTION OF CANINE HERPESVIRUS INFECTION IN PUPPIES BY INJECTION OF IMMUNE SERUM AND ACYCLOVIR
Author(s)	ASANO, Machi
Citation	Japanese Journal of Veterinary Research, 43(1), 68-68
Issue Date	1995-06-15
Doc URL	http://hdl.handle.net/2115/2511
Type	bulletin (article)
File Information	KJ00002398174.pdf



[Instructions for use](#)

AN EXPERIMENTAL TRIAL OF THE PREVENTION OF
CANINE HERPESVIRUS INFECTION IN PUPPIES
BY INJECTION OF IMMUNE SERUM AND ACYCLOVIR

Machi ASANO

*Veterinary Hospital,
Faculty of Veterinary Medicine,
Hokkaido University, Sapporo 060, Japan*

The efficacy of immune serum and acyclovir (ACV) was evaluated against experimentally induced canine herpesvirus (CHV) infection in newborn pups. *In vitro* trials to establish the efficacy of ACV on CHV were also carried out.

One- to 3-day-old pups from bitches with no history of CHV infection were inoculated with $10^{7.2}$ TCID₅₀ of CHV strain GCH-1 intranasally and subcutaneously (S.C.).

Two sera (one with a titer of 1 : 64 and the other with one of 1 : 16) from bitches that had previously lost their litters as a result of CHV infection were used as immune sera. Eight pups were injected with 2 or 3 ml/day of serum having a titer of 1 : 64 S.C. for 5 successive days after viral inoculation. Five of these 8 pups survived while 3 died. Virological and histopathological examination revealed that the death of the 3 pups resulted from causes other than CHV. Both of the pups that had received 2 ml S.C. of serum having a titer of 1 : 16 died of CHV infection with moderate lesions. CHV was isolated from the tissues of these pups.

For the *in vivo* test of ACV, eight pups were injected with CHV as described above. ACV was administered S.C. at doses of 2.5, 5 and 10 mg/kg/day to 3 pups, respectively, and was intraperitoneally (I.P.) administered at doses of 5 mg/kg/day for 5 successive days after viral inoculation. None of these administrations prevented the death of the pups.

The virus was isolated from the tissues of all these pups. The I.P.-injected group had more severe CHV lesions than those in the S.C.-injected group. The S.C. injected group showed differences in lesions that correlated with the dosage administered.

For the *in vitro* trial, 0.59 μ g/ml ACV was required to reduce 50% of the cytopathic effect.

In conclusion, it was demonstrated that the injection of convalescent bitch serum with a high antibody titer might be a useful measure to protect puppies from CHV infection. It was also shown that ACV had antiviral activity against CHV strain GCH-1 *in vitro*. However, further studies are needed to establish a method for prevention of CHV infection in pups.