



| | |
|------------------|---|
| Title | Cytokines : Communication Molecules that Influence the Process of Disease |
| Author(s) | SPLITTER, Gary |
| Citation | Japanese Journal of Veterinary Research, 45(1), 3-12 |
| Issue Date | 1997-05-30 |
| DOI | 10.14943/jjvr.45.1.3 |
| Doc URL | http://hdl.handle.net/2115/2592 |
| Type | bulletin (article) |
| File Information | KJ00002398323.pdf |



[Instructions for use](#)

Cytokines : Communication Molecules that Influence the Process of Disease

Gary Splitter

(Accepted for publication : March 1, 1997)

Abstract

Subsets of T lymphocytes that produce different cytokine patterns have reinforced the cellular- and humoral-mediated duality of the immune response. Thus, different cytokines selectively produced by different T cells results in the Th1/Th2 T cell paradigm. These T cell subsets, in an ever widening circle of examined diseases, contribute to the resolution or persistence of a disease. Now, it is timely to examine the contribution of the cytokines produced in different disease states for the improvement of vaccines and therapies for domestic animals.

Key words : cytokines, T cells, vaccine

The pathological consequences for a wide range of parasitic and infectious diseases are now linked to the pattern of cytokines produced by a particular T cell subset during the disease process. Presently, the connection between cytokine patterns and disease outcome is an intense area of interest in understanding disease mechanisms and prevention in domestic animals. The secreted cytokines play an important role in activating a variety of lymphoid cells, including, B cells, cytotoxic T cells, natural killer cells, macrophages and additional cells that participate in the immune response. Cytokine production by lymphoid cells occurs in distinct patterns allowing classification of cells into subpopulations. The most readily discernible cell populations are Th1 and Th2, also termed type 1 and type 2. These two cell populations have been best identified in CD4 T cells^{4,21}. CD4 cells of the

Th1 type produce interleukin (IL)-2, interferon (IFN)- γ and tumor-necrosis factor (TNF)- β , while Th2 cells produce IL-4, IL-5 and IL-6 and IL-13. Evidence has rapidly accumulated that Th1 and Th2 cells differ in the types of antibody response stimulated by B cells that Th1 cells mediate delayed type hypersensitivity reactions, and that Th2 cells are more potent helpers for antibody production, thus providing a simple explanation for the observed dichotomy in immune responses⁴⁴. Two additional features of CD4 T cells are important in understanding the generation of helper-T cell diversity. First, each T cell subset produces cytokines that serve as autocrine growth factors and promote the differentiation of naive T cells to a given pathway. Second, the two T cell subsets produce cytokines that regulate the development and activity of the opposite subset in a paracrine manner. Thus,

IFN- γ produced by Th1 cells amplifies Th1 development and inhibits proliferation of Th2 cells, while IL-10 and IL-4 produced by Th2 cells blocks activation of Th1 cells. This cytokine pattern evident in different T cell subsets supports the dual nature of T cell dependent effector and regulatory mechanisms, and the important roles played by cytokines in controlling pathological immune responses.

In the eleven years since the first description of helper T cell subsets, studying the roles of cytokine patterns is a major focus in understanding various diseases. Now, it is timely to propose hypotheses based on cytokine patterns to explain the differentiation and functions of T cell subsets and to explore these cytokine concepts as a basis for pathology, prophylactic vaccines and therapeutic approaches for immunologic diseases.

Function of Th1 and Th2 T cells

Controversy has existed regarding the primary functions for helper-T cell subsets. Th1 cells have been proposed as responsible for cell-mediated immunity and delayed type hypersensitivity, while Th2 cells were responsible for humoral immunity^{4,44}. However, IFN- γ , considered as a Th1 cytokine, clearly is involved in determining the IgG isotype produced by B cells. Similarly, other Th1 and Th2 cytokines have biologic effects on humoral or cell-mediated immunity, respectively. Thus, the classification of cytokines into the two groups is based on an expanded understanding of function. Classification of cytokines within a particular subset is presently based on known biological activities. For example, IFN- γ activates macrophages, enhancing microbicidal activity, as well as stimulating the production of IgG antibodies. Interestingly, IgG antibodies are central in opsonization and phagocytosis of particulate microbes by macrophages. In mice the IFN- γ induced isotypes are IgG2a and IgG3¹³.

However, the IgG homologues induced by IFN- γ in domestic animals have yet to be clearly resolved²⁰. Thus, a central function of Th1 cytokines appears to be a phagocyte-mediated defense against infections based on the ability of Th1 cytokines to activate macrophages to phagocytize and destroy microbes. In addition, Th1 cytokines, such as IFN- γ and IL-2 have autocrine and paracrine effects on natural killer and cytotoxic CD8 lymphocytes further substantiating a biologic role of Th1 cytokines in cell-mediated immunity.

In contrast, Th2 cytokines, e.g., IL-4 and IL-5, participate in the switch of B cell immunoglobulin production to IgE and eosinophil activation, respectively^{24,61}. Cytokines produced by Th2 cells are known to help B cells produce IgM and non-complement-fixing IgG isotypes, IgG1, in mice but the effect of these cytokines on isotype production in domestic animals requires additional study²⁰.

Interestingly, several recently discovered cytokines appear to possess either cell-mediated or humoral immune enhancing abilities. The best characterized of these recent cytokines is IL-12 that is produced primarily by macrophages and is known to bias the immune response to a Th1 profile in contrast to IL-4 that directs a Th2 T cell response. A second cytokine, IL-14, also known as high molecular weight B cell growth factor, is produced by B cells and induces the rapid proliferation of B cell²². High concentrations of IL-14 has been observed in certain B cell tumors²². Similarly, IL-15 is expressed in skeletal muscle and serves as an anabolic agent to increase skeletal muscle mass by increasing myosin heavy chain expression⁴⁹. Muscle cell division or the rate of myoblast differentiation appear unaffected by IL-15. IL-16 is produced by CD8 T cells and binds to the CD4 receptor present on CD4 T cells. Therefore, this cytokine may function in feedback communication between CD8 and CD4 cells. IL-16 is chemo-

tactic for CD4 T cells and induces functional IL-2 receptors on CD4 T cells^{15,32,35,37}). Interestingly, IL-16 induces transmembrane signaling events similar to those noted for gp120 of human immunodeficiency virus (HIV) when it engages the CD4 ligand^{7,8,39}). Perhaps one of the functions of the gp120 of HIV is to mimic a normal cytokine, like IL-16, to induce CD4 T cell activation. Another T cell cytokine that functions in T cell activation is IL-17. Northern analysis of RNA from various cells revealed IL-17 is produced in a tightly controlled manner by CD4 T cells and in stimulated peripheral blood T cells but not B cells or unstimulated T cells²³). However, the receptor for IL-17 is produced by a wide variety cell types indicating that IL-17 acts in a proinflammatory manner on many other cells and tissues. IL-17 activates transcription factor NF- κ B and induces expression of IL-6, IL-8, granulocyte colony stimulating factor, prostaglandin E2 and surface ICAM-1 and enhances proliferation of T cells induced by a sub-optimal costimulus, phytohemagglutinin (PHA)⁶³). These findings suggest that IL-17 may constitute an early initiator of the T cell dependent inflammatory reaction and an element of the cytokine network that bridges the immune system to hematopoiesis. As with IL-16, IL-17 has an infectious mimic in Herpesvirus *saimiri*, a lymphotropic virus where the open reading frame 13 of the virus contains a sequence that has biologic activity with the IL-17 receptor⁶⁴). Lastly, IL-18, also known as interferon-gamma inducing factor, augments natural killer cell activity similar to the structurally unrelated IL-12⁴³). IL-18 enhances the production of IFN- γ and GM-CSF, while inhibiting IL-10 production in mitogen-stimulated lymphocytes⁶⁰). IL-18 differs from IL-12 because IL-18 significantly enhances IL-2 and GM-CSF production in T cell cultures, while IL-18 and IL-12 similarly induce IFN- γ production by T cells. These findings suggest IL-18 enhances T cell proliferation through an IL-2

dependent pathway augmenting Th1 cytokine production⁴³) and cytotoxicity¹⁶).

These newly identified cytokines have received minimal examination in domestic animals. The opportunity to explore the physiologic role of these cytokines in alternative species, as well as changes that occur in response to disease provides many avenues that may lead to the understanding of disease pathogenesis as well as new innovative vaccines and therapies regarding diseases of domestic animals.

T cell subsets in disease

Polarization of Th1 and Th2 T cell subsets can provide insight into immunological mechanisms of disease outcome. A number of infectious, allergic and autoimmune disorders have been associated with Th1 or Th2 cytokine profiles, suggesting the resolution or progression of given diseases pivot on the particular T cell subsets and their elaborated cytokines^{31,51,52,56}). Again, the association of a cytokine pattern with a particular T cell subset support the concept of cell- and humoral-mediated immunity orchestrated by different T cell subsets. One of the first examples of this T cell subset duality associated with resistance or susceptibility to an infectious disease were inbred mice studies with cutaneous leishmaniasis caused by *Leishmania major*⁵¹). Impressively, cytokines produced by the Th1 subset (principally IFN- γ) in one mouse strain were associated with clearance of the parasite followed by disease resistance, while a different mouse strain produced cytokines associated with the Th2 subset (principally IL-4) resulting in disease persistence followed by death. The paradigm of cytokine profiles predicting the disease outcome has been reinforced using total tissue RNA obtained during the disease process and observing the Th1/Th2 cytokine patterns without the biases introduced by cell culture and cloning^{54,62}).

Resistance to many intracellular microbes,

including bacteria, protozoa and fungi, is founded on the induction of Th1 responses initiated by IFN- γ and TNF- α that activate macrophages^{31,56}. The Th1 cytokine pattern also is evident in viral infections where natural killer cells, cytolytic CD8 T cells and neutralizing antibodies (usually with the Th1 isotype pattern) play the major role in host resistance³. In these Th1 responses, IL-12 produced by macrophages plays a pivotal role in directing the Th1 response pattern. However, the triggering events that predict IL-12 induction and the consequences of IL-12 to regulate the Th1 cytokine pattern is an active area of research. Relevant to predicting the Th1/Th2 cytokine pattern is the fact that IL-12 directly triggers the induction of Stat 4 supporting the Th1 cytokine pathway²⁶, while IL-4 triggers the induction of Stat6 and insulin-response substrate signaling Th2 cytokine pathway in T cells^{25,36,53,55}. Supporting the role of these intracellular signaling pathways in development of Th1/Th2 cytokine patterns are the findings that deletion of the IL-12⁴¹ or Stat4^{30,59} gene result in deficient Th1 responses, while deletion of the IL-4^{33,34} or Stat6 gene^{29,57,58} result in deficient Th2 responses.

Relevant to Th1/Th2 cytokine patterns, progression of human leprosy to tuberculoid or lepromatous lesions are explained by the predominance of Th1 and Th2 cytokines, respectively⁶². Also, recombinant IFN- γ has been used to successfully treat the lesions of lepromatous leprosy supporting clinical cytokine application to influence disease progression in humans^(28,45,46). Similarly, progression of acquired immunodeficiency syndrome is predicted based on an early loss of IL-2, IFN- γ and IL-12 production^{9,10}. Further, human immunodeficiency virus-1 appears to preferentially infect Th2 T cells, suggesting a mechanism for viral persistence in a Th1-deficient state⁴⁰.

A number of studies have examined the role of individual cytokines produced in response to

pathogens of domestic animals; however, few studies have detailed the relationship of Th1/Th2 cytokine profiles in such diseases. Recently, studies of bovine leukemia virus (BLV) have determined that as disease progresses from the serologically positive to persistent lymphocytosis stage a shift in cytokine profiles of Th1 to Th2 occurs⁴⁸. In the serologically positive stage IL-2, IFN- γ , and IL-12 are prominent cytokines. However, with progression to persistent lymphocytosis and tumor formation, these Th1 type cytokines decrease, while IL-10 becomes a prominent regulatory cytokine. The IL-10 present in this disease is produced by monocyte/macrophages and possesses regulatory properties that may be influential in disease progression. IL-10, originally termed cytokine synthesis inhibitory factor²¹, inhibits the expression of other cytokines especially Th1 cytokines, as well as reducing the expression of major histocompatibility complex class II molecules, mainly on macrophages, to inhibit antigen presentation¹⁷. The inability of T cells to respond to presented foreign peptides could reduce IL-2, IL-6 and IFN- γ production. Second IL-10 also induces programmed cell death in Th1 cells^{11,12}. Th1 cells express high levels of Fas ligand and can be induced to programmed cell death, whereas Th2 cells express low levels of Fas and fail to undergo apoptosis⁵⁰. Fas antigen mediates Th1 cell activation early in the immune response as well as apoptosis of Th1 cells late in the immune response¹, and only activated T cells are killed by Fas-Fas ligand interaction^{27,38}. Therefore, early in BLV infection Th1 cells may be activated, and later these activated Th1 cells may undergo programmed cell death. Thus, IL-2 and IFN- γ production might be reduced following apoptosis of Th1 cells. Such an imbalance between Th1 and Th2 cells may provoke the next disease stage of BLV infection as suggested for AIDS¹². Exploration of Th1/Th2 cytokine patterns in diseases of domestic animals provides the oppor-

tunity to discern the mechanisms that underlie the basis of a given disease. A variety of reagents and approaches now exist to identify cytokines of domestic animals.

Identification of cytokines

Cytokines are most biologically active at short range and are often confined to the radius of several cells or in the intercellular space between two engaging cells, e.g., an antigen presenting cell and a T cell. Further, cytokines usually have an extremely short half life, with biological activity lasting only a few minutes in the blood prior to enzymatic inactivation. Further, cytokines that characteristically exist as membrane bound, e.g., IL-1 α or have already bound a receptor will be problematic in detection¹⁸⁾. Thus, when selecting an assay for measuring cytokines, a particular assay may not be applicable for all circumstances. Although measurement of cytokine production in mice and humans has been aided by commercial enzyme-linked immunoabsorbent assays (ELISA), the general absence of species-specific commercial products has hampered cytokine analysis in domestic animals. In addition to ELISA methodology, bioassays, RT-PCR and flow cytometry can be used to assess cytokine production.

First, ELISA techniques are frequently used for cytokine assessment, this assay can only measure free, secreted cytokine. Further, monoclonal antibodies produced to murine or human cytokines most often detect a species-specific cytokine epitope, making such ELISA reagents useless for detection of domestic animal cytokines. Monoclonal antibodies to domestic animal cytokines are now receiving considerable attention. However, measuring cytokines in serum as an assessment of cytokine patterns suffers from the fact that cytokines principally act at short range. Similarly, measurement of cytokines from long-term cell cultures where oligoclonal expansion may not reflect the predominant

cytokine pattern or cell population that existed in the animal. A modification of the ELISA is the ELISA-spot assay where cytokine production by single cells is detected⁴²⁾. This approach may allow cytokine assessment directly from an animal.

Second, bioassays utilize a cell line possessing the cytokine receptor. However, cells may possess receptors for additional cytokines that may be present in the sample to be tested. The use of cytokine specific monoclonal antibodies that compete for the cytokine, as well as recombinant cytokine as a standard are essential assay components. Presently, the monoclonal antibodies or recombinant cytokine are unavailable for most domestic animal cytokines limiting the utility of this assay system.

Third, reverse transcriptase-polymerase chain reaction (RT-PCR) assays can be used to measure cytokine transcription. This technique offers sensitivity, rapid assessment, small amount of cells are required and many cytokines can be simultaneously analyzed. RT-PCR allows the evaluation of immune responses occurring either systemically or within a local immune environment. Reagents to detect multiple domestic animal cytokines are often unavailable; however, RT-PCR requires only the short segments of the DNA sequence from the respective cytokine gene¹⁴⁾. Further, consensus sequence regions of similar cytokine genes of other species, such as mouse and human, often can be used to design appropriate primers. Using internal competitive standards (competitors) the absolute amounts of mRNA transcripts can be determined in small numbers of cells. Although cytokine transcription is assessed and not synthesis, reports document the strong relationship between transcription levels and amount of cytokine synthesized^{2,5)}. Similarly, oligonucleotide fragments may be used in Northern blot analysis⁶⁾ for the presence of cytokines.

Fourth, flow cytometry assays can detect

cytokine synthesis in the cytosol using cytokine specific monoclonal antibodies following cell membrane permeabilization^{19,47}. This approach offers rapid, measurable detection of cytokines. Cell phenotype can also be determined; however, the number of fluorochromes that can be used simultaneously limits the number of cytokines detected in a given sample.

Conclusions

Identifying Th1/Th2 cytokine patterns produced by domestic animals during different stages of a disease provides insight into the mechanisms that underlie the basis of resistance or susceptibility. Thus, the scientific community is now positioned to test hypotheses regarding mechanisms of disease progression or resolution. With such knowledge comes the opportunity to influence the type and level of immunity through newly engineered vaccines that can preference a Th1 or Th2 cytokine pattern. Also, as concerns increase regarding antibiotic resistant bacteria, alternative methods to antibiotics are necessary to augment the host's own immune system to combat infectious agents. Knowing the role of Th1 and Th2 cytokines in particular diseases, it will be possible using chemical and biologic materials to enhance host immune responses by directing and augmenting a desirable Th1 or Th2 cytokine response. Lastly, because selected cytokines are instrumental in pivoting the development of a Th1 or Th2 cytokine response, certain cytokines or the genes encoding a given cytokine may be useful in domestic animal vaccines. Veterinarians and immunologists have a remarkable opportunity to creatively explore new solutions to long-standing diseases of domestic animals using the knowledge of individual cytokines and Th1/Th2 cytokine patterns.

Acknowledgments

This work was supported in part by the College of Agricultural and Life Sciences and the

following grants: BARD US-2367-94, NIH R01-CA59127, and USDA 96-35204-3670.

References

- 1) Alderson, M. R., Armitage, R. J. Maraskovsky, E., Tough, T. W., Roux, E., Schooley, K., Ramsdell, F. and Lynch, D. H. 1993. Fas transduces activation signals in normal human T lymphocytes *J. Exp. Med.* 178 : 2231-2235.
- 2) Barnes, P. F., Lu, S., Abrams, J. S., Wang, E., Yamamura, M. and Modlin, R. L. 1993. Cytokine production at the site of disease in human tuberculosis. *Infect. Immun.* 61 : 3482-3489.
- 3) Biron, C. A. 1994. Cytokines in the generation of immune responses to, and resolution of, virus infection. [Review] *Curr. Opin. Immunol.* 6 : 530-538.
- 4) Bottomly, K. 1988. A functional dichotomy in CD4⁺ T lymphocytes. *Immunol. Today* 9 : 268-274.
- 5) Brorson, K. A., Berverly, B., Kang, S., Lenardo, M. and Schwartz, R. H. 1991. Transcriptional regulation of cytokine genes in nontransformed T cells. Apparent constitutive signals in run-on assays can be caused by repeat sequences. *J. Immunol.* 147 : 3601-3606.
- 6) Brown, W. C., Woods, V. M., Chitko-Mckown, C. G., Hash, S. M., and Rice-Fight, A. C. 1994. IL-10 is expressed by bovine type 1 helper (Th1), type 2 helper (Th2) and unrestricted (Th0) parasite-specific T cell clones, and inhibits proliferation of all three subsets in an accessory cell dependent manner. *Infect. Immun.* 62 : 4697-4708.
- 7) Broxmeyer, H. E. 1996. Is interleukin 17, an inducible cytokine that stimulates production of other cytokines, or merely a redundant player in a sea of other biomolecules. *J. Exp. Med.* 183 : 2411-2415.
- 8) Center, D. M., Kornfeld, H. and Cruikshank, W. W. 1996. Interleukin 16 and its function as a CD4 ligand. *Immunol. Today* 17 : 476-481.
- 9) Chehimi, J., Starr, S. E., Frank, I., D'Andrea, A., Ma, X., MacGregor, R. R., Sennelier, J., and Trinchieri, G. 1994. Impaired interleukin

- 12 production in human immunodeficiency virus-infected patients. *J. Exp. Med.* 179 : 1361–1366.
- 10) Clerici, M., and Shearer, G. M. 1993. A Th2 \rightarrow Th1 switch is a critical step in the etiology of HIV infection. *Immunol. Today* 14 : 107–111.
 - 11) Clerici, M., Sarin, A., Coffman, R. L., Wynn, T. A., Blatt, S. P., Hendrix, C. W., Wolf, S. F., Shearer, G. M., and Henkart, P. A. 1994. Type 1/type 2 cytokine modulation of T-cell programmed cell death as a model for human immunodeficiency virus pathogenesis. *Proc. Natl. Acad. Sci. USA* 91 : 11811–11815.
 - 12) Clerici, M., and Shearer, G. M. 1994. The Th1–Th2 hypothesis of HIV infection: new insights. *Immunol. Today* 15 : 575–581.
 - 13) Coffman, R. L., Leberman, D. A. and Rothman, P. 1993. Mechanism and regulation of immunoglobulin isotype switching [Review]. *Adv. Immunol.* 54 : 229–270.
 - 14) Covert, J. and Splitter, G. 1995. Detection of cytokine transcriptional profiles from bovine peripheral blood mononuclear cells and CD4⁺ lymphocytes by reverse transcriptase polymerase chain reaction. *Vet. Immunol. Immunopath.* 49 : 39–50.
 - 15) Cruikshank, W., Kornfeld, H., Berman, J., Chupp, G., Keane, J., and Center, D. 1996. Biological activity of interleukin-16. *Nature* 382 : 501–502.
 - 16) Dao, T., Ohashi, K., Kayano, T., Kurimoto, M., and Okamura, H. 1996. Interferon-gamma-inducing factor, a novel cytokine, enhances Fas ligand-mediated cytotoxicity of murine T helper 1 cells. *Cell. Immunol.* 173 : 230–235.
 - 17) deWaal Malefyt, R., Haanen, J., Spits, H., Roncarolo, M.-G., teVelde, A., Figdor, C., Johnson, K., Kastelein, R., Yssel, H., and deVries, J. E. 1991. Interleukin 10 (IL-10) and viral IL-10 strongly reduce antigen-specific human T cell proliferation by diminishing the antigen-presenting capacity of monocytes via downregulation of class II major histocompatibility complex expression. *J. Exp. Med.* 174 : 915–924.
 - 18) Dinarello, C. A. 1990. An overview of the pathophysiology of cytokines. In : *Progress in Leukocyte Biology, The Physiological and Pathological Effects of Cytokines*. C. A. Dinarello, M. J. Kluger, M. C. Powanda, J. J. Oppenheim, eds., Vol. 10B. pp. 1–6, Wiley-Liss, New York.
 - 19) Elson, L. H., Nutman, T. B. Metcalfe, D. D. and Prussin, C. 1995. Flow cytometric analysis for cytokine production identifies T helper 1, T helper 2, and T helper 0 cells within the human CD4⁺CD27⁻ lymphocyte subpopulation. *J. Immunol.* 154 : 4294–4301.
 - 20) Estes, M. D. 1996. Differentiation of B cells in the bovine. Role of cytokines in immunoglobulin isotype expression. *Vet. Immunol. Immunopath.* 54 : 61–67.
 - 21) Fiorentino, D. F., Bond, M. W., and Mosmann, T. R. 1989. Two types of mouse helper T cells IV: Th2 clones secrete a factor that inhibits cytokine production by Th1 clones. *J. Exp. Med.* 170 : 2081–2095.
 - 22) Ford, R., Tamayo, A., Martin, B., Miu, K., Claypool, K. 1995. Identification of B-cell growth factors (interleukin-14; high molecular weight-B-cell growth factors) in effusion fluids from patients with aggressive B-cell lymphomas. *Blood.* 86 (1) : 283–293.
 - 23) Fossiez, F., Djossou, O., Chomarat, P., Flores-Romo, L., Ait-Yahia, S., Maat, C., Pin, J., Garrone, P., Garcia, E., Saeland, S., Branchard, D., Gaillard, C., DasMahapatra, B., Rouvier, E., Golstein, P., Banchereau, J., and Lebecque, S. 1996. T cell interleukin-17 induces stromal cells to produce proinflammatory and hematopoietic cytokines. *J. Exp. Med.* 183 : 2593–2603.
 - 24) Galli, S. J. New concepts about mast cells [review]. *N. Engl. J. Med.* 1993. 328 : 257–265.
 - 25) Hou, J., Schindler, U., Henzel, W. J., Ho, T. C., Brasseur, M., McKnight, S. L. 1994. An interleukin-4-induced transcription factor: IL-4 Stat. *Science* 265 : 1701–1706.
 - 26) Jacobson, N. G., Szabo S. J., Weber-Nordt, R. M., Zhong, Z., Schreiber, R. D., Darnell, J. E. Jr., and Murphy, K. M. 1995. Interleukin 12 signaling in T helper type 1 (Th1) cells involves tyrosine phosphorylation of signal

- transducer and activator of transcription (Stat)3 and Stat4. *J. Exp. Med.* 181 : 1755–1762.
- 27) Kabelitz, D. and Wesselborg, S. 1992. Life and death of superantigen-reactive human CD4⁺ T cell clone : staphylococcal enterotoxins induce death by apoptosis but simultaneously trigger a proliferative response in the presence of HLA-DR⁺ antigen-presenting cells. *Int. Immunol.* 4 : 1381–1388.
 - 28) Kaplan, G., Mathur, N. K., Job, C. K., Nath, I., and Cohn, Z. A. 1989. Effect of multiple interferon gamma injections on the disposal of *Mycobacterium leprae*. *Proc. Nat. Acad. Sci. USA.* 86 : 8073–8077.
 - 29) Kaplan, M. H., Schindler, U., Smiley, S. T., and Grusby, M. J. 1996. Stat6 is required for mediating responses to IL-4 and for development of Th2 cells. *Immunity* 4 : 313–319.
 - 30) Kaplan, M. H., Sun, Y.-L., Hoey, T. and Grusby, M. J. 1996. Impaired IL-12 responses and enhanced development of Th2 cells in Stat4-deficient mice. *Nature* 382 : 174–177.
 - 31) Kaufmann, S. H. E. 1993. Immunity to intracellular bacteria [review]. *Annu. Rev. Immunol.* 11 : 129–163.
 - 32) Kazatchkine, M. D. 1996. Interleukin 16 (IL-16). *Eur. J. Immunol.* 26 : 1196.
 - 33) Kope, M., LeGros, G., Bachmann, M., Lamers, M. C., Bluethmann, H., Kohler, G. 1993. Disruption of the murine IL-4 gene blocks Th2 cytokine responses. *Nature* 362 : 245–248.
 - 34) Kuhn, R., Rajewsky, K., and Muller, W. 1991. Generation and analysis of interleukin-4 deficient mice. *Science* 254 : 707–710.
 - 35) Laberge, S., Cruikshank, W. W., Beer, D. J., and Center, D. M. 1996. Secretion of IL-16 (lymphocyte chemoattractant factor) from serotonin-stimulated CD8⁺ T cells in vitro. *J. Immunol.* 156 : 310–315.
 - 36) Lederer, J. A., Perez, V. L., DesRoches, L., Kim, S. M., Abbas, A. K., and Lichtman, A. H. Cytokine transcriptional events during helper T cell subset differentiation. *J. Exp. Med.* 184 : 397–406, 1996.
 - 37) Lim, K. G., Wan, H. C., Bozza, P. I. T., Resnick, M. B., Wong, D. T., Cruikshank, W. W., Kornfeld, H., Center, D. M., Weller, P. F. 1996. Human eosinophils elaborate the lymphocyte chemoattractants IL-16 (lymphocyte chemoattractant factor) and RANTES. *J. Immunol.* 156 : 2566–2570.
 - 38) Lynch, D. H., Ramsdell, F. and Alkerson, M. R. 1995. Fas and FasL in the homeostatic regulation of immune responses. *Immunol. Today* 16 : 569–574.
 - 39) Maciaszek, J. W., Parada, N. A., Cruikshank, W. W., Center, D. M., Kornfield, H., and Viglianti, G. A. 1997. IL-16 represses HIV-1 promoter activity. *J. Immunol.* 158 : 5–8.
 - 40) Maggi, E., Mazzetti, M., Ravina, A., Annunziato, F., deCarli, M., Piccinni, M. P., Manetti, R., Carbonari, M., Pesce, A. M., del Prete, G. 1994. Ability of HIV to promote a Th1 to Th0 shift and to replicate preferentially in Th2 and Th0 cells. *Science* 265 : 244–248.
 - 41) Magram, J., Connaughton, S. E., Warriar, R. R., Carvajal, D. M., Wu, C. Y., Ferrante, J., Stewart, C., Sarmiento, U., Faherty, D. A., Gately, M. K. 1996. IL-12-deficient mice are defective in IFN gamma production and type 1 cytokine responses. *Immunity* 4 : 471–481.
 - 42) Merville, P., Pouteil-Noble, C., Wijdenes, J., Potaux, L., Touraine, J. L., and Bancheau, J. 1993. Detection of single cells secreting IFN-gamma, IL-6, and IL-10 in irreversibly rejected human kidney allografts, and their modulation by IL-2 and IL-4. *Transplantation* 55 : 639–646.
 - 43) Micallef, M. J., Ohtsuki, T., Kohno, K., Tanabe, F., Ushio, S., Namba, M., Tanimoto, T., Torigoe, K., Fuji, M., Ikeda, M., Fukuda, S., Kurimoto, M. 1996. Interferon-gamma-inducing factor enhances T helper 1 cytokine production by stimulated human T cells - synergism with interleukin-12 for interferon-gamma production. *Eur. J. Immunol.* 26 : 1647–1651.
 - 44) Mosmann, T. R. and Coffman, R. L. 1989. Th1 and Th2 cells : different patterns of lymphokine secretion lead to different functional properties. *Annu. Rev. Immunol.* 7 : 145–173.
 - 45) Murray, H. W. 1994. Interferon-gamma and host antimicrobial defense : current and future

- clinical applications. *Am. J. Med.* 97 : 459–467.
- 46) Nathan, C., Squires, K., Griffo, W., Levis, W., Varghese, M., Job, CK., Nusrat, AR., Sherwin, S., Rappoport, S., Sanchez, E. 1990. Widespread intradermal accumulation of mononuclear leukocytes in lepromatous leprosy patients treated systemically with recombinant interferon gamma. *J. Exp. Med.* 172 : 1509–1512.
 - 47) Prussin, C., and Metcalfe, D. D. 1995. Detection of intracytoplasmic cytokine using flow cytometry and directly conjugated anti-cytokine antibodies. *J. Immunol. Methods* 188 : 117–128.
 - 48) Pyeon, D., O'Reilly, K. L. and Splitter, G. A. 1996. Increased Interleukin-10 mRNA expression in tumor-bearing or persistently lymphocytotic animals infected with bovine leukemia virus. *J. Virol.* 70 : 5706–5710.
 - 49) Quinn, L. S., Haugk, K. L., and Grabstein, K. H. 1995. Interleukin-15: a novel anabolic cytokine for skeletal muscle. *Endocrinology.* 136 : 3669–3672.
 - 50) Ramsdell, F., Seaman, M. S., Miller, R. E., Picha, K. S., Kennedy, M. K. and Lynch, D. H. 1994. Differential ability of Th1 and Th2 T cells to express Fas ligand and to undergo activation-induced cell death. *Int. Immunol.* 6 : 1545–1553.
 - 51) Reiner, S. L., and Locksley, R. M. 1995. The regulation of immunity to *Leishmania major*. [review] *Annu. Rev. Immunol.* 13 : 151–177.
 - 52) Romagnani, S. 1994. Lymphokine production by human T cells in disease states. *Annu. Rev. Immunol.* 12 : 227–280.
 - 53) Ryan, J. J., McReynolds, L. J., Keegan, A., Wang, L. H., Garfein, E., Rothman, P., Nelms, K., and Paul, W. E. 1996. Growth and gene expression are predominantly controlled by distinct regions of the human IL-4 receptor. *Immunity* 4 : 123–132.
 - 54) Robinson, D. S., Hamid, Q., Ying, S., Tsicopoulos, A., Barkans, J., Bentley, A. M., Corrigan, C., Durham, S. R., and Kay, A. B. 1992. Predominant Th2-like bronchoalveolar T-lymphocyte population in atopic asthma. *N. Engl. J. Med.* 326 : 298–304.
 - 55) Seder, R. A. and Paul, W. E. 1994. Acquisition of lymphokine-producing phenotype by CD4+ T cells. *Annu. Rev. Immunol.* 12 : 635–673.
 - 56) Sher, A. and Coffman R. L. 1992. Regulation of immunity to parasites by T cells and T cell-deprived cytokines. *Annu. Rev. Immunol.* 10 : 385–409.
 - 57) Shimoda, K., vanDeursen, J., Sangster, M. Y., Sarawdr, S. R., Carson, R. T., Tripp, R. A., Chu, C., Quelle, F. W., Nosaka, T., Vignali, D. A., Doherty, P. C., Grosveld, G., Paul, W. E., Ihle, J. N. 1996. Lack of IL-4-induced Th2 response and IgE class switching in mice with disrupted Stat6 gene. *Nature* 380 : 630–633.
 - 58) Takeda, K., Tanaka, T., Shi, W., Matsumoto, M., Minami, M., Kashiwamura, S., Nakanishi, K., Yoshida, N., Kishimoto, T., Akira, S. 1996. Essential role of Stat6 in IL-4 signalling. *Nature* 380 : 627–630.
 - 59) Thierfelder, W. E., vanDeursen, J. M., Yamamoto, K., Tripp, R. A., Sarawar, S. r., Carson, R. T., Sangster, M. Y., Vignali, D. A., Doherty, P. C., Grosveld, G. C., Ihle, J. N. 1996. Requirement for Stat4 in interleukin-12-mediated responses of natural killer and T cells. *Nature* 382 : 171–174.
 - 60) Ushio, S., Namba, M., Okura, T., Hattori, K., Nukada, Y., Akita, K., Tanabe, F., Konishi, K., Micallef, M., Fujii, M., Torigoe, K., Tanimoto, T., Fukuda, S., Ikeda, M., Okamura, H., and Kurimoto, M. 1996. Cloning of the cDNA for human IFN-gamma-inducing factor, expression in *Escherichia coli*, and studies on the biologic activities of the protein. *J. Immunol.* 156 : 4247–4279.
 - 61) Wardraw, A. J., Moqbel, R. and Kay, A. B. 1995. Eosinophils: biology and role in disease. [Review] *Adv. Immunol.* 60 : 151–266.
 - 62) Yamamura, M., Uyemura, K., Deans, R. J., Weinberg, K., Rea, T. HY., Bloom, B. R., Modlin, R. L. 1991. Defining protective responses to pathogens: cytokine profiles in leprosy lesions. *Science* 254 : 277–279.
 - 63) Yao, Z., Fanslow, W. C., Seldin, M. F., Rousseau, A. M., Painter, S. L., Comeau, M. R., Crhen, J. I. and Spriggs, M. K. 1995.

- Herpesvirus Saimiri encodes a new cytokine, IL-17, which binds to a novel cytokine receptor. *Immunity* 3 : 811–821.
- 64) Yao, Z., Maraskovsky, E., Spriggs, M. K., Cohen, J. I., Armitage, R. J., and Alderson, M. R. 1996. Herpesvirus saimiri open reading frame 14, a protein encoded by T lymphotropic herpesvirus, binds to MHC class II molecules and stimulates T cell proliferation. *J. Immunol.* 156 : 3260–3266.