Title	Seasonal changes in stable carbon isotopic composition of n-alkanes in the marine aerosols from the western North Pacific: Implications for the source and atmospheric transport
Author(s)	Bendle, James A.; Kawamura, Kimitaka; Yamazaki, Koji
Citation	Geochimica et Cosmochimica Acta, 70(1), 13-26 https://doi.org/10.1016/j.gca.2005.08.013
Issue Date	2006-01-01
Doc URL	http://hdl.handle.net/2115/26123
Type	article (author version)
File Information	GCA70-1.pdf

[Instructions for use](#)

**Seasonal changes in stable carbon isotopic composition of n-alkanes
in the marine aerosols from the western North Pacific: implications
for the source and atmospheric transport.**

JAMES A. BENDLE¹, KIMITAKA KAWAMURA¹, KOJI YAMAZAKI²

¹Marine and Atmospheric Sciences Division, Institute of Low Temperature Science, Hokkaido
University, Sapporo, Japan.

²Graduate School of Environmental Earth Sciences, Hokkaido University, Kita-Ku, Sapporo, Japan.

Abstract

In order to constrain seasonal changes in the long-range atmospheric transport of land derived lipid biomarker compounds (LDLBs) over the open ocean, marine aerosol samples were collected on a biweekly basis from 1990 to 1993 at a remote island, Chichi-Jima (27°04'N, 142°13'E) in the western North Pacific. We report results of the compound specific stable carbon isotope analysis of the *n*-alkanes isolated from the marine aerosols fraction using gas chromatography/isotope ratio/mass spectrometry (GC-ir-MS). The isotopic data reveal seasonal variations in vegetation source, atmospheric pathways and transport processes. Concentrations of *n*-alkanes ranged from 0.1 to 14.1 ngm⁻³, with concentrations highest in the winter. Compound-specific isotope analysis revealed, in particular, strong seasonal changes in the $\delta^{13}\text{C}$ values of the C₂₉ and C₃₁ *n*-alkanes (biomarkers for higher plants). Lighter $\delta^{13}\text{C}$ values were observed in winter (typically -32 to -34‰) – with a transition to heavier values in summer (typically -28 to -31‰). Using a mixing equation and typical end members for C₃ and C₄ plants, we found that this is due to relative increases in the contributions from C₄ plants in the summer season. However, in a few samples significant biasing of the $\delta^{13}\text{C}$ signal by anthropogenic fossil fuel contributions was also inferred. Using backward air mass trajectory analyses, we found that the Asian continent was the major source region for C₃ plant material during winter/spring, whereas Indonesia/Australia and possibly the Americas are the source regions for C₄ material during the summer/autumn. We also found that enhanced atmospheric transport of *n*-alkanes from C₄ plants occurred in 1991 summer/autumn during a strong El Nino event, which was associated with forest and bush fires in Indonesia and Australia. In addition to providing information on contemporary processes, this study also provides a base for future paleoclimatological work in ocean sediments.

1 INTRODUCTION

Aeolian dust fallout to the open ocean was first described in the Atlantic by Darwin (Darwin, 1846) and the importance of atmospheric transport of terrestrial materials to deep-sea sediments – including the Pacific basin - has long been a major issue of oceanography (Blank et al., 1985; Rex and Goldberg, 1958; Schramm and Leinen, 1987). The study of the fate of organic aerosols has been given further impetus recently, due to a significant role in climate forcing; directly through radiative forcing effects (e.g. Chung and Seinfeld, 2002; e.g. Jayaraman, 1999; Penner et al., 1992) and indirectly, by altering the microphysical properties of clouds (e.g. Novakov and Penner, 1993; Yu, 2000). A prominent contemporary example of the importance of organic aerosols is given by the phenomenon known as the Atmospheric Brown Cloud – a coalescent cloud of anthropogenic pollutants and natural wind-bourn dust - that is having a measurable impact on Asian climate (UNEP and C⁴, 2002).

Analytical advances during the last few decades have made possible studies describing in detail the molecular and isotopic composition of the organic component of aeolian materials. A result of this work has been the demonstration that homologous series of high molecular weight (HMW) *n*-alkanes, *n*-alcohols, and fatty acids are typical land derived lipid biomarkers* (LDLBs) found in atmospheric dust (Conte and Weber, 2002; Fang et al., 2002; Gagosian et al., 1987; Kawamura, 1995; Kawamura et al., 2003; Matsumoto et al., 2001; Sicre and Peltzer, 2004), marine sediments (Gagosian and Peltzer, 1986; Huang et al., 2000; Ikehara et al., 2000; Ohkouchi et al., 1997a; Ohkouchi et al., 1997b; Ohkouchi et al., 2000; Schefuss et al., 2003a; Schefuss et al., 2003b; Westerhausen et al., 1993; Zhao et al., 2003) and lacustrine sediments (Huang et al., 2001; Huang et al., 1999; Street-Perrott et al., 1997; Zhang et al., 2004).

These lipids are synthesised by terrestrial higher plants as constituents of the epicuticular waxes (Eglinton et al., 1962; Eglinton and Hamilton, 1967). Particulate matter containing such compounds may be deflated to the troposphere either directly, by being sloughed off plant surfaces by winds (especially by sandblasting), or alternately they may be eroded from the soil reservoir during dust storms (Simoneit et al., 1977; Simoneit and Eglinton, 1977). Therefore, plant-wax lipids form a significant component of the “aerosol veil”, even above remote ocean areas such as the North Pacific (Gagosian and Peltzer, 1986; Gagosian et al., 1981; Kawamura, 1995; Kawamura et al., 2003), South Pacific (Gagosian et al., 1987; Sicre and Peltzer, 2004) and Atlantic (e.g. Simoneit et al., 1991; Simoneit et al., 1977).

The refractory nature of some LDLB compounds (e.g. HMW *n*-alkanes, *n*-alcohols and *n*-monocarboxylic acids) (e.g. Cranwell, 1981; Kawamura, 1995) means that information contained within them (e.g. isotopic composition of the source material) may be preserved in modern and geological samples without suffering major modification. Studies of contemporary LDLB samples may elucidate a number of

* The term “biomarker” refers to an organic compound with an unambiguous link to a known natural product. In this case the biomarkers are compounds emanating from the epicuticular waxes of terrestrial higher plants.

modern earth processes, such as long-range atmospheric transport, circulation (Kawamura, 1995; Ohkouchi et al., 1997a) and deposition of organic carbon (OC) (Gagosian and Peltzer, 1986; Kawamura, 1995; Prah and Muehlhausen, 1989) (including organic pollutants (Simoneit, 1997; Simoneit, 2002)) as well as the relative contributions to aerosol OC from C_3 and C_4 plant sources (Chesselet et al., 1981; Conte and Weber, 2002; Fang et al., 2002; Ohkouchi et al., 1997a; Prah and Muehlhausen, 1989; Schefuss et al., 2003a). Moreover, in addition to providing information on processes of contemporary importance, such studies also provide a base for palaeoclimatological work (Bird et al., 1995; Ohkouchi et al., 1997b; Prah and Muehlhausen, 1989; Schefuss et al., 2003b; Westerhausen et al., 1993).

n-Alkanes typically have carbon chain-lengths ranging from C_{25} to C_{35} and a strong predominance of odd-carbon-number over even-carbon-number homologues (C_{27} , C_{29} , C_{33} are most common) (Eglinton et al., 1962; Eglinton and Hamilton, 1967; Kolattukudy, 1976). High molecular *n*-alkanes are more resistant to degradation than shorter-chain homologues and other *n*-alkyl components (order of stability: *n*-alkanes > *n*-alkanoic acids > *n*-alkanols) (Cranwell, 1981). The carbon isotopic analysis of tissues from plants using the different pathways of photosynthetic carbon fixation, C_3 , C_4 , and CAM, show different levels of ^{13}C depletion (Smith and Epstein, 1971). This carbon isotopic heterogeneity between plants types is also observed within specific compound classes, including *n*-alkanes (Chikaraishi and Naraoka, 2003; Collister et al., 1994). Thus compound-specific carbon isotope (CSIA) analyses of leaf-wax lipids can determine the plant types from which the lipids were derived. C_3 plants have *n*-alkane $\delta^{13}C$ values around -35‰ (-30 to -40‰ vs. Pee Dee belemnite), whereas C_4 plants have $\delta^{13}C$ values around -20‰ (-17 to -24‰) (Chikaraishi and Naraoka, 2003; Collister et al., 1994). The third type, CAM plants, can use both carbon fixation pathways, so their $\delta^{13}C$ values are intermediate (Chikaraishi and Naraoka, 2003; O'Leary, 1981; Rieley et al., 1993). Virtually all trees, most shrubs, and cool-season grasses and sedges use the C_3 pathway, whereas C_4 photosynthesis is found in warm-season grasses, sedges and maize. CAM plants form a much smaller proportion of the global biomass and include many succulents, such as cacti (Cerling and Quade, 1993; Spicer, 1989). The stable carbon isotope compositions of *n*-alkanes have been used to determine relative contributions by C_3/C_4 plant material in marine sediments (Bird et al., 1995; Huang et al., 2000; Kuypers et al., 1999; Schefuss et al., 2003b) and aerosol dust in the eastern Atlantic (Schefuss et al., 2003a). To date, no CSIA analysis of *n*-alkanes in dust in the western Pacific has been conducted.

Here we present data on the seasonal variations of stable carbon isotope compositions of *n*-alkanes from aerosols collected from Chichi-Jima in the western North Pacific (see Figure 1). We compare the variations in carbon isotope compositions of the *n*-alkanes collected at Chichi Jima with air mass back trajectory analysis and a new quantitative map of global C_3/C_4 abundance (Still et al., 2003) (Figure 1). The data provides insight into the higher plant and anthropogenic sources and transport pathways of terrigenous lipid materials in the continental dust carried out to the ocean. Moreover, in addition to providing information on contemporary processes, this study also provides a base for future paleoclimatological work in ocean sediments. CSIA of *n*-alkanes in sediments can help resolve palaeo-wind patterns and deposition of OC over the Pacific. Alternately, if wind patterns can be constrained (e.g. by modelling) sedimentary *n*-alkanes could reveal changes in the evolution of past C_3/C_4 plant abundances and hence the carbon isotopic discrimination of

terrestrial photosynthesis - an important parameter for accurate estimates of paleo- ρCO_2 (Conte and Weber, 2002).

2 EXPERIMENTAL SECTION

2.1 Regional Setting and Sampling

Chichi-Jima is a small island ($\sim 8 \times 5$ km in size) located at $27^{\circ}04'\text{N}$; $142^{\circ}13'\text{E}$, approximately 2000 km west of the Asian continent and around 1000 km south of Tokyo (Figure 1). The region is strongly influenced by the Asian monsoon, with wind systems characterized by Westerlies during the winter to spring seasons, whereas Trade Winds dominate in summer to autumn. Figure 1 shows variations of wind direction at 700 hPa over Chichi-Jima Island over the 4-year sampling period basis from April 1990 to November 1993. During this period 69 aeolian dust samples were collected on a biweekly basis at the Satellite Tracking Center of National Aeronautic and Science Develop Agency (NASDA, elevation: 254 m). The climate of Chichi-Jima is sub-tropical with an annual rainfall of 850mm. The biomass is dominated by C_3 plants ($>99\%$ Hara, pers. comm.). There are a few species of C_4 plant, mostly sedges which are found in streams near the coast (Toyoda, 1981; Hara, pers. comm.). It is highly unlikely that there would be a significant C_4 plant seasonal influence on the aerosols collected at the elevated sampling station.

2.2 Extraction, derivatization and analysis of n-alkanes.

Aerosol samples were collected on a pre-combusted (500°C , 3 h) Pallflex quartz fiber filter (20×25 cm) using a high volume air sampler ($1 \text{ m}^3 \text{ min}^{-1}$). Detailed procedures were described in Kawamura *et al.* (2003). The sampling time was approximately 100 h to collect enough material for chemical analysis. After the sample collection, the quartz filter was placed in a pre-cleaned glass jar with a Teflon-lined screw cap and stored in a freezer at -20°C until analysis. An aliquot of the filters (10 – 20%) was extracted under reflux for 2 h with 100 mL of 0.1 M KOH in methanol, containing ca. 5% distilled water. The analytical method was described in Kawamura *et al.* (2003). Briefly, the extracts were isolated by filtration with a pre-cleaned Whatmann GF/A filter. The residue was further extracted with methanol and then methylene chloride under ultrasonification for 3 min. The extracts were combined and concentrated under vacuum and then separated into neutral and acidic fractions using the methods of Kawamura (1995). The neutral fraction was further separated into four sub-fractions using a silica gel column chromatography, that is, aliphatic hydrocarbons/*n*-alkanes (N1), aromatic hydrocarbons (N2), aldehydes and ketones (N3) and alcohols (N4) by elution with *n*-hexane, *n*-hexane/methylene chloride (2:1) mixture, methylene chloride and methylene chloride/methanol (95:5) mixture, respectively. Sub-fractionation of the acid classes is detailed in Kawamura *et al.* (2003) and Fang *et al.* (2002).

The *n*-alkanes were analyzed on a Carlo Erba Mega 5160 gas chromatograph (GC) equipped with a cool on-column injector, fused silica capillary column (Chrompack, CP-Sil 60m $\times$ 0.32mm i.d. $\times$ 0.25 μm) with an FID detector. The oven program was 70°C to 120°C at $30^{\circ}\text{C}/\text{min}$ and to 300°C (22 min) at $5^{\circ}\text{C}/\text{min}$. Structural identification was confirmed using a GC/MS (Thermoquest, Voyager) with similar GC column conditions. Authentic *n*- C_{29} alkane was used for quantification, recoveries were better than 80%. Analytical errors of the procedures were within 15%.

Laboratory procedural blanks showed that blank levels were less than 1% of the samples.

2.3 Stable Carbon Isotope Analysis.

Isotopic ratios of *n*-alkanes were determined using an HP 6890 gas chromatograph coupled with a Finnigan MAT Delta Plus isotope ratio mass spectrometer by a Finnigan MAT combustion furnace containing Cu and Ni wires which were doped with oxygen and maintained at a temperature of 850 °C. The GC was equipped with a cool on-column injector and a HP-5MS fused silica capillary column (30m × 0.32mm i.d., 0.25µm film thickness). The GC oven was programmed from 50 to 120 °C at 30 °C/min, from 120 to 300 °C at 5 °C/min and maintained at 300 °C for 22 min. The flow rate for He carrier gas was set at 1.6 mL/min. Two microliters of sample solution spiked with C₂₀ methyleicosanoate (C₂₀ FAME) as an internal standard of known isotopic composition were injected through an on-column injector. Peaks were simultaneously detected in Faraday collectors at *m/z* 44 (¹²C¹⁶O₂), 45 (¹³C¹⁶O₂ and ¹²C¹⁷O¹⁶O), and 46 (¹²C¹⁶O¹⁸O, ¹²C¹⁷O¹⁷O, and ¹³C¹⁷O¹⁶O), amplified, corrected for the presence of ¹⁷O at mass 45 using the Craig correction (Craig, 1957). The isotopic composition of *n*-alkanes is reported in the delta notation relative to the Pee Dee Belemnite (PDB) standard as follows:

$$\delta^{13}\text{C} (\text{‰}) = [({}^{13}\text{C}/{}^{12}\text{C})_{\text{sample}} / ({}^{13}\text{C}/{}^{12}\text{C})_{\text{PDB}} - 1] \times 10^3 \quad (1)$$

Isotope values reported were determined by averaging duplicate analyses whenever concentrations were sufficient. A minority of the samples (30%) contained only enough material for a single isotopic analysis of the dominant C₂₉ and C₃₁ homologues. For the duration of the analysis the condition of the machine was checked by injection of an external standard of *n*-alkanes (mixture of various carbon numbers from C₁₅ to C₃₃) of known isotopic composition. This gave a reproducibility of ~0.5‰ for all compounds (e.g. δ¹³C 0.45‰, 0.5‰ and 0.57‰ for C₂₇, C₂₉ and C₃₁ respectively). The isotopic composition of sample *n*-alkanes was calculated by co-injection with a non-coeluting internal standard (C₂₀ FAME) of known isotopic composition (δ¹³C -26.21‰).

2.4 Back-Trajectory Analysis.

Back-trajectory analyses were conducted using the Fifth-Generation NCAR/Penn State mesoscale model (MM5). The trajectory calculation was based on backward tracking of a selected air parcel released at a target location, assuming the parcel was moving along the ambient airflow. The flow pattern was updated every 6 h.

3 RESULTS AND DISCUSSION

3.1 Distribution of *n*-alkanes in aerosols

The concentrations and molecular distributions of the *n*-alkanes (0.17–14 ngm⁻³, average 1.7 ngm⁻³) collected at Chichi-Jima have been reported previously, with other lipid classes (Kawamura et al., 2003). The seasonal data is summarized in Table 1. The *n*-alkanes were characterized by a strong odd-carbon number predominance (CPI ratios, average 4.5) with a maximum at C₂₉ or C₃₁, indicating that terrestrial higher plant waxes were the primary source. Figure 3 illustrates a typical *n*-alkane class distribution, as seen in most of the samples. Absolute concentrations generally increased in winter/spring and decreased toward summer/autumn season, except for

the El Niño year of 1991 when the concentrations were relatively high in the summer/autumn (Kawamura et al., 2003). Such a seasonal trend suggested that atmospheric transport of terrestrial organic materials over the western North Pacific are enhanced during winter/spring season probably due to the strong westerly winds (see Figure 2). CPI values were generally high suggesting a minimal contribution from petroleum residues/pollution (Kawamura et al., 2003). Unresolved complex mixtures of hydrocarbons (UCM) were detected in some samples. UCM hydrocarbons are emitted to the atmosphere by the incomplete combustion of fossil fuels (e.g. Boyer and Laitinen, 1975; Simoneit and Mazurek, 1981) and are typically abundant in aerosol and rainwater samples from the urban atmosphere (e.g. Kawamura and Kaplan, 1991; Kawamura et al., 1995; Simoneit, 1984; Simoneit et al., 1988). The presence of UCMs may demonstrate the influence of polluted air from Asian countries at Chichi-Jima. However, the concentrations were 2 or 3 orders of magnitude lower than for urban samples, thus confirming (as with the high CPI values) the relatively low levels of pollution at the remote Chichi-Jima site. There were no clear seasonal trends of the CPI ratios or UCM concentrations (Kawamura et al., 2003).

3.2 Stable isotopic compositions of *n*-alkanes.

Results of the stable isotopic measurements for the major *n*-alkanes are given in Table 2. The stable isotopic compositions of the dominant HMW *n*-alkanes (C₂₉, C₃₁) fell in the range -26.6 to -34.4‰ indicating that all the samples consisted primarily of varying relative amounts of C₃ and C₄ plant-derived leaf-wax lipids (Table 2). In general, for the dominant C₂₉ and C₃₁ *n*-alkanes, lighter $\delta^{13}\text{C}$ values were observed in winter (typically -31 to -34‰) with a transition to heavier values in summer (typically -28 to -33‰) (Figure 4 a,b). Within individual samples the odd numbered *n*-alkanes are relatively more depleted in ^{13}C with increasing chain length, a typical isotopic distribution is illustrated in Figure 3. In general the C₂₉ and C₃₁ alkanes were the most ^{13}C depleted compounds in each sample. The longer-chain odd-numbered compounds, C₃₃ and C₃₅, if measurable, were isotopically enriched relative to C₂₉ and C₃₁.

Fang *et al* (2002) previously reported the compound specific $\delta^{13}\text{C}$ values measured in the fatty acid compound class extracted from the same Chichi-Jima sample set. The isotopic values measured by Fang *et al* (2002) for the HMW fatty acids were also consistent with a higher plant source of varying relative amounts of C₃ and C₄ lipids (-30‰ to -28.2‰ with a mean of -29.1‰ $\pm$ 0.7‰). They suggested that heavier ^{13}C values (-26.9‰ to -24.8‰, with a mean of -25.8‰) measured in the LMW fatty acids (C₁₂ -C₁₉) may have indicated a contribution from marine biota (Fang et al., 2002), as marine organic carbon in the 40-50°N latitude range has a $\delta^{13}\text{C}$ of around -21‰ (Chesselet et al., 1981; Fontugne and Duplessy, 1978). However, the *n*-alkane compound class data does not suggest a significant marine algal input. *n*-Alkanes synthesised by aquatic algae typically have C₁₅ or C₁₇ as the dominant homologue (Blumer et al., 1971; Gelpi et al., 1970), but these compounds were not present as aerosol particles in concentrations measurable by GC-FID or GC-ir-MS.

As well as marine derived *n*-alkanes a significant contribution of fossil fuel *n*-alkanes to the aerosols collected at Chichi-Jima is also a possibility and may bias the $\delta^{13}\text{C}$ signal derived from higher-plant waxes. There is no consensus on the most

appropriate method to assess and/or correct for non-higher plant contributions. To estimate whether a potential fossil fuel/marine contribution of *n*-alkanes influenced our samples we applied a mass balance approach adapted from the method of Ishiwatari *et al.*, (1994):

$$\delta^{13}C_{nw} = \frac{(\delta^{13}C_{nx} \times A_x) - \delta^{13}C_n}{A_x - 1} \quad (2)$$

where $\delta^{13}C_{nw}$ is the higher plant wax $\delta^{13}C$ fraction and $\delta^{13}C_{nx}$ is the fossil fuel/marine $\delta^{13}C$ fraction of an *n*-alkane $\delta^{13}C_n$ with carbon number *n* (C_n) and A_x is the fraction of the fossil fuel/marine *n*-alkanes contributing to the absolute abundance of C_n . For values of $\delta^{13}C_{nx}$, we use the $\delta^{13}C$ value of the C_{24} *n*-alkane in each sample, assuming that it is equal to the $\delta^{13}C$ value of the fossil fuel contribution. The C_{24} *n*-alkane is often absent in pure plant wax samples (Chikaraishi and Naraoka, 2003; Collister *et al.*, 1994; Eglinton *et al.*, 1962; Eglinton and Hamilton, 1967), but occurs (in small quantities) in many of the samples from Chichi-Jima. Therefore, we assume that the C_{24} *n*-alkane originates from fossil fuel/marine sources. To calculate A_x we used the ratio C_{24}/C_{29} *n*-alkane as a measure of the fraction contributed by fossil fuel/marine *n*-alkanes. Some previous authors have used CPI ratios to calculate A_x (Ishiwatari *et al.*, 1994; Schefuss *et al.*, 2003a). However, we are not satisfied that an uncontaminated/higher plant wax end member can be assigned to the mixing equation used for this method – given that natural CPI values vary so greatly (e.g. 1.4 to 40.3 in Chikaraishi and Naraoka, 2003; Collister *et al.*, 1994).

The corrected $\delta^{13}C$ values for the C_{29} and C_{31} *n*-alkanes are given in Table 2 and illustrated in Figure 4c,d with the uncorrected values for comparison. Corrections could be made for only the samples where the C_{24} *n*-alkane was abundant enough for a successful compound-specific $\delta^{13}C$ measurement. This was possible for 55% of the samples for which compound-specific $\delta^{13}C$ measurements had also been made for the C_{29} and C_{31} *n*-alkanes. For two thirds of these samples the differences in the uncorrected and corrected $\delta^{13}C$ values were insignificant ($< 0.5\text{‰}$), for one third of the samples the corrections were $> 0.5\text{‰}$ and for five samples (QFF- 194, -214, -217, -223, -420) the corrections were $> 1\text{‰}$. The potential source of the non-higher plant contributions to these samples is considered in more detail in section 3.5.

Figure 4 illustrates that the data-set which includes the corrected $\delta^{13}C$ values does not greatly affect the discernable seasonal trend in isotopic values (which become heavier in the summer/autumn seasons). This suggests that the seasonal trend must contain a highly significant natural signal that is a function of changes in the proportions of delivered C_3 and C_4 plant material. This is confirmed by the fact that measurements of UCMs and the CPI values at Chichi-Jima (which are sometimes used to infer the contribution from fossil fuel sources) did not have a similar, clear seasonal trend (Kawamura *et al.*, 2003). For consistency within the dataset we use the uncorrected values in section 3.2.1 to assess C_3 and C_4 plant contributions and to compare with the modeled backward trajectories. This is because the corrections could not be applied to all of the samples (due to variable availability of C_{24}).

Another way to access both the potential biasing of the natural C_3 / C_4 plant signal by fossil fuel/marine inputs and whether the uncorrected or partially corrected *n*-alkane data set is preferable is to compare the *n*-alkane isotopic data with that

previously obtained from fatty acids. Fatty acids are less susceptible to non-higher plant contributions than *n*-alkanes as their concentrations are typically one order of magnitude higher in higher plant waxes. Figure 5 illustrates the linear correlation between the carbon isotopic compositions ($\delta^{13}\text{C}$) of the C_{29} *n*-alkane (this paper) and the C_{24} fatty acid (Fang *et al.*, 2000). Linear correlations of higher significance existed between the C_{24} fatty acid and the uncorrected C_{29} (R^2 of 0.47) and C_{31} (R^2 of 0.35) *n*-alkane isotopic data, than for the data-sets which included the corrected values (R^2 of 0.39 and 0.19 for the C_{29} and C_{31} respectively). The greater consistency of the uncorrected *n*-alkanes with the fatty acid supports the use of the uncorrected data-set.

3.3 Backward trajectories and sources of C_4 plant-derived wax lipids in aerosols

In general lighter $\delta^{13}\text{C}$ values were observed in winter (typically -31 to -34‰) with a transition to heavier values in summer (typically -28 to -33‰) (Figure 4). If the $\delta^{13}\text{C}$ values are averaged by season this trend is even clearer for the dominant C_{29} and C_{31} *n*-alkanes (Figure 6). The difference in the $\delta^{13}\text{C}$ of the same *n*-alkanes between the winter and summer seasons may reflect a seasonal variability in the source areas and/or atmospheric transport pathways and hence differential source areas. The isotopic composition of plants can change in response to seasonal environmental stimuli. Lighter values of $\delta^{13}\text{C}$ for the C_{29} and C_{31} *n*-alkanes in several deciduous tree species have been observed in autumn leaves compared with leaves sampled at the start of the growing season ($\Delta \delta^{13}\text{C}$ -0.6 to -5‰) (Chikaraishi *et al.*, 2004; Lockheart *et al.*, 1997). Spring lipids may be isotopically heavier due to the remobilization of stored carbohydrate at the beginning of the growing season (Lockheart *et al.*, 1997). However, such phenomena can not explain the observed seasonal signal at Chichi-Jima, which is the opposite in sign (*n*-alkanes become isotopically heavier from the spring to autumn). We expect that, given the strong seasonality recorded in the prevailing winds at Chichi-Jima (Figure 2), changes in transport pathways dominate the signal at Chichi-Jima. This suggests that, every year, for a particular season the *n*-alkanes may be derived from relatively consistent sources and the aerosols are transported by similar atmospheric transport pathways and processes. In order to explore this in more detail, we estimated the source contributions of C_3 and C_4 plant material (derived from the *n*-alkane $\delta^{13}\text{C}$) and the transport pathways (by back-trajectory analysis).

Assuming that there is no significant isotopic fractionation during aerosol transport over the ocean the relative contributions to the *n*-alkanes from terrestrial C_3 and C_4 plants can be estimated by using a two-component mixing equation (Chesselet *et al.*, 1981):

$$C_{\text{alk}} = C_3 + C_4 \quad (3)$$

$$C_{\text{alk}} \delta^{13}C_{\text{alk}} = \delta^{13}C_3 \times C_3 + \delta^{13}C_4 \times C_4 \quad (4)$$

Where C_{alk} is the total concentration of an *n*-alkane, C_3 and C_4 are *n*-alkane concentrations from C_3 and C_4 plants, and $\delta^{13}C_3$ and $\delta^{13}C_4$ are the isotopic composition of the *n*-alkanes from C_3 and C_4 plant sources respectively. The end-member values for the mixing equation were averages taken from the literature and from measurements made at ILTS (see Table 3): for C_{29} , C_4 plant *n*-alkanes = 20.2‰

and C₃ plant *n*-alkanes = -35.1‰; for C₃₁, C₄ plants *n*-alkanes = 20.4‰ and C₃ plant *n*-alkanes = -35.5‰ (Chikaraishi and Naraoka, 2003; Collister et al., 1994). This results in estimates of C₄ plant contribution varying from 6 to 37% for the individual winter samples and 7 to 59% for the individual summer samples (Table 2 and Figure 4, right axis).

Fang *et al.* (2002) estimated C₄ plant contributions to the Chichi-Jima aerosols from the isotopic composition of the C₂₈ fatty acid. They detected a similar seasonal trend of relatively higher inputs of C₄ material in the summer than the winter seasons (Table 4). However the absolute estimates of C₄ contributions are consistently lower when calculated from the C₂₈ fatty acid $\delta^{13}\text{C}$ (‰C₄ 6 to 27) (Fang et al., 2002) than from the *n*-alkanes (‰C₄ of 15 to 35%) (this paper). This discrepancy may result from differential seasonal metabolism of *n*-fatty-acids and *n*-alkanes. It may also be due the use of inaccurate end-member numbers for either or both the fatty acid and *n*-alkanes and perhaps highlights that caution is needed interpreting absolute source input contributions calculated from end-member models.

The relative contributions from the C₃ and C₄ plant sources are determined by the regional meteorological conditions and C₃/C₄ production and deflation to the atmosphere in the source areas. Comparison of composite back-trajectory analysis maps for each season (Figure 7) with a global map of the fraction of C₄ biomass (Still et al., 2003) (Figure 1) reveals the potential source areas of eroded plant material that are responsible for the seasonal changes in the C₂₉ and C₃₁ *n*-alkane $\delta^{13}\text{C}$ values observed at Chichi-Jima. In general, in winter/spring, more air masses originated in north Asia where for large regions C₄ plants comprise 0% of the mean annual biomass (Figure 1) (Still et al., 2003). In contrast in summer/fall, backward trajectories show a greater diversity of source areas: less air masses originated in north Asia, instead they were more often marine (easterly trades) Southeast Asian, Indonesian and sometimes (1991 and 1992) Australian in origin. In the latter regions C₄ plants form a more significant proportion of the annual mean biomass than in northern and western Asia, for example: up to 50% in eastern Chinese provinces such as Hebei and Shanxi, up to 70% in parts of Thailand and Cambodia and up to 100% in much of northern Australia (Figure 1) (Still et al., 2003).

3.4 C₃ and C₄ plant contribution and air mass source categories

To further confirm that changes in air-mass source areas control the seasonal $\delta^{13}\text{C}$ *n*-alkane signal at Chichi-Jima each sample was allocated an air mass source category, based on the dominant source area observed in the 10-day back trajectory. The categories were: Northeast Asia (NEA), Southeast Asia (SEA), Indonesia and Australia (IA) and Marine easterlies (ME) (Figure 1, Table 2). 96% percent of winter samples fall into the SEA or NEA source area categories, in contrast only 21% of the summer samples were of NEA or SEA origin, being of IA (30%) and ME (49%) origin respectively (Table 2). Box plots of the C₂₉ and C₃₁ $\delta^{13}\text{C}$ data for each of these major source areas are illustrated in Figure 8. The box plots show that samples in the NEA and SEA categories were significantly isotopically lighter (median values of around -32 to -33‰) than those of the ME and IA categories (median values of around -30 to -31‰). This confirms that the isotopically heavier values recorded at Chichi-Jima during the summer-autumn are a function of material carried by the IA

and ME air mass sources. The interquartile ranges (25 – 75% of the data) also demonstrate that - on the basis of CSIA measurements of the *n*-alkanes - it is largely possible to discriminate between samples with an Asian continental source (NEA or SEA) and the easterly (ME) or southerly (IA) sources. However, using this method further categorisation was not possible – there was a major overlap of the interquartile ranges of the NEA and SEA categories and of the IA and ME categories (Figure 8). The lack of significant difference between SEA and NEA categories and between IA and ME categories may be due to the entrainment and mixing by air masses of *n*-alkanes from wide geographic areas. It seems that the $\delta^{13}\text{C}$ of the aerosol *n*-alkanes at Chichi-Jima are only sensitive to the major seasonal changes in the prevailing wind systems. This suggests that paleoclimatic reconstructions using sedimentary material recovered from the Chichi-Jima sector of the western Pacific may record C_3/C_4 plant abundance changes over wide geographical areas.

In order to explore the potential for various combinations of higher plant biomarker parameters to further discriminate between the different source areas we looked at a matrix of scatterplots (Figure 9). A number of combinations of parameters showed a significant separation of the NEA and SEA categories from the IA and ME categories (e.g. $\delta^{13}\text{C}$ of C_{24} fatty acids vs $\delta^{13}\text{C}$ of C_{29} *n*-alkanes; $\delta^{13}\text{C}$ of C_{31} *n*-alkanes vs UCM) but no combinations could separate out the NEA from SEA categories or the IA from ME categories.

Prior to this comparison of the global C_4 biomass with back-trajectory analysis Fang *et al* (2000) had suggested that the increase in C_4 inputs in the summer at Chichi-Jima were a result of material carried by the easterly trade winds from the North and Central Americas, where a diverse C_4 flora exists (Still *et al.*, 2003; Teeri and Stowe, 1976) (Figure 1). Indeed 6 of the 8 isotopically heaviest *n*-alkane samples ($>29.5\%$ or around $>40\%$ C_4) were of marine easterly (ME) origin. However, even 20 day back-trajectory analysis did not place the source air-mass parcels in the Americas (data not shown). Therefore, while there may be a high proportion of C_4 material of American provenance in these samples, we can not confirm the thesis of Fang *et al.*, (2000) that it is transported directly to Chichi-Jima by specific meteorological events. Our analysis further suggests that the increase in C_4 contribution observed in the summer seasons at Chichi-Jima was also the result of relatively less C_4 poor material being transported from north Asia and relatively more C_4 rich material being transported from Southeast Asia and Australia (Figure 7) as well as from the Americas.

3.5 Investigation of individual samples

In section 3.2 (Figure 4c,d) we identified five samples for which we estimated the $\delta^{13}\text{C}$ contribution from non-higher plant sources to be greater than 1%. The back trajectory fields for the samples included the following densely populated regions:

- Sample QFF-194: Japan, Korea and Northeast China.
- Sample QFF-214: Eastern China and Northern Japan.
- Sample QFF-217: Taiwan, the Philippines and Southern Japan.
- Sample QFF-222: Southern Japan and Eastern China.
- Sample QFF-420: Japan, Korea and Northeast China.

Therefore, in all of the above cases it appears that the source of the non-higher plant *n*-alkanes must be anthropogenic materials exported to Chichi-Jima as part of the “Asian outflow” which is frequently polluted by fossil fuel/biofuel derivatives (e.g. Fuelberg et al., 2003; Liu et al., 2003; Woo et al., 2003). However, many of the other samples collected at Chichi-Jima also had back-trajectories which included these regions. Therefore a precise reason for the above samples having a higher anthropogenic signal at Chichi-Jima may be due to temporal changes in the production of or transport efficiency of anthropogenic material.

Previously Kawamura *et al.* [2003] noted that absolute concentrations (ngm^{-3}) of terrestrial biomarkers were generally lower in the summer to autumn seasons due to the predominance of maritime trade winds. However, higher concentrations were recorded during the summer and fall of 1991. During this period the highest proportion of C_4 material at Chichi-Jima was discovered for the C_{29} *n*-alkane (Figure 4a). The year of 1991 was characterized by a strong El Niño event, expressed meteorologically by weaker subtropical high-pressure systems in the western Pacific and more frequent development of tropical low systems (typhoons). The back-trajectories demonstrated that more air masses associated with this year originated in Southeast Asia, including Indonesia, Kalimantan and Australia (see Figure 7 and Kawamura et al., 2003). The C_4 contributions to aerosols at Chichi-Jima were therefore enhanced by emissions from these areas. Samples QFF-216 and QFF-219 (Figure 4a) had high concentrations of C_4 plant material (45 and 55% respectively for the C_{29} *n*-alkane), these samples also had back-trajectories which reached to northern Australia (see Figure 10), a region with a biomass of up to 100% C_4 plants over large areas (Still et al., 2003). Also during the 1991 El Niño event, the air over the western equatorial Pacific became dry and hot and many forest fires occurred in southeast Asia (Kita et al., 2000) and Australia (Verdon et al., 2004), as well as the following year. During the forest fires and biomass burning, lipid class compounds, originally present as leaf waxes, could have been significantly emitted as smoke particles (without severe oxidative degradation) via volatilization (Abas et al., 1995; Oros and Simoneit, 2001). Thereafter lipids associated with smoke produced by smoldering and flaming processes may have been uplifted by convection and transported long distances over the Pacific Ocean, hence, enhancing the input of C_4 plant material detected at Chichi-Jima.

4 CONCLUSIONS

The *n*-alkane class - in the remote marine aerosols collected from the western North Pacific - contains predominantly long-chain compounds with a strong dominance of odd carbon numbered compounds. These lipids are derived from epicuticular waxes of terrestrial plants and indicate the importance of vegetation sources for organic matter in aerosols. Compound Specific Isotope Analysis (CSIA) of the *n*-alkanes reveals relatively greater contribution of C_3 plant material in the winter/spring and C_4 material in the summer/fall at Chichi-Jima. Comparison of backward trajectory analysis and a global map of C_3/C_4 biomass showed that higher plant *n*-alkanes transported from the east Asian continent were isotopically distinct from those of a marine easterly or Indonesian/Australian origin. But further discrimination between Northeast Asian and Southeast Asian sources or between marine easterly and Indonesian/Australian sources was not possible using $\delta^{13}\text{C}$ of the

HMW *n*-alkanes or by various combinations of higher plant biomarker parameters. In general in winter/spring, more air masses transported C₃ material from north Asia, whereas in summer/fall relatively more air masses transported material from south east Asia, Indonesia and Australia where C₄ plants are more abundant. Marine easterlies were also an important source of isotopically heavier *n*-alkanes in the summer, which may have been transported from the Americas. The highest mean C₄ contributions in the summer of 1991 may have been enhanced by strong El Niño conditions which influenced meteorology and biomass burning emissions.

Acknowledgements

We thank S. Ogishima for his assistance in collecting the aerosol sample at the Chichi-Jima site and the National Aeronautic and Science Develop Agency (NASDA) for allowing use the Satellite Tracking Center at the Island and Y. Ishimura for the extraction and isolation of *n*-alkanes from aerosol samples. This study was partly supported by a Grant-in-Aid (10144101) to K.K. from the Japanese Ministry of Education, Science, and Culture. J.B. appreciates the financial support for a Research Fellowship and grant-in-aid from the Japanese Society for the Promotion of Science.

References

- Abas M. R., Simoneit B. R. T., Elias V., Cabral J. A., and Cardoso J. N. (1995) Composition of higher molecular weight organic matter in smoke aerosol from biomass combustion in Amazonia. *Chemosphere* **30**, 995-1015.
- Bird M. I., Summons R. E., Gagan M. K., Roksandic Z., Dowling L., Head J., Fifield L. K., Cresswell R. G., and Johnson D. P. (1995) Terrestrial vegetation change inferred from n-alkane $\delta^{13}\text{C}$ analysis in the marine environment. *Geochimica et Cosmochimica Acta* **59**(13), 2853-2857.
- Blank M., Leinen M., and Prospero J. M. (1985) Major Asian aeolian inputs indicated by the mineralogy of aerosols and sediments in the Western North Pacific. *Nature* **314**, 84-86.
- Blumer M., Guillard R. R. L., and Chase T. (1971) Hydrocarbons of marine phytoplankton. *Marine Biology* **8**, 183.
- Boyer K. K. and Laitinen H. A. (1975) Automobile exhaust particules. *Environ. Sci. Technol.* **9**, 457-459.
- Cerling T. E. and Quade J. (1993) Stable carbon and oxygen isotopes in soil carbonates. In *Climate change in continental isotopic records* (ed. P. K. Swart, K. C. Lohmann, J. McKenzie, and S. Savin). AGU.
- Chesselet R., Fontugne M., Buat-Ménard P., Ezat U., and Lambert C. E. (1981) The origin of particulate organic carbon in the marine atmosphere as indicated by its stable carbon isotopic composition. *Geophysical Research Letters* **8**(4), 345-348.
- Chikaraishi Y. and Naraoka H. (2003) Compound-specific delta D-delta C-13 analyses of n-alkanes extracted from terrestrial and aquatic plants. *Phytochemistry* **63**(3), 361-371.
- Chikaraishi Y., Naraoka H., and Poulson S. R. (2004) Carbon and hydrogen isotopic fractionation during lipid biosynthesis in a higher plant (*Cryptomeria japonica*). *Phytochemistry* **65**(3), 323-330.
- Chung S. H. and Seinfeld J. H. (2002) Global distribution and climate forcing of carbonaceous aerosols. *Journal Of Geophysical Research-Atmospheres* **107**(D19).
- Collister J. W., Rieley G., Stern B., Eglinton G., and Fry B. (1994) Compound-specific $\delta^{13}\text{C}$ analyses of leaf lipids from plants with differing carbon dioxide metabolism. *Organic Geochemistry* **21**(6-7), 619-627.
- Conte M. H. and Weber J. C. (2002) Plant biomarkers in aerosols record isotopic discrimination of terrestrial photosynthesis. *Nature* **417**, 639-641.
- Craig H. (1957) Isotopic standards for carbon and oxygen and correction factors for mass spectrometric analysis of carbon dioxide. *Geochimica et Cosmochimica Acta* **12**, 133-149.
- Cranwell P. A. (1981) Diagenesis of free and bound lipids in terrestrial detritus deposited in a lacustrine sediment. *Organic Geochemistry* **3**(3), 79-89.
- Darwin C. (1846) An account of the fine dust which often falls on vessels in the Atlantic Ocean. *Journal of the Geological Society London* **2**, 26-30.
- Eglinton G., Gonzales A. G., Hamilton R. J., and Raphael R. A. (1962) Hydrocarbon constituents of the wax coatings of plant leaves, a taxonomic survey. *Phytochemistry* **1**(89-102).
- Eglinton G. and Hamilton R. J. (1967) Leaf epicuticular waxes. *Science* **156**, 1322-1335.
- Fang J., Kawamura K., Ishimura Y., and Matsumoto K. (2002) Carbon isotopic composition of fatty acids in the marine aerosols from the western North Pacific: implication for the source and atmospheric transport. *Environmental Science and Technology* **36**, 2598-2604.
- Fontugne M. and Duplessy J. C. (1978) Carbon isotope ratio of marine plankton related to surface water masses. *Earth and Planetary Science Letters* **41**, 365-371.
- Fuelberg H. E., Kiley C. M., Hannan J. R., Westberg D. J., Avery M. A., and Newell R. E. (2003) Meteorological conditions and transport pathways during the Transport and Chemical

- Evolution over the Pacific (TRACE-P) experiment. *Journal Of Geophysical Research Atmospheres*, Research-Atmospheres.
- Gagosian R. B. and Peltzer E. T. (1986) The importance of atmospheric input of terrestrial organic material to deep sea sediments. *Advances in Organic Geochemistry* **10**, 661-669.
- Gagosian R. B., Peltzer E. T., and Merrill J. T. (1987) Long-range transport of terrestrially derived lipids in aerosols from the South Pacific. *Nature* **325**, 800-803.
- Gagosian R. B., Peltzer E. T., and Zafriou O. C. (1981) Atmospheric transport of continentally derived lipids to the tropical North Pacific. *Nature* **291**, 312-314.
- Gelpi E., Schneider J., Mann J., and J. O. (1970) Hydrocarbons of geochemical significance in microscopic algae. *Phytochemistry* **9**, 603-612.
- Huang Y., Street-Perrott F. A., Metcalfe S. E., Brenner M., Moreland M., and Freeman K. H. (2001) Climate change as the dominant control on glacial-interglacial variations in C-3 and C-4 plant abundance. *Science* **293**(5535), 1647-1651.
- Huang Y. S., Dupont L., Sarnthein M., Hayes J. M., and Eglinton G. (2000) Mapping of C-4 plant input from North West Africa into North East Atlantic sediments. *Geochimica et Cosmochimica Acta* **64**(20), 3505-3513.
- Huang Y. S., Street-Perrott F. A., Perrot R. A., Metzger P., and Eglinton G. (1999) Glacial-interglacial environmental changes inferred from molecular and compound-specific $\delta^{13}\text{C}$ analyses of sediments from Sacred Lake, Mt. Kenya. *Geochimica et Cosmochimica Acta* **63**(9), 1383-1404.
- Ikehara M., Kawamura K., Ohkouchi N., Murayama M., Nakamura T., and Taira A. (2000) Variations of terrestrial input and marine productivity in the Southern Ocean (48 – S) during the last two deglaciations. *Paleoceanography* **15**(2), 170-180.
- Ishiwatari R., Uzaki M., and Yamada K. (1994) Carbon isotope composition of individual *n*-alkanes in recent sediments. *Organic Geochemistry* **21**(6/7), 801-808.
- Jayaraman A. (1999) Results in direct radiative forcing of aerosols obtained over the Indian Ocean. *Current Science* **76**, 924-930.
- Kawamura K. (1995) Land-derived lipid class compounds in the deep-sea sediments and marine aerosols from North Pacific. In *Biogeochemical Processes and Ocean Flux in the Western Pacific* (ed. H. Sakai and Y. Nozaki), pp. 31-51. TERRAPUB.
- Kawamura K., Ishimura Y., and Yamazaki K. (2003) Four years' observations of terrestrial lipid class compounds in marine aerosols from the western North Pacific. *Global Biogeochemical Cycles* **17**(1).
- Kawamura K. and Kaplan R. I. (1991) Organic compounds in rainwater. In *Organic Chemistry of the Atmosphere* (ed. L. D. Hansen and D. J. Eatough), pp. 233-284. CRC.
- Kawamura K., Kosaka M., and Sempere R. (1995) Distributions and seasonal changes in hydrocarbons in urban aerosols and rainwaters (in Japanese with an English abstract). *Chiyukagaku (Geochemistry)* **29**, 1-15.
- Kita K., Fujiwara M., and Kawakami S. (2000) Total ozone increase associated with forest fires over the Indonesian region and its relation to the El Nino-Southern oscillation. *Atmospheric Environment* **34**(17), 2681-2690.
- Kolattukudy P. E. (1976) *The Chemistry and Biochemistry of Natural Waxes*. Elsevier.
- Kuypers M. M. M., Pancost R. D., and Sinninghe D. J. S. (1999) A large and abrupt fall in atmospheric CO₂ concentration during Cretaceous times. *Nature* **399**(6734), 342-345.
- Liu H. Y., Jacob D. J., Bey I., Yantosca R. M., Duncan B. N., and Sachse G. W. (2003) Transport pathways for Asian pollution outflow over the Pacific: Interannual and seasonal variations. *Journal Of Geophysical Research Atmospheres*, Research-Atmospheres.
- Lockheart M. J., VanBergen P. F., and Evershed R. P. (1997) Variations in the stable carbon isotope compositions of individual lipids from the leaves of modern angiosperms: Implications for

- the study of higher land plant-derived sedimentary organic matter. *Organic Geochemistry* **26**(1-2), 137-153.
- Matsumoto K., Kawamura K., Uchida M., Shibata Y., and Yoneda M. (2001) Compound specific radiocarbon and $\delta^{13}\text{C}$ measurements of fatty acids in a continental aerosol sample. *Geophysical Research Letters* **28**(24), 4587-4590.
- Novakov T. and Penner J. E. (1993) Large Contribution of Organic Aerosols to Cloud-Condensation-Nuclei Concentrations. *Nature* **365**(6449), 823-826.
- O'Leary. (1981) Carbon isotope fractionation in plants. *Phytochemistry* **20**, 553-568.
- Ohkouchi N., Kawamura K., Kawahata H., and Taira A. (1997a) Latitudinal distributions of terrestrial biomarkers in the sediments from the Central Pacific. *Geochimica et Cosmochimica Acta* **61**(9), 1911-1918.
- Ohkouchi N., Kawamura K., and Taira A. (1997b) Molecular paleoclimatology: reconstruction of climate variabilities in the late Quaternary. *Organic Geochemistry* **27**(3/4), 173-183.
- Ohkouchi N., Kawamura K., Takemoto N., Ikehara M., and Nakatsuka T. (2000) Implications of carbon isotope ratios of C_{27} - C_{33} alkanes and C_{37} alkenes for the sources of organic matter in the Southern Ocean surface sediments. *Geophysical Research Letters* **27**(2), 233-236.
- Oros D. R. and Simoneit B. R. T. (2001) Identification and emission factors of molecular tracers in organic aerosols from biomass burning, 2, Deciduous trees. *Applied Geochemistry* **16**, 1545-1565.
- Penner J. E., Dickinson R., and O'Neill C. (1992) Effects of aerosol from biomass burning on the global radiation budget. *Science* **256**, 1432-1434.
- Prahl F. G. and Muehlhausen L. A. (1989) An organic geochemical assessment of oceanographic conditions at Manop site C over the past 26, 000 years. *Paleoceanography* **4**, 495-510.
- Rex R. W. and Goldberg E. D. (1958) Quartz contents of pelagic sediments of the Pacific Ocean. *Tellus* **10**, 153-159.
- Rieley G., Collister J. W., Stern B., and Eglinton G. (1993) Gas-chromatography/Isotope ratio mass spectrometry of leaf wax *n*-alkanes from plants with differing carbon dioxide metabolisms. *Rap. Comm. Mass. Spectrom.* **7**, 488-491.
- Schefuss E., Ratmeyer V., Stuut J. B. W., Jansen J. H. F., and Damste J. S. S. (2003a) Carbon isotope analyses of *n*-alkanes in dust from the lower atmosphere over the central eastern Atlantic. *Geochimica Et Cosmochimica Acta* **67**(10), 1757-1767.
- Schefuss E., Schouten S., Jansen J. H. F., and Damste J. S. S. (2003b) African vegetation controlled by tropical sea surface temperatures in the mid-Pleistocene period. *Nature* **422**(6930), 418-421.
- Schramm C. T. and Leinen M. S. (1987) Eolian transport to Hole 595A from the late Cretaceous through Cenozoic. *Initial Reports of the Deep Sea Drilling Project* **88**, 469-473.
- Sicre M.-A. and Peltzer E. T. (2004) Lipid geochemistry of remote aerosols from the southwestern Pacific Ocean sector. *Atmospheric Environment* **38**, 1615-1624.
- Simoneit B. R. T. (1984) Organic matter of the troposphere-III. Characterization and sources of petroleum and pyrogenic residues in aerosols over the western United States. *Atmospheric Environment* **18**, 51-67.
- Simoneit B. R. T. (1997) Compound-specific carbon isotope analysis of individual long-chain alkanes and alkanolic acids in Harmattan aerosols. *Atmospheric Environment* **31**(15), 2225-2233.
- Simoneit B. R. T. (2002) Biomass burning - a review of organic tracers for smoke from incomplete combustion. *Applied Geochemistry* **17**, 129-162.
- Simoneit B. R. T., Cardoso J. N., and Robinson N. (1991) An assessment of terrestrial higher molecular weight lipid compounds in aerosol particulate matter over the South Atlantic from about 30-70°S. *Chemosphere* **23**(4), 447-465.
- Simoneit B. R. T., Chester R., and Eglinton G. (1977) Biogenic lipids in particulates from the lower atmosphere over the eastern Atlantic. *Nature* **267**, 682-685.

- Simoneit B. R. T., Cox R. E., and Standley L. J. (1988) Organic matter of the troposphere, IV, Lipids in Harmattan aerosols of Nigeria. *Atmospheric Environment* **22**, 983-1004.
- Simoneit B. R. T. and Eglinton G. (1977) Organic matter of eolian dust and its input to marine sediments. In *Advances in Organic Geochemistry* (ed. R. Campos and J. Goni), pp. 415-430. Enadisma.
- Simoneit B. R. T. and Mazurek M. A. (1981) Air pollution: The organic components. *Crit. Rev. Environ. Control.* **11**, 219-276.
- Smith B. N. and Epstein S. (1971) Two categories of $^{13}\text{C}/^{12}\text{C}$ ratios for higher plants. *Plant Physiology* **47**, 380-384.
- Spicer R. A. (1989) Physiological characteristics of land plants in relation to environment through time. *Trans. Royal Soc. Edinburgh: Earth Sci.* **80**, 321-329.
- Still C. J., Berry J. A., Collatz G. J., and DeFries R. S. (2003) Global distribution of C3 and C4 vegetation: Carbon cycle implications. *Global Biogeochemical Cycles* **17**(1), doi:10.1029/2001GB001807.
- Street-Perrott F. A., Huang Y. S., Perrott R. A., Eglinton G., Barker P., BenKhelifa L., Harkness D. D., and Olago D. O. (1997) Impact of lower atmospheric carbon dioxide on tropical mountain ecosystems. *Science* **278**(5342), 1422-1426.
- Teeri J. A. and Stowe L. G. (1976) Climatic patterns and the distribution of C₄ grasses in North America. *Oecologia* **23**, 1-12.
- Toyoda T. (1981) *Flora of the Bonin Islands*. ABOC-sha.
- UNEP and C⁴. (2002) *The Asian Brown Cloud: Climate and Other Environmental Impacts*. UNEP.
- Verdon D. C., Kiem A. S., and Franks S. W. (2004) Multi-decadal variability of forest fire risk in eastern Australia. *International Journal of Wildland Fire* **13**(2), 165-171.
- Westerhausen L., Poynter J., Eglinton G., Erlenkeuser H., and Sarnthein M. (1993) Marine and terrigenous origin of organic matter in modern sediments of the equatorial East Atlantic: the $\delta^{13}\text{C}$ and molecular record. *Deep-Sea Research I* **40**(5), 1087-1121.
- Woo J. H., Streets D. G., Carmichael G. R., Tang Y. H., Yoo B., Lee W. C., Thongboonchoo N., Pinnock S., Kurata G., Uno I., Fu Q. Y., Vay S., Sachse G. W., Blake D. R., Fried A., and Thornton D. C. (2003) Contribution of biomass and biofuel emissions to trace gas distributions in Asia during the TRACE-P experiment. *Journal Of Geophysical Research*, Art. No. 8812.
- Yu S. (2000) Role of organic acids (formic, acetic, pyruvic and oxalic) in the formation of cloud condensation nuclei (CCN): a review. *Atmospheric Research* **53**(4), 185-217.
- Zhang Z., Zhao M., Yang X., Wang S., Jiang X., Oldfield F., and Eglinton G. (2004) A hydrocarbon biomarker record for the last 40 kyr of plant input to Lake Heqing, southwestern China. *Organic Geochemistry* **35**, 595-613.
- Zhao M., Dupont L., Eglinton G., and Teece M. (2003) n-Alkane and pollen reconstruction of terrestrial climate and vegetation for N.W. Africa over the last 160 kyr. *Organic Geochemistry* **34**(1), 131-143.

Table 1. Summarized data for air temperature (T, °C), total carbon (TC, ug/m³), normalized concentration of *n*-alkanes, carbon preference index (CPI, dimensionless), unresolved carbon mixture (UCM) and ACL.

Season ^a	Number of samples	Air T(°C)	TC (µg/m ³)	normalized concn ^b			CPI ^c	UCM	ACL ^d
				ΣC ₁₉₋₂₆	ΣC ₂₇₋₄₀	Σ <i>n</i> -alkanes			
Winter 90	2	19.9	1.45 ± 0.3	1.77± 0.9	6.07± 2.5	7.84± 3.3	8.4	3.8	29.55
Summer 90	12	26.5	0.47 ± 0.9	5.53± 0.7	19.88± 2.2	25.38± 2.8	3.3	5.8	29.75
Winter 91	5	21.5	0.94 ± 1.2	1.23± 0.2	7.92± 0.5	9.14± 0.6	4.8	4.9	29.50
Summer 91	10	27.2	0.50 ± 1.6	5.86± 0.5	30.33± 3.2	36.20± 3.7	12.4	3.7	30.11
Winter 92	10	19.9	0.78 ± 1.1	4.44± 0.5	29.78± 4.7	34.21± 5.2	3.8	3.8	29.80
Summer 92	9	25.0	0.43 ± 1.7	2.84± 0.4	22.49± 3.2	25.33± 3.6	2.4	5.0	29.70
Winter 93	10	19.0	0.71 ± 0.5	5.56± 0.4	29.56± 2.3	35.04± 2.6	3.8	4.2	29.75
Summer 93	11	26.6	0.40 ± 1.0	3.06± 0.3	22.88± 1.9	25.93± 2.2	6.6	4.0	30.03

^aSummer: May-October; winter: November-April. ^bConcentration of *n*-alkanes normalized to total carbon (TC), ng of *n*-alkanes /µg of C. ^c $CPI_{n-alkanes} = 2 \Sigma \text{ odd } (C_{25} - 35) / [\Sigma \text{ even } (C_{24} - 34) + \Sigma \text{ even } (C_{26} - 36)]$. ^dAverage chain length: the concentrated-weighted mean carbon chain length).

Table 2. Stable carbon isotopic analyses of *n*-alkanes and the calculated corrections for non-higher plant contributions.

Sample	Season	Source Allocation*	$\delta^{13}\text{C}$ values			$\delta^{13}\text{C}_{29}$ corr. ^a	$\delta^{13}\text{C}_{29}$ Δ^b	$\delta^{13}\text{C}_{31}$ corr. ^a	$\delta^{13}\text{C}_{31}$ Δ^b	%C4 (C ₂₉)	%C4 (C ₃₁)
			C ₂₇	C ₂₉	C ₃₁						
QFF-181	Winter 90	NE A	-31.81	-32.62	-33.21	-33.29	-0.67	-33.99	-0.78	17	15
QFF-182		SE A									
QFF-183	Summer 90	ME	-29.45	-30.47	-31.33	-30.66	-0.19	-31.57	-0.24	31	28
QFF-184		ME	-31.21	-32.41	-33.39	-33.01	-0.60	-34.11	-0.73	18	14
QFF-185		ME	-30.14	-30.88	-32.02	-31.35	-0.47	-32.70	-0.68	28	23
QFF-187		ME		-31.34	-32.65	-31.34		-32.65		25	19
QFF-188		ME									
QFF-189		ME		-31.02	-30.48	-31.02		-30.48		27	33
QFF-190		ME		-30.05	-31.07	-30.05		-31.07		34	29
QFF-191		ME									
QFF-192		ME									
QFF-193		ME		-28.63	-28.40	-28.63		-28.40		43	47
QFF-194		ME	-29.16	-30.48	-32.75	-32.29	-1.81	-35.88	-3.13	31	18
QFF-195		ME									
QFF-207	Winter 91	ME	-29.19	-31.75	-30.97	-31.75		-30.97		22	30
QFF-208		NE A	-30.34	-32.04	-32.37	-32.98	-0.95	-33.41	-1.04	21	21
QFF-209		NE A	-32.21	-32.66	-31.80	-33.00	-0.34	-32.08	-0.29	16	25
QFF-210		SE A	-30.24	-30.88	-31.40	-30.94	-0.06	-31.47	-0.07	28	27
QFF-211		SE A	-30.46	-30.73	-31.94	-30.73		-31.94		29	24
QFF-212	Summer 91	ME									
QFF-214		SE A	-28.28	-29.31	-29.15	-30.72	-1.42	-30.49	-1.35	39	42
QFF-216		IA	-28.31	-28.69	-30.00	-29.17	-0.49	-30.74	-0.74	43	36
QFF-217		IA	-30.20	-31.01	-30.94	-33.38	-2.37	-33.25	-2.31	27	30
QFF-218		IA									
QFF-219		IA	-28.08	-27.23	-27.56	-27.23		-27.56		53	53
QFF-220		IA	-29.07	-30.00	-29.95	-30.00		-29.95		34	37
QFF-221		IA	-33.12	-33.38	-34.44	-33.50	-0.12	-34.61	-0.17	12	7
QFF-222		ME	-29.09	-29.48	-29.23	-31.28	-1.80	-30.88	-1.65	38	42
QFF-223		NE A	-31.22	-32.82		-32.82				15	
QFF-393	Winter 92	NE A	-30.92	-32.55	-31.92	-33.36	-0.81	-32.58	-0.67	17	24
QFF-394		NE A									
QFF-395		NE A	-29.27	-33.07	-32.94	-33.44	-0.37	-33.30	-0.36	14	17
QFF-396		NE A	-30.81	-32.88	-32.64	-33.43	-0.55	-33.16	-0.52	15	19
QFF-397		NE A	-32.38	-33.35	-32.71	-33.83	-0.49	-33.13	-0.42	12	19
QFF-399		NE A	-32.02	-33.14	-31.66	-31.51	-0.34	-29.97	-0.28	13	25
QFF-400		SE A	-30.44	-32.83	-31.83	-32.83		-31.83		15	24
QFF-401		SE A	-31.10	-33.05	-32.36	-33.22	-0.17	-32.49	-0.13	14	21
QFF-402		NE A	-31.58	-32.17	-32.84	-32.34	-0.17	-33.07	-0.23	20	18
QFF-403		SE A	-31.43	-33.29	-34.32	-32.21		-32.89		12	8
QFF-404	Summer 92	NE A	-30.22	-30.68	-32.56	-30.68		-32.56		30	19
QFF-405		SE A	-30.04	-33.49	-32.89	-33.49		-32.89		11	17
QFF-406		ME	-29.52	-29.96	-29.32	-30.01	-0.05	-29.36	-0.04	34	41
QFF-407		IA	-31.49	-32.44	-30.94	-33.04	-0.60	-31.36	-0.42	18	30
QFF-408		ME		-29.11	-28.66	-29.11		-28.66		40	45
QFF-410		IA	-29.95	-30.83	-30.51	-30.83		-30.51		29	33
QFF-412		IA	-28.47	-29.80	-30.02	-29.94	-0.14	-30.16	-0.14	36	36
QFF-413		ME		-29.11	-32.10	-29.11		-32.10		40	23
QFF-414		NE A	-30.55	-30.83	-32.68	-31.48	-0.17	-32.90	-0.23	29	19
QFF-415	Winter 93	NE A	-30.35	-31.03	-29.86	-31.03		-29.86		27	37
QFF-416		NE A									
QFF-417		NE A	-32.32	-33.71	-33.21	-33.96	-0.25	-33.42	-0.21	9	15
QFF-418		SE A	-29.54	-31.71	-31.52	-32.18	-0.47	-31.96	-0.45	23	26
QFF-419		NE A	-33.53	-34.26	-33.79	-34.45	-0.19	-33.95	-0.17	6	11
QFF-420		NE A	-29.87	-31.42	-31.65	-33.28	-1.86	-33.59	-1.94	25	25
QFF-421		NE A	-31.47	-33.42	-33.15	-33.42		-33.15		11	16
QFF-422		NE A	-30.87	-32.45	-31.76	-32.71	-0.26	-31.97	-0.22	18	25
QFF-423		NE A	-30.73	-32.12	-32.07	-32.12		-32.07		20	23
QFF-424		NE A	-31.69	-32.87	-33.65	-33.15	-0.29	-34.05	-0.40	15	12
QFF-426	Summer 93	SE A	-29.91	-33.49	-32.23	-33.49		-32.23		11	22
QFF-427		IA	-29.75	-31.55	-32.92	-31.55		-32.92		24	17
QFF-428		IA									
QFF-429		NE A									
QFF-430		ME	-28.64	-28.32	-26.56	-28.32		-26.56		46	59
QFF-431		IA		-30.05	-30.07	-30.05		-30.07		34	36
QFF-432		ME									
QFF-433		IA	-29.40	-28.83	-28.77	-28.83		-28.77		42	45
QFF-434		ME									
QFF-435		ME									
QFF-436		NE A									
QFF-437		NE A	-30.94	-32.57	-32.67	-32.57		-32.67		17	19

* Predominant source area in the 10 day back-trajectory. Acronyms are: NEA = Northeast Asia; SEA = Southeast Asia; Marine Easterlies; IA = Indonesia and Australia

n.d. = not determined, only isotopic data for *n*-alkanes with carbon numbers from C₂₄ to C₃₃ are shown, measurements for other carbon numbers (C₁₉ – C₂₃, C₃₄ – C₃₇) were sparse and have been omitted.

^a $\delta^{13}\text{C}$ values of respective *n*-alkanes corrected for fossil fuel/ marine contribution.

^b Difference in $\delta^{13}\text{C}$ between uncorrected and corrected values for respective *n*-alkanes.

Table 3. Stable carbon isotopic compositions (per mil) of *n*-alkanes from C₃ and C₄ plants.

Plant Type	Source	C ₂₉ δ ¹³ C (s.d.)	C ₃₁ δ ¹³ C (s.d.)
C₃ Plants	Collister et al, 1997. 9 species	-35.4‰ (1.7)	-36.1‰ (1.3)
	Chikaraishi and Naraoka, 2003. 22 species	-35‰ (2.7)	-35.2‰ (2.9)
	ILTS this paper 1 species	-33.7‰	
	Mean	-35.1‰ (1.7)	-35.5‰ (1.3)
C₄ Plants	Collister et al, 1997. 3 species	-21.5‰ (4.3)	-20.2‰ (2.3)
	Chikaraishi and Naraoka, 2003. 7 species	-19.2‰ (2.4)	-20.2‰ (2.3)
	ILTS this paper 1 species	-19.6‰	-20.3‰
	Mean	-20.2‰ (2.3)	-20.4‰ (2)

* Measurements made at ILTS, plant leaves were extracted with DCM/MeOH (5:1) and then the extracts were saponified using 0.5 M KOH/MeOH. The *n*-alkanes were isolated and analysed by the methods described in the text.

Table 4. Calculated % C₄ source input of plants from end members* of C₃ and C₄ plants.

Compound	W-90	S-90	W-91	S-91	W-92	S-92	W-93	S-93
C ₂₉ <i>n</i> -alkane (this paper)	17	30	23	33	15	30	17	29
C ₃₁ <i>n</i> -alkane (this paper)	15	26	25	35	19	29	21	33
C ₂₈ fatty acid Fang <i>et al</i> (2000)	6	20	22	27	9	14	8	18

* For the C₂₈ fatty acid Fang *et al* (2000) used end members of -30.9‰ for C₃ plants and -20.8‰ for C₄ plants.

Figure 1. Global map showing the sampling location (Chichi-Jima) in the western Pacific, the global C4 percentage of terrestrial biomass and the source areas of the four major air mass trajectories. Abbreviations: NE A = North east Asia; SE A = South east Asia; IA = Indonesia and Australia; ME = Marine easterlies. The C4 biomass is adapted from Still et al., 2003, the C4 biomass fraction was modelled by combining remote sensing products, physiological modelling, a spatial distribution of global crop fractions, and national harvest area data for major crop types.

Figure 1: Variations of wind direction at 700 hPa over Chichi-Jima in the western North Pacific (see a map in Figure 1).

Figure 3: Typical distribution of n-alkane concentrations and isotopic values from a North east Asian sample (QFF-417-N1) collected at Chichi-Jima, December 21-25, 1992.

Figure 4: Distribution of carbon isotopic composition ($\delta^{13}\text{C}$, left axis) of C29 and C31 *n*-alkanes and estimates of the percentage contribution of C4 plant material (%C4, right axis) in aerosol samples collected at Chichi-Jima Island between 1990 and 1993. a & b) uncorrected $\delta^{13}\text{C}$ values of C29 and C31. c & d) $\delta^{13}\text{C}$ values of C29 and C31 corrected for non-higher plant inputs (open circles) compared to uncorrected values (black circles), the samples with the largest corrections are highlighted by arrows. Grey shading delimits the summer (May-October) and white the winter seasons (November-April).

Figure 5: Linear correlation between carbon isotopic compositions ($\delta^{13}\text{C}$) of the C29 *n*-alkane (this paper) and the C24 Fatty Acid (Fang *et al*, 2000) in aerosol samples collected at Chichi-Jima Island between 1990 and 1993.

Figure 6: Seasonal mean distributions of carbon isotopic composition ($\delta^{13}\text{C}$) of C29 & C31 *n*-alkanes in aerosol samples collected at Chichi-Jima Island between 1990 and 1993.

Figure 7: Composite seasonal 10 day Back-trajectory analyses for the sampling periods. Conducted using the Fifth-Generation NCAR/Penn State mesocale model (MM5). The trajectory calculations were based on backward tracking of air parcels released at the target location (for the sampling periods) assuming the parcel was moving along the ambient airflow. The flow pattern was updated every 6 hours.

Figure 8: Boxplots of the carbon isotopic composition ($\delta^{13}\text{C}$) of C29 & C31 *n*-alkanes in aerosol samples collected at Chichi-Jima Island between 1990 and 1993, categorised by back-trajectory source area. Each box shows the median (black line), the interquartile range (box), the min and max values that are *not* outliers (whiskers), mild outliers (circles) and extreme outliers (stars).

Figure 9: Data matrix showing plots of all possible combinations of the following variables measured in the aerosols collected at Chichi-Jima: C29 *n*-alkane $\delta^{13}\text{C}$; C31 *n*-alkane $\delta^{13}\text{C}$; C24 FA (fatty acid) $\delta^{13}\text{C}$; CPI; UCM –total *n*-alkanes; ACL (C27-C33). The coloured markers correspond to the source allocations.

Figure 10: Composite seasonal 15 day Back-trajectory analyses for a) QFF-216 and b) QFF-219. Conducted using the Fifth-Generation NCAR/Penn State mesocale model (MM5). The trajectory calculations were based on backward tracking of air

parcels released at the target location (for the sampling periods) assuming the parcel was moving along the ambient airflow. The flow pattern was updated every 6 hours.

Wind direction at 700 hPa over Chichi-Jima

Westerly

Trade wind

Wind direction

QFF-417-N1 21-25st December 1992

1990 Summer trajectories
(10 days backward trajectories)

Winter 1990
(10 days backward trajectories)

Summer 1991
(10 days backward trajectories)

Winter 1991
(10 days backward trajectories)

Summer 1992
(10 days backward trajectories)

Winter 1992
(10 days backward trajectories)

Summer 1993
(10 days backward trajectories)

Winter 1993
(10 days backward trajectories)

a)

QFF-216 June 1991 925-hPa
(15 days backward trajectories)

b)

QFF-219 August 1991 925-hPa
(15 days backward trajectories)