



Title	Differentiative potential of a mouse parthenogenetic embryonic stem cell line revealed by embryoid body formation in vitro
Author(s)	PARK, Jong-Im; YOSHIDA, Ikuya; TADA, Takashi; TAKAGI, Nobuo; TAKAHASHI, Yoshiyuki; KANAGAWA, Hiroshi
Citation	Japanese Journal of Veterinary Research, 46(1), 19-28
Issue Date	1998-05-30
DOI	10.14943/jjvr.46.1.19
Doc URL	<a href="http://hdl.handle.net/2115/2635">http://hdl.handle.net/2115/2635</a>
Type	bulletin (article)
File Information	KJ00002398607.pdf



[Instructions for use](#)

## Differentiative potential of a mouse parthenogenetic embryonic stem cell line revealed by embryoid body formation *in vitro*

Jong-Im Park, Ikuya Yoshida<sup>1</sup>, Takashi Tada<sup>2</sup>, Nobuo Takagi<sup>1</sup>,  
Yoshiyuki Takahashi and Hiroshi Kanagawa.

(Accepted for publication: Jan. 22, 1998)

### Abstract

The *in vitro* differentiative potential of mouse parthenogenetic (PG) embryonic stem (PGES) cells were investigated in the formation of embryoid bodies (EBs). EBs derived from PGES cells retarded in growth and showed restricted differentiation compared to their fertilized counterpart. In chimeric EBs from the aggregation of PGES and fertilized ES cells, morphological examination revealed that PGES cells were reduced in their population and distributed in endodermal layer as culture periods proceeded. These findings were comparable to those in aggregation chimeras of fertilized and PG embryos, and suggest that the differentiation of PGES cells *in vitro* is restricted in the formation of EBs.

Key words: differentiation, embryoid bodies, embryonic stem cells, parthenogenesis,

### Introduction

Mammalian parthenogenetic (PG) embryos do not survive until term<sup>36,38</sup>. In the laboratory mouse, PG embryos are lost at various stages of development; most of them are resorbed immediately, a majority of embryos surviving implantation are lost before 9 days of gestation with no or extremely underdeveloped extraembryonic structures, and even exceptionally well-grown parthenogenones die by midgestation, probably mainly due to poor differentiation of extraembryonic tissues<sup>38</sup>. On the other hand, PG cells are capable of contributing to adult somatic cells as well as germ cells in chimeras produced by aggregation of PG and fertilized embryos<sup>1,12-</sup>

<sup>15,27,39</sup>. PG cells in chimeras distributed randomly early stages after aggregation<sup>7,12</sup>, but they are lost from the trophoblastic cell lineage substantially in early gastrulation stage<sup>7</sup>. Elimination of the PG component from mesodermal tissues such as skeletal muscle became evident later<sup>12-15</sup>. Through midgestation and post natal period, PG cells are restricted to tissues of mostly ectodermal origin<sup>1,21</sup>. It is interesting to notice that cells of androgenone origin are mainly distributed in mesodermal tissues such as skeletal muscles and bones<sup>22,23</sup>. This contrasting contribution of PG and androgenic (AG) cells in chimera may due to genomic imprinting<sup>40,41</sup>. More than twenty endogenous imprinted genes have now been reported in human and mouse<sup>10</sup>,

Laboratory of Theriogenology, Department of Veterinary Clinical Sciences, Graduate School of Veterinary Medicine, Hokkaido University, Sapporo, 060–0818,

<sup>1</sup>) Division of Bioscience, Graduate School of Environmental Earth Science, Hokkaido University, Sapporo, 060–0810,

<sup>2</sup>) School of Medicine, Tottori University, Yonago, 683–0826.

but no candidate gene involved in the differential elimination of AG and PG cells is identified.

Recently, embryonic stem (ES) cells bearing uniparental genomes are frequently used for studies related to imprinting<sup>1,19,21,23,37,42</sup>. ES cells are undifferentiated cultured cell lines established from preimplantation embryos<sup>11,24</sup>. Since ES cells were isolated first in the laboratory mouse more than a decade ago, they have been indispensable tools for dissecting early mammalian embryogenesis<sup>2,4,30</sup>. ES cells have been shown to maintain remarkable pluripotency in chimera formation with host embryos; they can differentiate into functional gametes as well as various types of somatic cells<sup>2,3</sup>. Totipotency of certain ES cell lines has been demonstrated by the birth of mice consisting exclusively of ES derived cells<sup>26,48</sup>. In addition to these unique properties *in vivo*, ES cells undergo spontaneous differentiation under appropriate culture conditions<sup>8,9</sup>. When grown in suspension, ES cells stick one another to form large cell aggregates that differentiate into a structure called embryoid bodies (EBs) closely resembling embryonic region of the egg cylinder stage embryo<sup>2,9</sup>. There are various lines of evidence to indicate that EBs can be used as an *in vitro* model system for studying certain events of occurring in early embryogenesis<sup>5,20,29</sup>.

ES cells are especially valuable for the study for the PG and AG cells because both types of embryos are hard to obtain routinely, although behavior of PG embryonic cells and that of parthenogenetic ES (PGES) cells in chimeric embryos are slightly different<sup>1,37</sup>. It is possible that certain changes in imprinting occurring during establishment and maintenance of ES cell lines are responsible for the differences found between them<sup>25,37,42</sup>. The aim of this study is to understand why PG conceptuses ultimately fail to survive in mammals. This study describes the distribution and behavior of PGES cells in EBs formation by using newly isolated cell line

with *lac Z* gene as *in situ* maker.

## Material and Methods

### Cell lines

$\beta$ -galactosidase-expressing ES cell line NR2 (a kind gift from Dr. M. Azim Surani, Wellcom/CRC, UK.) was derived from a male transgenic mice line bearing *E. coli lac Z* transgene<sup>1</sup>. TMA-24 and TMA-48P cells, which arose from a fertilized male and a PG blastocyst of 129/Sv background, respectively<sup>35</sup>. T48PZ4 cell line was isolated from TMA-48P cells transfected with *lac Z* reporter gene. Briefly, pENL containing *lac Z* gene under the control of elongation factor 1a promoter (a generous gift from Dr. Kazunori Hanaoka, Kitasato University) and pSTneo<sup>18</sup> (a kind gift from Drs. Kazuto Katoh and Hisato Kondoh, Osaka University) as a selection marker were linearized, and introduced into TMA-48P cells by Shimadzu GTE-1 electroporator. After electroporation, cells were grown in a selection medium containing 250  $\mu$ g/ml Geneticin (G418, Sigma) for 10 days. T48PZ4 cells thus isolated was used for this study.

### Culture conditions

ES cells used in this study were maintained in an undifferentiated state on mitomycin C (10  $\mu$ g/ml) treated STO cell feeder layer in Dulbecco's Modified Eagle's Medium (DMEM, Dainippon pharmaceutical Co.) supplemented with 10% fetal bovine serum and  $1 \times 10^3$  unit/ml of recombinant leukemia inhibitory factor (LIF, Amrad) at 37°C, in humidified air containing 5% CO<sub>2</sub><sup>43</sup>. ES cells were grown then in collagen coated plastic petri dishes (Iwaki) without STO cell feeder layer.

### Embryoid bodies preparation

EBs were produced as described before<sup>32,43</sup>. About  $1 \times 10^7$  ES cells grown without feeder layer were trypsinized and plated onto a bacteriological petri dish (Eiken) to which they do not adhere. Chimeric EBs were produced by aggregation of TMA-48P (*lac Z*-negative)/NR2

(*lac Z*-positive), T48PZ4 (*lac Z*-positive)/TMA-24 (*lac Z*-negative) and NR2/TMA-24 cell lines. About  $5 \times 10^6$  cells from each cell line were seeded onto a bacteriological petri dish allowing free aggregation. EBs were maintained in DMEM supplemented with 10% heat-inactivated fetal calf serum without LIF. The medium was changed everyday thereafter. EBs were harvested for histological and morphological examination 3, 5, 7, 10 and 14 days after initiation of differentiation.

#### *Histological examination of embryoid bodies*

To compare the growth of EBs derived from NR2, TMA-48P and aggregation chimera of the two cell lines, EBs were fixed with 4% (w/v) paraformaldehyde in phosphate-buffered saline (0.1 M PBS, containing NaCl 8.0 g/l, KCl 0.2 g/l,  $\text{Na}_2\text{HPO}_4$  1.44 g/l and  $\text{KH}_2\text{PO}_4$  0.24g/l), embedded in paraffin, sectioned serially at  $3 \mu\text{m}$  in thickness, and stained with haematoxylin and eosin. The contribution and distribution of PGES cells in chimeric EBs were determined using X-gal histochemistry<sup>16</sup>). Briefly, chimeric EBs were fixed with 4% (w/v) paraformaldehyde and incubated in X-gal reaction mixture which contained 2 mM  $\text{MgCl}_2$ , 5 mM potassium ferrocyanide, 5 mM potassium ferricyanide, 0.02% NP40 and 0.01% sodium deoxycholate, and 1 mg/ml X-gal in PBS, at  $37^\circ\text{C}$  overnight. After X-gal staining, specimens were dehydrated and embedded in paraffin, sectioned serially at  $3 \mu\text{m}$  in thickness, and stained with haematoxylin and eosin.

#### *Data Analysis*

Numbers of total cells and *lacZ*-positive cells

in each EB were counted using the preparation which gave the maximum area of the EB. Contribution of PGES cells was evaluated by calculating the percentage of the cells derived from PGES cells based on the total cell number. In older EBs, dead cells present at the central region of the section were omitted from counting. Data were compared using Student's *t* test.

## Results

### *Growth of embryoid bodies*

Most EBs derived from NR2 cells tended to be larger than those derived from TMA-48P cells in cell number (Table 1). In fact, the cell number in average-sized EBs in each culture was lower in PGES than fertilized ES cells for the first 3 days in culture ( $P < 0.05$ ). Cell number continued to increase slowly in PGES-derived EBs beyond day 5. Growth of chimeric and fertilized EBs was slightly better than PGES-derived EBs. Dead cells increased in the core region of fertilized and chimeric EBs thereafter, which made it difficult to count the cell number precisely. PGES-derived EBs scarcely developed into a cystic type, and the core region remained more or less homogeneous.

### *Morphology of embryoid bodies*

#### *Fertilized ES cells*

NR2 and TMA-24 cells formed aggregates within 12 hr of suspension culture. Three days after initiation of differentiation, EBs were still simple cell aggregates encircled with endoderm-like cells (Fig. 1, A). By day 5, most of EBs were composed of embryonic ectoderm-like cells and cell debris surrounded by endoderm-like cell

Table 1. Cell numbers of embryoid bodies (EBs) counted at day 3 to day 10

EBs derived from	No. of cells in EBs (No. of EBs examined)			
	Day 3	Day 5	Day 7	Day 10
NR2	151.9 ± 7.5(10) <sup>a)</sup>	208.6 ± 7.6(10) <sup>a)</sup>	382.7 ± 7.3(7) <sup>a)</sup>	469.6 ± 9.1(5) <sup>a)</sup>
TMA-48P	105.7 ± 3.3(10) <sup>b)</sup>	144.8 ± 5.7(10) <sup>b)</sup>	204.4 ± 4.5(6) <sup>b)</sup>	289.4 ± 4.5(7) <sup>b)</sup>
NR2/TMA-48P	156.4 ± 8.4(10) <sup>a)</sup>	282.9 ± 4.9(10) <sup>a)</sup>	293.1 ± 8.0(5) <sup>a)</sup>	459.8 ± 4.1(5) <sup>a)</sup>

Values are mean ± SEM.

a, b) Values with different superscript differ significantly ( $P < 0.05$ ).




Figure 1. Histological sections of embryoid bodies (EBs) at day 3 (A, E and I), day 5 (B, F and J), day 7 (C, G and K), and day 10 (D, H and L). EBs derived from NR2 cell line (A-D) and chimeric EBs derived from TMA-24/T48PZ4 cell lines (I-L) showing well developed endoderm (en) with Reichert's membrane-like material (rm), embryonic ectoderm-like cells (ec), dead cells (dc) and cavity (c) formation. EBs derived from T48PZ4 cell line (E-H) showing flattened embryonic ectoderm-like cells with sparse endodermal layer and poorly organized core. Bar ; 0.5 mm. Haematoxylin and eosin staining.

layer developed fully on the surface of most of EBs (Fig. 1, B). They kept growing, embryonic ectoderm-like cells showed columnar arrangement and an eccentric cavity began to appear in the core. The cavities initiated to expand in most EBs on day 7 (Fig. 1, C). About half of EBs developed balloon-like cysts by day 10 (Fig. 1, D). Under a stereo microscope, beating muscle was first observed in these EBs. Over the next few days, a large proportion of the EBs developed in a similar manner. A few of EBs continued to enlarge and reached several millimeters in diameter. Development ceased at this stage.

#### *Parthenogenetic ES cells*

TMA-48P and T48PZ4 cells completed aggregation within 12 hr as in fertilized ES cell lines. Aggregations were homogeneous on day 3 (Fig. 1, E). Endoderm-like cells began to appear on the surface on day 5, but they developed poorly than fertilized counterparts (Fig. 1, F). Cell death began to be observed in core structure. By day 7, a small cavity appeared in only exceptional EBs. Endoderm-like cells in PG EBs were consistently more sparse in density than those in fertilized EBs of the same age (Fig. 1, G). Extensive cell death disclosed by the accumulation of a large amount of cell




Figure 2. Morphology of chimeric embryoid bodies (EBs) derived from TMA-24 and T48PZ4 cell lines stained with X-gal at day 3 (A), day 5 (B), day 7 (C), and day 10 (D). Note that the distribution of T48PZ cells in endodermal layer (en). Bar ; 0.5 mm (A, B, and C in the same magnification), 0.2 mm (D).

Table 2. Contribution of parthenogenetic ES cells to chimeric embryoid bodies

	% of contribution (No. of EBs examined)			
	3 day	5 day	7 day	10 day
PGES <sup>a)</sup> /FtES <sup>b)</sup>				
T48PZ4/TMA-24	57.8 ± 4.1 (23)	29.3 ± 3.8 (12) <sup>d)</sup>	18.1 ± 3.1 (21) <sup>d)</sup>	4.0 ± 2.1 (46) <sup>d)</sup>
TMA-48P/NR-2	49.5 ± 4.1 (16)	31.4 ± 3.5 (15) <sup>d)</sup>	20.3 ± 2.9 (15) <sup>d)</sup>	6.1 ± 2.6 (10) <sup>d)</sup>
FtES/FtES <sup>c)</sup>				
NR2/TMA-24	48.4 ± 3.6 (10)	49.8 ± 4.1 (10) <sup>e)</sup>	55.2 ± 3.4 (10) <sup>e)</sup>	57.0 ± 4.0 (5) <sup>e)</sup>

Values are mean ± SEM.

a) parthenogenetic ES cells. b) fertilized ES cells. c) contribution of NR2.

d, e) Values with different superscript differ significantly ( $P < 0.05$ )

debris on the bottom of the petri dish should be responsible for the slower growth of PG EBs. At day 10, though EBs were larger in size, no orderly differentiated structure was observed, and only a few exceptional EBs developed into cystic form (Fig. 1, H). After prolonged culture, only a few EBs became cystic, and developed a balloon-like structure, but never showed further differentiation including beating muscles.

#### *Chimeric embryoid bodies*

The size of EBs derived from mixture of PGES cells and fertilized ES cells was slightly smaller than that of fertilized EBs. Overall growth and morphological feature (Fig. 1, I-L) were almost comparable to fertilized EBs, although beating muscle was never observed under stereo microscope in chimeric EBs.

Distribution of parthenogenetic and fertilized ES cells in chimeric embryoid bodies

While degree of chimerism varied among individual EBs, PGES cells decreased steadily throughout the entire culture period (Fig. 2). Table 2 shows the contribution of PGES cells and fertilized ES cells in chimeric EBs. PGES cells were randomly distributed in the EBs 3 days after differentiation. Cell count showed that there was no significant difference in the frequency and distribution of PGES and fertilized ES cells in chimeric EBs at this stage ( $P < 0.01$ ). By day 5, PGES cells had been lost from core structure considerably and remaining PGES scattered in

the core as small colonies of a few cells. Loss of PGES cells from chimeric EBs continued thereafter, but PGES cells remained in the outer endoderm and the ectoderm layer adjacent to it in 7-day EBs.

#### Discussion

EBs from fertilized ES cells mimic certain developmental events occurring in early post-implantation embryos including proamniotic cavity formation and X chromosome inactivation<sup>2,9,35,44</sup>. The present study was initiated to understand the nature of PG cells using this *in vitro* model system. Findings obtained in the present study invariably show that PGES cells are deficient in undergoing differentiation via EB formation. PGES cells were capable of aggregation as fertilized ES cells, but subsequent poor differentiation of endodermal cells on the surface of the EBs and extensive cell death seemed major factors responsible for the defective EBs formation. Reduced differentiative potential into endoderm in PGES cells might reflect the abnormal endoderm differentiation in PG embryos *in vivo* reported by Sturm *et al.*<sup>38</sup>. Recent findings point to important inductive roles played by the visceral endoderm in implanting embryos<sup>6,17,29</sup>. It is possible that defects in endoderm formation is a crucial role trait that explains, at least partly, the abnormal development of PG embryos.

Contrary to the above findings, PGES cells persisted in the outer endodermal layer in

chimeric EBs consisting of fertilized ES and PGES cells. The cellular position in the cell aggregate is an important factor for determining the fate of each cell<sup>34</sup>). Thus, outside cells at the morula stage are destined to differentiate into trophectoderm cells, and inside cells remain undifferentiated as inner cell mass cells<sup>31</sup>). In the case of the EBs, it may be assumed that outside cells transform into endoderm cells. In view of the reduced differentiative potential into endoderm, various assumptions have to be made elucidate the prevalence of PGES cells in the endodermal layer of chimeric EBs. Most probably, PGES cells were forced to take the outside position due to slight differences in the cell surface structure between the two ES cell lines comprising the EBs. This consistent with finding reported by Rosenstraus and Levine<sup>33</sup>) in EBs formed by nullipotent and pluripotent embryonal carcinoma (EC) cell lines. In this case, nullipotent cells were allocated at the center and pluripotent cells were allocated to outer region to form the endodermal layer. Interestingly, there was apparent correlation between the allocation of PGES cells in EBs and their size in the present study (data was not shown). Chimeric EBs tended to be smaller if PGES cells were present in the inner ectodermal portion. While they were bigger than fertilized EBs in size if PGES cells were present mainly in the outer endoderm layer. It was reported that EBs that developed into cystic stage had the outer endodermal layer composed mainly of visceral endoderm-like cells<sup>9</sup>). An intriguing possibility would be that the coexisting fertilized ES cells complemented the defective differentiative potential of the PGES cells into endoderm.

Another remarkable feature of EBs formed by PGES cells was inability of cavitation, and hence formation of neuroepithelial tissue lining the cavity, which may be attributed to either lack of inductive signal from the defective endoderm or failure of response by inner cells to exogenous

signals. Previously Sado *et al.*<sup>35</sup>) obtained evidence to show that fertilized ES cells helped differentiation of PGES cells and female embryonic germ (EG) cells in chimeric EBs. The frequency of PGES cells and female EG cells having an inactive X chromosome increased significantly in differentiating chimeric EBs than in EBs formed by either PGES cells or female EG cells alone. In agreement with this finding Coucouvanis and Martin<sup>8</sup>) succeeded in inducing cavitation at the center of homogenous aggregates formed by nullipotent EC cells by enveloping with visceral endoderm layer. They proposed that the cavitation in early embryos and EBs is the result of the interaction of two signals, the dead signal from the outer endodermal layer and the survival signal from the basement membrane.

Gradual but consistent decrease of PGES cells from chimeric EBs hindered pursuit of the initial aim of the present study. This phenomenon resembles the exclusion of PG cells from PG/fertilized chimeric embryos *in vivo*<sup>12-15,26</sup>), which could be elucidated by a low mitotic activity and/or precocious cell death in PG cells<sup>28,44,46</sup>). PG embryos are probably short of certain growth factors such as insulin-like growth factor-II coded by maternally imprinted gene at specific stages of development and EB formation<sup>28,47</sup>). However, LT-1 EC cell line established from PG ovarian teratocarcinoma differentiated into balloon-like cystic EBs<sup>45</sup>). It is likely that imprinting is more stable in ES cell lines than in EC cell lines.

Although *in vitro* model system have various limitations as shown in this study, further careful studies of differentiating PG and AG ES cells may throw a new light on the role of genomic imprinting in early mammalian embryogenesis.

## References

- 1) Allen, N. D., Barton, S. C., Hilton, K., Norris, M. L., and Surani, M. A. 1994. A functional analysis of imprinting in parth-

- enogenetic embryonic stem cells. *Development*, 120 : 1473–1482.
- 2) Bradley, A. 1990. Embryonic stem cells: proliferation and differentiation. *Curr. Opin. Cell Biol.*, 2 : 1013–1017.
  - 3) Bradley, A., Evans, M., Kaufman, M. H. and Robertson, E. 1984. Formation of germ-line chimaeras from embryo-derived teratocarcinoma cell lines. *Nature*, 309 : 255–256.
  - 4) Camper, S. A., Saunders, T. L., Kendall, S. K., Keri, R. A., Seasholtz, A. F., Gordon, D., Birkmeier, T. S., Keegan, C. E., Karolyi, I. J., Roller, M. L., Burrows, H. L. and Samuelson, L. C. 1995. Implementing transgenic and embryo stem cell technology to study gene expression, cell-cell interactions and gene function. *Biol. Reprod.*, 52 : 246–257.
  - 5) Chen, U., Kosco, M. and Kosco, M. 1992. Establishment and characterization of lymphoid and myeloid mixed-cell population from mouse late embryoid bodies, “embryonic-stem-cell fetuses”. *Proc. Natl. Acad. Sci. USA.*, 89 : 2541–2545.
  - 6) Chen, W. S., Manova, K., Weinstein, D. C., Duncan, S. A., Plump, A. S., Prezioso, V. R., Bachvarova, R. F. and Darnell, J. E., Jr. 1994. Disruption of the HNF-4 gene, expressed in visceral endoderm, leads to cell death in embryonic ectoderm and impaired gastrulation of mouse embryos. *Genes Develop.*, 8 ; 2466–2477.
  - 7) Clarke, H., J. Varmuza, S., Prideaux, V. R. and Rossant, J. 1988. The developmental potential of parthenogenetically derived cells in chimeric mouse embryos: implications for action of imprinted genes. *Development*, 104 : 175–182
  - 8) Coucouvanis, E. and Martin, G. R. 1995. Signals for death and survival: A two-step mechanism for cavitation in the vertebrate embryo. *Cell*, 83 : 279–287.
  - 9) Doetschman, T. C., Eistetter, H., Katz, M., Schmidt, W. and Kemler, R. 1985. The in vitro development of blastocyst-derived embryonic stem cell lines: . formation of visceral yolk sac, blood islands and myocardium. *J. Embryo. exp. Morph.*, 87 : 27–45.
  - 10) Efstratiadis, A. 1994. Parental imprinting of autosomal mammalian genes. *Curr. Opin. Genes Dev.*, 4 : 265–280.
  - 11) Evans, M. J. and Kaufman, M. H. 1981. Establishment in culture of pluripotential cells from mouse embryos. *Nature*, 292 : 154–156.
  - 12) Fundele, R., Bober, E., Arnold, H. H., Grim, M., Bender, R., Wilting, J. and Christ, B. 1994. Early skeletal muscle development proceeds normally in parthenogenetic mouse embryos. *Dev. Biol.*, 161 : 30–36.
  - 13) Fundele, R., Howlett, S. K., Kothary, R., Norris, M. L., Mills, W. E. and Surani, M. A. 1991. Developmental potential of parthenogenetic cells: role of genotype-specific modifiers. *Development*, 113 : 941–946.
  - 14) Fundele, R. H., Norris, M. L., Barton, S. C., Fehlau, M., Howlett, S. K., Mills, W. E. and Surani, M. A. 1990. Temporal and spatial selection against parthenogenetic cells during development of fetal chimeras. *Development*, 108 : 203–211.
  - 15) Fundele, R. H., Norris, M. L., Barton, S. C., Reik, W. and Surani M. A. 1989. Systemic elimination of parthenogenetic cells in mouse chimeras. *Development*, 106 : 29–35.
  - 16) Hogan, B., Beddington, R., Costantini, F. and Lacy, E. 1994. *Manipulating the mouse embryo: a laboratory manual*. 2nd ed., pp. 340–343, Cold Spring Harbor Laboratory press, Cold Spring Harbor, New York
  - 17) Hogan, B. L. M. and Tilly, R. 1981. Cell interactions and endoderm differentiation in cultured mouse embryos. *J. Embryol. exp. Morph.*, 62 : 379–394.
  - 18) Katoh, K., Takahashi, Y., Hayashi, S. and Kondoh, H. 1987. Improved mammalian vectors for high expression of G418 resistance. *Cell Struct. Funct.*, 12 : 575–580.
  - 19) Kaufman, M. H., Robertson, E. J., Handyside, A. H. and M. J. Evans. 1983. Establishment of pluripotential cell lines from haploid mouse embryos. *J. Embryol. exp. Morph.*, 73 : 249–261.
  - 20) Keller, G. M. 1995. *In vitro* differentiation of embryonic stem cells. *Curr. Opin. Cell Biol.*, 7 : 862–869.
  - 21) Keverne, E. B., Fundele, R., Narasimba, M.,

- Barton, S. C. and Surani, M. A. 1996. Genomic imprinting and the differential roles of parental genomes in brain development. *Brain Res. Dev. Brain Res.*, 92 : 91–100.
- 22) Mann, J. R. 1992. Properties of androgenetic and parthenogenetic mouse embryonic stem cell lines; are genetic imprints conserved? *Seminars in Dev. Biol.*, 3 : 77–85.
- 23) Mann, J. R., Gadi, I., Harbison, M. L., Abbondanzo, S. J. and Stewart, C. L. 1990. Androgenetic mouse embryonic stem cells are pluripotent and cause skeletal defects in chimeras: Implications for genetic imprinting. *Cell*, 62 : 251–260.
- 24) Martin, G. R. 1981. Isolation of a pluripotent cell line from early mouse embryos cultured in medium conditioned by teratocarcinoma stem cells. *Proc. Natl. Acad. Sci. USA.*, 78 : 7634–7638.
- 25) McKarney, L. A., Overall, M. L. and Dziadek, M. 1996. Expression of H19 and Igf2 genes in uniparental mouse ES cells during *in vitro* and *in vivo* differentiation. *Differentiation*, 60 : 75–86.
- 26) Nagy, A., Gocza, E., Diaz, E. M., Prideaux, V. R., Ivanyi, E., Markkula, M. and Rossant, J. 1990. Embryonic stem cells alone are able to support fetal development in the mouse. *Development*, 110 : 815–821.
- 27) Nagy, A., Sass, M. and Markkula, M. 1989. Systemic non-uniform distribution of parthenogenetic cells in adult mouse chimaeras. *Development*, 106 : 321–324.
- 28) Newman-Smith, E. D. and Werb, Z. 1995. Stem cell defects in parthenogenetic preimplantation embryos. *Development*, 121 : 2069–2077.
- 29) Palis, J., McGrath, K. E. and Kingsley, P. D. 1995. Initiation of hematopoiesis and vasculogenesis in murine yolk sac explants. *Blood*, 86 : 156–163.
- 30) Pedersen, R. A. 1994. Studies of *in vitro* differentiation with embryonic stem cells. *Reprod. Fertil. Dev.*, 6 : 543–552.
- 31) Pedersen, R. A., Wu, K. and Balakier, H. 1986. Origin of the inner cell mass in mouse embryos: Cell lineage analysis by microinjection. *Dev. Biol.*, 117 : 581–595.
- 32) Robertson, E. J. 1987. Embryo-derived stem cell lines. In: *Teratocarcinomas and embryonic stem cells: a Practical Approach*, pp. 71–112, Robertson, E. J. ed., IRL Press, Oxford.
- 33) Rosenstraus, M. J. and Levine, A. J. 1979. Alterations in the developmental potential of embryonal carcinoma cells in mixed aggregates of nullipotent and pluripotent cells. *Cell*, 17 : 337–346.
- 34) Rosenstraus, M. J. and Spadaro, J. P. and Nilsson, J. 1983. Cell position regulates endodermal differentiation in embryonal carcinoma cell aggregates. *Dev. Biol.*, 98 : 110–116.
- 35) Sado, T., Tada, T. and Takagi, N. 1996. Mosaic methylation of Xist gene before chromosome inactivation in undifferentiated female mouse embryonic stem and embryonic germ cells. *Dev. Dyn.*, 205 : 421–434.
- 36) Solter, D. 1988. Differential imprinting and expression of maternal and paternal genomes. *Annu. Rev. Genet.*, 22 : 127–146.
- 37) Sturm, K. S., Berger, C. N., Zhou, S. X., Dunwoodie, S. L., Tan S.-S. and Tam, P. P. L. 1997. Unrestricted lineage differentiation of parthenogenetic ES cells. *Dev. Genes. Evol.*, 206 : 377–388.
- 38) Sturm, K. S., Flannery, M. L. and Pedersen, R. A. 1994. Abnormal development of embryonic and extraembryonic cell lineages in parthenogenetic mouse embryos. *Dev. Dyn.*, 201 : 11–28.
- 39) Surani, M. A. H. and Barton, S. C. 1977. Development to term of chimaeras between diploid parthenogenetic and fertilized embryos. *Nature*, 270 : 601–603.
- 40) Surani, M. A. H., Barton, S. C. and Norris, M. L. 1987. Experimental reconstruction of mouse eggs and embryos: An analysis of mammalian development. *Biol. Reprod.*, 36 : 1–16.
- 41) Surani, M. A., Kothary, R., Allen, N. D., Singh, P. B., Fundele, R., Ferguson-Smith, A. C. and Barton, S. C. 1990. Genomic imprinting and development in the mouse. *Development*, Suppl. 89–98.
- 42) Szabo, P. and Mann, J. R. 1994. Expression and methylation of imprinted genes during *in vitro* differentiation of mouse parthenogenetic

- and androgenetic embryonic stem cell lines. *Development*, 120 : 1651–1660.
- 43) Tada, T., Tada, M. and Takagi, N. 1993. X chromosome retains the memory of its parental origin in murine embryonic stem cells. *Development*, 119 : 813–821.
- 44) Tada, T. and Takagi, N. 1992. Early development and X-chromosome inactivation in mouse parthenogenetic embryos. *Mol. Reprod. Dev.*, 31 : 20–27.
- 45) Takagi, N. and Martin, G. R. 1984. Studies of the temporal relationship between the cytogenetic and biochemical manifestations of X-chromosome inactivation during the differentiation of LT-1 teratocarcinoma stem cells. *Dev. Biol.*, 103 : 425–433.
- 46) Uranga, J. A. and Arechaga, J. 1997. Cell proliferation is reduced in parthenogenetic mouse embryos at the blastocyst stage : A quantitative study. *Anat. Rec.*, 247 : 243–247.
- 47) Walsh, C., Glaser, A., Fundele, R., Ferguson-Smith, A., Barton, S., Surani, M. A., Ohlsson, R. 1994. The non-viability of uniparental mouse conceptuses correlates with the loss of the products of imprinted genes. *Mech. Dev.*, 46 : 55–62.
- 48) Wood, S. A., Pascoe, W. S., Schmidt, C., Kemler, R., Evans, M. J. and Allen, N. D. 1993. Simple and efficient production of embryonic stem cell-embryo chimeras by coculture. *Proc. Natl. Acad. Sci. USA.*, 90 : 4582–4585.