

Title         Effects of Some Chemicals on the Facet Number Variation in Bar-Eyed Mutant of Drosophila melanogaster					
Author(s)	KAWABE, Masata				
Citation	北海道大學理學部紀要, 13(1-4), 243-247				
Issue Date	1957-08				
Doc URL	http://hdl.handle.net/2115/27235				
Туре	bulletin (article)				
File Information	13(1_4)_P243-247.pdf				



# Effects of Some Chemicals on the Facet Number Variation in Bar-Eyed Mutant of Drosophila melanogaster<sup>1)</sup>

By

## Masata Kawabe

(Biological Institute, Kôbe University)

Sevster found in 1911 that the facet number of the Bar-eyed mutant in Drosophila melanogaster varied under the influence of temperature. Then the Bar-eved mutant was used by several investigators as a valuable tool in the study of physiological genetics. In addition to the temperature, numerous factors were found to modify the phenotype of the Bar-eyed mutant : they are X-rays (Luce, Quastler & Chase '51), oxygen (Margolis '39), nutrition (Krafka '20), extract of insect (Ephrussi, Khouvine & Chevais '38), and various chemicals (Hinshow '41, and Khouvine, Chevais & Grégoiré '43). Further it was found that the chemicals such as amino acids, imidazol derivatives, ureids, and acid amides strongly increased the facet number of Bar-eyes, and that in some cases they converted the phenotype of Bar-eves into the wild or the heterozygotic type (Kaji '54). On the other hand, there are reports that the facet number of Bar-eye is reduced through the treatment of formalin (Hinshow '41, Luce '41). It is of interest to investigate whether any chemicals other than formalin influence or not to reduce the facet number of Bar-eye from the viewpoint of physiological genetics. In this paper, the author wishes to report the results of some preliminary experiments undertaken from the above viewpoint.

#### Material and methods

The Bar and Ultra-Bar strains of Drosophila melanogaster employed as material in the present study were derived from the stocks maintained at the Kyôto University. A few pairs of B or BB flies were cultured in the glass tube, 14 cm in length, 2.5 cm in diameter, containing 5 cc of the culture medium with living yeast. Chemicals which are to be examined were added in definite concentration to the ordinary culture medium. Percentages shown in Tables 1 and 2 denote the rate of chemicals to the culture medium, which consists of white flour 65 g, sugar 25 g, agar 7 g, and water 550 cc. In oxyquinolline, scarlet and Sudan III, the concentrations were indicated by cc of the alcoholic saturated solutions in the culture medium, and then the letter S is used to designate the con-

<sup>1)</sup> Contribution No. 45 from the Biological Institute, Faculty of Science, Kôbe University, Kôbe, Japan.

Dedicated to Professor Tohru Uchida in honor of his 60th birthday.

Jour, Fac. Sci. Hokkaido Univ. Ser. VI, Zool. 13, 1957 (Prof. T. Uchida Jubilee Volume).

#### M. Kawabe

centration in Tables 1 and 2.

In each experiment, the culture medium without chemicals was prepared to use as a control. Throughout the experiments the temperature was kept at  $25^{\circ}$  to  $28^{\circ}$ C. The facet number of the left eye was counted in the next progeny. In some exceptional cases the facet number of the right eye was counted. Flies for observation were sampled randomly from early hatched flies.

	Chemicals	Concen- tration	Obs. no.	ې و Facet number MinMax. Mean			Obs. no.	් ඊ Facet number MinMax. Mean		
а	Formalin	{Control 0.08%	10 20	34 22	45 35	39.2 28.4*	10 20	40 26	69 49	53.6 38.5*
b	Potassium cyanide	{Control 0.04%	10 20	34 34	53 53	41.2 41.7	10 16	43 32	66 69	52.6 43.7
с	Thiosulphate sodium	{Control 0.05%	10 12	29 44	78 75	49.6 53.3	10 5	37 41	74 76	60.5 62.6
d	Dinitrophenol	Control 0.00002M 0.0001 M 0.0002 M	9 9 10 20	32 31 23 19	47 49 39 40	40.2 40.4 30.3* 30.9*	10 3 10 19	35 43 38 25	61 51 72 75	45.2 45.6 46.5 49.7
e	Cresol	Control	42	23	57	38.6	38	44	79	57.2
		$Ort- \begin{cases} 1\% \\ 2\% \\ 3\% \\ 4\% \\ 5\% \end{cases}$	5 14 35 39 18	26 34 20 27 28	45 58 49 63 43	38.6 43.0 32.7 35.7 33.1	2 12 14 25 21	59 47 44 33 41	77 74 68 68 62	68.0 59.4 53.2 45.5 48.3
		$\left\{ \begin{array}{c} 1\% \\ 2\% \\ 3\% \\ 4\% \\ 5\% \end{array} \right\}$	9 18 10 9 24	24 22 34 24 24	42 50 54 44 38	31.5 31.5 44.5 32.8 30.8	7 24 10 4 15	32 39 39 36 37	70 72 66 48 56	47.3 58.8 52.9 41.6 44.4
		$   Para -    \begin{cases}     1\% \\     2\% \\     3\% \\     4\% \\     5\%   \end{cases} $	13 15 5 30 25	26 23 37 25 25	46 40 46 41 44	36.0 29.4 42.0 33.6 35.6	11 13 7 30 48	40 31 42 36 37	75 67 85 77 63	59.8 51.8 61.3 47.5 51.3
f	Oxyquinolline	$ \left\{ \begin{array}{c} I \\ I \\ \\ \\ \\ \\ \\ \\ \\ \\ \\ \\ \\ \\ \\ \\ \\ $	10 10 10	34 34 30 35	53 50 48 53	41.2 40.3 37.2 42.5	10 10 10 5	43 33 25 38	66 61 74 62	52.6 50.8 40.2 51.8
g	Piridin	$ \begin{array}{c cccc}  & 11 & 1. & 5 \\  & & & \\  & & $	7 3 10 7	28 34 35 42	31 40 56 57	29.7 36.3 43.3 47.7	5 6 10 5	33 34 41 42	70 65 81 63	40.4 51.8 57.1 51.4

Table 1. Facet numbers of Ultra-Bar flies fed on various chemicals.

\* means that difference from its control is statistically significant with more 95% of confidence.

#### 244

## **Results of experiments**

Effects on Ultra-Bar flies: Examination of reducing effect of formalin on Ultra-Bar was made in a 0.08% solution using the BB strain. The results obtained are shown in Table 1, a. The facet number of BB flies fed on 0.08% formalin are reduced significantly in comparison with the control flies.

	Chemicals	Concen- tration	Obs. no.	çç Facet number MinMax. Mean			Obs. no.	<sup>8</sup> <sup>8</sup> Facet number MinMax. Mean		
a	Phenol	$\begin{cases} Control \\ 0.1\% \\ 0.2\% \end{cases}$	17 15 6	25 35 36	79 67 62	63.2 45.9 49.5	17 10 8	45 64 64	105 102 76	75.2 82.7 70.2
b	Hydroquinnone	$ \begin{cases} Control \\ 0.2\% \end{cases} $	17 15	25 49	79 68	63.2 54.6	17 8	<b>45</b> 60	105 73	75.2 68.4
c ,	Phloroglucinol	Control 0.1% 0.2%	17 3 6	25 51 30	79 56 46	63.2 53.3 36.3*	17 7 12	45 64 49	105 89 78	75.2 73.2 60.6
d	Ethanol	Control 19%	10 10	38 32	74 47	58.0 39.4*	5 10	54 43	115 111	73.2 62.5
e	Boric acid	{Control 0.01%	20 2	46 32	103 43	65.1 37.5	19 11	50 44	92 77	72.7 60.2
f	Oxyquinolline	{Control 1S	4 7	56 35	87 52	66.5 44.0	4 1	75	110	92.5 50.0
g	Acriflavine	$ \int_{0}^{I} \begin{cases} Control \\ 0 & 01\% \end{cases} $	9 10	53 39	87 50	64.4 42.7*	6 11	60 35	85 66	74.5 50.3*
		$II \left\{ \begin{smallmatrix} \mathrm{Control} & 0.08\% \\ 0.08\% \end{smallmatrix} \right\}$	20 12	46 28	103 48	65.1 41.6*	19 8	50 53	92 77	72.7 66.2*
h	Butter yellow	{Control 0.02% 0.05%	20 10 11	46 39 27	103 80 46	65.1 55.7 33.8*	19 10 9	50 51 39	92 98 60	72.7 75.1 47.6*
ì	Scarlet	$ \left\{ \begin{array}{c} I \\ \\ 0.01S \\ 0.03S \\ 0.05S \\ \end{array} \right. $	9 4 3 10	53 35 37 27 38	87 39 42 45 74	64.4 37.0 39.3 33.2* 58.0	6 11 10 10	60 51 44 26 54	85 96 87 55	74.5 70.2 60.7 45.4*
			17 10	41 27	65 45	51.5* 33.2	7 8	75 41	116 94	87.4 56.2*
j	Sudan	$III \begin{cases} Control \\ 0.03S \\ 0.05S \\ 0.1 & S \end{cases}$	20 7 10 5	46 58 44 33	103 75 61 42	65.1 66.4 52.7 38.1*	19 5 9 3	50 65 58 44	92 75 80 65	72.7 70.0 70.3 55.6*

Table 2. Facet number of Bar flies fed on various chemicals.

\* means that difference from its control is statistically significant with more 95% of confidence.

#### M. Kawabe

Potassium cyanide and 2-4, dinitrophenol, both of which were commonly known as inhibitors of oxidative enzymes, were used to observe reducing effect of the facet number of BB flies. The reducing effect on the facet number was not clearly indicated by the treatment with 0.04% potassium cyanide (Table 1, b), but the facet numbers of female BB flies were reduced significantly in 0.0002 M and 0.0001 M of dinitrophenol (Table 1, d).

Thiosulphate sodium, ort-, meta-, and para-cresol, oxyquinolline, and piridin were examined to learn the reducing effect on the facet number of BB flies. The results obtained here are shown in Table 1, c,e,f, and g. They gave no clear effect of reduction on facet number, except oxyquinolline. Through the treatment with oxyquinolline (1 S), the facet number of BB flies was reduced in comparison with the results of the controls in both of I and II experiments, though not statistically significant.

Effects on Bar flies: Since the reducing effect of dinitrophenol on the facet number of BB flies was seen, phenols (such as phenol, hydroquinnon and phloro-glucinol), ethyl alcohol, and oxyquinollin were examined to see a similar effect on Bar flies. The results of these experiments are as shown in Table 2, a to e, 0.2% phloroglucinol and 19% ethanol showed a slight reducing effect, especially in female flies. No clear effects were given by treatment with other chemicals.

Boric acid and carcinogenic dyes such as acriflavin, butter yellow, scarlet and Sudan III come under examination for reducing effect on the facet number of *Bar* flies. From the results of these experiments as indiacted in Table 2, f to j, it is possible to say that all of carcinogenic chemicals give a remarkable reducing effect on the facet number of *Bar* flies, with exception of boric acid, since the flies examined with the latter chemical are too small in number. A similar reducing effects were shown in the repeated experiments of the examination with acriflavine and scarlet.

#### **Remarks and conclusion**

Reviewing the data presented in this study, two points call attention for discussion. The first deals with a comparison between the action of so-called facet increasing substances (B<sup>+</sup>-substances) and those of chemicals examined here. B<sup>+</sup>substances act more intensely on the male than on the female (Kaji '51). On the contrary, the reducing effect was more manifest in the female than in the male. The second point is a fact that carcinogenic dyes show a reducing act on the facet number of the Bar-eyed mutant like nitrogen mustard. Kaji and Ogaki ('51) found that phenocopies of eyeless or Bar-eyed mutants in the wild flies were induced by the treatment of nitrogen mustard.

It seems probable from the preliminarly experiments reported here that carcinogenic dyes such as acriflavine, scarlet and Sudan III, and some kinds of phenols such as dinitrophenol and phloroglucinol have a similar reducing effect as formalin on the facet number of Bar-eyed mutant, though the effects were not very remark-

246

able, in comparison with the effects of B+-substances such as acid amides. Since these chemicals affect indirectly the process of Bar-eye formation, it is necessary to find chemicals with more direct action on the reducing facet number of Bar-eyed mutant in the future.

Acknowledgment: I am much indebted to Professor S. Fujii, Kôbe University, for his invaluable criticisms and encouragement throughout the work. I wish to express also my sincere gratitude to Dr. Sajiro Makino, Professor of Hokkaido University, for his revision of this manuscript. Thanks are due to Assistant professor Y. Okuda who kindly place his unpublished data regarding cresol and piridin at my disposal. This work was aided in part by a grant from the Scientific Research Fund of the Department of Education.

### Literature cited

Ephrussi, B., Y. Khouvine and S. Chevais 1938. Nature 141: 204.

Hinshow, B. M. 1941. Trans. Ill. State Acad. Sci. 34: 223-224.

Kaji, S. 1954. Annot. Zool. Jap. 26: 224-227.

Kaji, S. and M. Ogaki 1951. Zool. Mag. (Tokyo) 60: 254-257.

Khouvine, Y., S. Cheavis et J. Grégoiré 1943. C. R. Acad. Sci. (Paris) 217 : 161-163.

Krafka, J. J. 1920. Jour. Gen. Physiol. 3: 207-210.

Luce, W. M. 1941. Rec. Genet. Soc. Amer. 7: 79.

Luce, W. M., H. Quastler, and H. B. Chase 1951. Genetics 36: 488-499.

Margolis, O. S. 1939. Genetics 24: 15-21.

Seyster, E. W. 1911. Biol. Bull. 37: 168-181.