



Title	Parasitic forms of a myxosporean in the kidney of <i>Lampetra japonica</i> : An ultrastructural study
Author(s)	MORI, Koshi
Citation	Japanese Journal of Veterinary Research, 47(1-2): 65-65
Issue Date	1999-08-31
Doc URL	<a href="http://hdl.handle.net/2115/2745">http://hdl.handle.net/2115/2745</a>
Type	bulletin
File Information	KJ00003408077.pdf



[Instructions for use](#)

Parasitic forms of a myxosporean in the kidney of *Lampetra japonica*  
: An ultrastructural study

Koshi Mori

*Laboratory of Anatomy,  
Department of Biomedical Sciences,  
School of Veterinary Medicine,  
Hokkaido University, Sapporo 060-0818, Japan*

No information have been available on myxosporea in cyclostomata. The present study was originally planned to investigate "macrophagic cells" in the kidney of arctic lamprey *Lampetra japonica*. Subsequent ultrastructural study revealed that the "macrophagic cells" were unidentified myxosporean trophozoites parasitized at high prevalences in the fish kidney. The trophozoites (pseudoplasmodia with/without sporoblasts) existed predominantly in the lumen of proximal urinary tubules, but were rarely found in any other regions of the kidney. Since no mature spores were found, exact identification of species was impossible. Two types of parasitic forms were observed in proximal urinary tubules: one attaching to the epithelial cells of

renal tubules, and the other free-floating in the lumen of tubules. Ultrastructurally, the attaching trophozoites developed microvilli-like projections towards the apical surface of epithelial cells and interdigitated with microvilli of the brush border. In contrast, the whole surface of floating trophozoites was smooth without any cell projections. The developed projections of trophozoites may contribute to their firm attachment to the epithelial cells and/or absorption of nutrients via the epithelial cells. Against the myxosporean infection, the lamprey as the host provoked a local immune reaction by disposition of numerous lymphocytes and macrophages into the epithelium of urinary tubules.

Contribution of dihydropyridine-insensitive mechanisms to  $\text{Ca}^{2+}$  influx  
in rat pancreatic  $\beta$  cells

Mikio Hayashi

*Laboratory of Physiology, Department of Biomedical Sciences,  
Faculty of Veterinary Medicine, Hokkaido University,  
Sapporo 060-0818, Japan*

1. The present study was carried out to clarify the role of  $\text{Ca}^{2+}$  influx through dihydropyridine-insensitive mechanisms in stimulus-secretion coupling in rat pancreatic  $\beta$  cells. To characterize the dihydropyridine-insensitive  $\text{Ca}^{2+}$  influx, intracellular  $\text{Ca}^{2+}$  concentration ( $[\text{Ca}^{2+}]_i$ ) was directly monitored by a microfluorometric method using Fura-2 in isolated perfused prepa-

rations of rat pancreatic islets. The changes in  $[\text{Ca}^{2+}]_i$  induced by high concentrations of glucose and potassium ion ( $\text{K}^+$ ) were analyzed quantitatively, and influence of  $\text{Ca}^{2+}$  entry blockers upon  $[\text{Ca}^{2+}]_i$  changes was examined.

2. Stimulation of pancreatic islets with 20 mM glucose caused a biphasic  $[\text{Ca}^{2+}]_i$  dynamics: a transient increase followed by a sustained in-