Title	An antifungal compound involved in symbiotic germination of <i>Cypripedium macranthos</i> var. <i>rebunense</i> (Orchidaceae)
Author(s)	Shimura, Hanako; Matsuura, Mayumi; Takada, Noboru; Koda, Yasunori
Citation	Phytochemistry, 68(10), 1442-1447 https://doi.org/10.1016/j.phytochem.2007.03.006
Issue Date	2007-05
Doc URL	http://hdl.handle.net/2115/27966
Type	article (author version)
File Information	PH68-10.pdf

[Instructions for use](#)

Title

An antifungal compound involved in symbiotic germination of *Cypripedium macranthos* var. *rebunense* (Orchidaceae)

Authors

Hanako Shimura, Mayumi Matsuura, Noboru Takada,
Yasunori Koda*

*Department of Botany, Graduate School of Agriculture, Hokkaido
University, Sapporo, 060-8589, Japan*

*Corresponding author. Tel.: +81 11 706 2478; fax: +81 11 706 2482.
E-mail address: yasunori@res.agr.hokudai.ac.jp (Y. Koda)

Received 1 September 2006; received in revised form 26 January
2007

Abstract

Germination of orchid seeds fully depends on a symbiotic association with soil-borne fungi, usually *Rhizoctonia* spp. In contrast to the peaceful symbiotic associations between many other terrestrial plants and mycorrhizal fungi, this association is a life-and-death struggle. The fungi always try to invade the cytoplasm of orchid cells to obtain nutritional compounds. On the other hand, the orchid cells restrict the growth of the infecting hyphae and obtain nutrition by digesting them. It is likely that antifungal compounds are involved in the restriction of fungal growth. Two antifungal compounds, lusianthrin and chrysin, were isolated from the seedlings of *Cypripedium macranthos* var. *rebunense* that had developed shoots. The former had a slightly stronger antifungal activity than the latter, and the antifungal spectra of these compounds were relatively specific to the nonpathogenic *Rhizoctonia* spp. The level of lusianthrin, which was very low in aseptic protocorm-like bodies, dramatically increased following infection with the symbiotic fungus. In contrast, chrysin was not detected in infected protocorm-like bodies. These results suggest that orchid plants equip multiple antifungal compounds and use them at specific developmental stages; lusianthrin maintains the perilous symbiotic association for germination and chrysin helps to protect adult plants.

Keywords: *Cypripedium macranthos* var. *rebunense*; Orchidaceae; Antifungal compounds; Symbiotic germination; Chrysin; Lusianthrin

1. Introduction

Symbiosis with soil-borne fungi is indispensable for germination of orchid seeds, which contain insufficient stores of substances required for germination. In contrast to the peaceful and sustainable symbiotic associations between many other terrestrial plants and mycorrhizal fungi, where the fungi accept photosynthetic products from host plants in exchange for nutritional salts and water, the association between orchids and fungi is strictly a life-and-death struggle. The hyphae of fungi always try to invade the cytoplasm of orchid cells to obtain nutritional compounds, as in a saprophytic infection. On the other hand, orchid cells inhibit the growth of the infecting hyphae to keep them in the apoplast, enclosed by plasma membranes (Peterson et al., 1996), and make the mycelium form dense coils called pelotons. In the developing protocorm, the pelotons are limited to cortical cells and are not found in the apical meristem (Peterson et al., 1998, Shimura and Koda, 2005). Orchid cells appear to obtain nutritional compounds by digesting pelotons through secretion of enzymes (Blakeman et al., 1976). When orchids cells fail to control fungal growth, the fungi prevail and prey on the developing protocorms. In contrast, when the orchids exhibit an excessive resistant response, a hypersensitive reaction occurs and the fungus fails to grow inside the protocorms (Hadley, 1970). In such cases, the developing protocorms starve. Thus, germination of orchid seeds requires quite a delicate balance.

Orchid plants appear to use both chemical and physical devices to control invasion by symbiotic fungi. Thickening of the cell wall and deposition of phenolic compounds seem to act as physical barriers (Beyrle et al., 1995). Blakeman et al. (1976) hypothesized that an increase in oxidative enzyme activity upon fungal infection, which was similar to the oxidative activity of plants against pathogens, may play a role in the defense response of the orchid. As more clear-cut chemical devices, some antifungal compounds are thought to be involved in the control of fungal growth (Hadley, 1982). Although some antifungal compounds

(phytoalexin) have been found in adult orchid plants (Fisch et al., 1973, Ward et al., 1975), their role in regulation of fungal growth during symbiotic seed germination remains to be elucidated. Beyrle et al. (1995) reported that a small amount of an antifungal compound was present in aseptic protocorms of *Orchis morio*, but the amount did not change with the inoculation of a symbiotic fungus, suggesting that this compound is not involved in regulating this symbiotic association during germination.

Cypripedium macranthos var. *rebunense*, a kind of lady-slipper orchid, is one of the most famous wild orchids in Japan, because it has become a symbol of threatened-plant conservation. In 1994, the "Law for the Conservation of Endangered Species of Wild Fauna and Flora, Japan" designated this variety as a domestic threatened species, and collection from natural habitats was strictly prohibited. Recently, we have developed two protocols for the micropropagation of this variety: by aseptic culture of protocorm-like bodies (PLBs) (Shimura and Koda, 2004), and by symbiotic germination with a symbiotic fungus (Shimura and Koda, 2005). In the latter study, we found that both nutritional deficiency and a lengthy culture period at a high temperature contribute to loss of the symbiotic association, and as a consequence, the protocorms turn brown and rot. These results suggest that thermal stress and nutrient deficiency reduce the production of antifungal compounds and as a result the protocorms fail to control the invasiveness of the fungus. Some of the characteristics of the symbiotic fungus (designated WO-97) were similar to those of *Rhizoctonia repens* (perfect stage; *Tulasnella* spp), which is known to be a symbiotic fungus for many orchid species. However, the sequence homology of 5.8S rDNA was less than 60% (unpublished data), suggesting that this fungus is different from *R. repens*.

In this paper we describe the isolation and structural elucidation of antifungal compounds from plantlets of this orchid,

and discuss the involvement of these compounds in symbiotic germination.

Abbreviation: PLBs, protocorm like bodies

2. Results and discussion

Ethanol extract of plantlets was sub-fractionated into *n*-hexane, EtOAc, butanol, and aqueous soluble fractions. Each solvent fraction obtained from plantlets (equivalent to 3 g fresh weight) was added to medium and their antifungal activities against WO-97 were examined. The EtOAc fraction showed the strongest antifungal activity, and the fungus failed to grow completely (Fig. 1). Considerable activity was also found in the *n*-hexane fraction.

The EtOAc fraction was subjected to further bioassay-guided purification based on the presence of antifungal activity. The activity was separated into two fractions after chromatography on a Sephadex LH-20 column, and the fractions were designated A and B in the order of elution. Fraction A was subjected to TLC and silica gel column chromatography because it showed low solubility in MeOH. One mg of active compound was obtained from fraction A. The ¹H NMR spectroscopic data for this compound were identical to those of chrysin (**1**) (Chen et al., 2003), a commercially available flavonoid (Acros Organics, New Jersey, USA). This compound had been reported to be an anticonvulsant found in *Passiflora* (Medina et al., 1990), a chemical component of an orchid, *Bulbophyllum odoratissimum* (Majumder and Sen, 1991), and a phytoalexin found in the sapwood of Rosaceae (Kokubun and Harborne, 1995).

Final purification of the active compound in fraction B was performed by successive HPLC procedures, and 1.2 mg of pure compound was obtained. The ¹H and ¹³C NMR spectroscopic data for compound B agreed with those of lusianthrin (**2**) (Majumder and

Lahiri, 1990). Although this compound has previously been isolated as a chemical component of a dried sample of an orchid plant, *Lusia indivisa* (Majumder and Lahiri, 1990), it has not been shown to have antifungal activity. The position of the methoxyl group was supported by NOE's between the methoxyl group and H-1 (8.4%) and H-3(2.2%) in a differential NOE experiment. The chemical structure of lusianthrin (2) resembles those of orchinol and hircinol, which have been isolated as phytoalexins from adult orchid plants (Fisch et al., 1973, Ward et al., 1975). Lusianthrin (2) showed slightly stronger antifungal activity than chrysin (1); the ED₅₀ values of both compounds against WO-97 were ca. 10 μ M.

The antifungal spectra of lusianthrin (2) and chrysin (1) were studied using 5 kinds of fungi, including 3 representative pathogens. Lusianthrin (2) showed strong antifungal activity against WO-97 and its closely related fungi, *Rhizoctonia repens*, but weak or no activity against the 3 pathogens (Fig. 2). Chrysin (1) also showed its strongest activity against WO-97 (Fig. 3).

To confirm whether or not lusianthrin (2) and chrysin (1) help to defend against infection by a symbiotic fungus, changes in the levels of these two compounds after the inoculation of WO-97 were examined. Due to the shortage of seeds, it is practically impossible to produce a sufficient amount of aseptic protocorms. Therefore, this experiment was carried out using aseptic protocorm-like bodies (PLBs) derived from germinated seeds (Shimura and Koda, 2004). To facilitate fungal infection, PLBs that had begun to differentiate roots were selected and inoculated with WO-97, and then sampled monthly. Microscopic observation revealed that invasion by hyphae into the cortical cells of the roots occurred at 1-2 months. After 3 months, pelotons were observed in the cells. Four months after inoculation, some of the PLBs turned brown and then rotted, and some remained white and healthy. Pelotons and digested pelotons were found in the cells of healthy PLBs (Fig. 4A). In contrast, no pelotons were found in the cells of brown PLBs, and monilioid cells,

which are nutritional storage hyphae, were observed (Fig. 4B), suggesting that the fungus preyed PLBs. Monilioid cells were not found in the cells of healthy PLBs.

Healthy PLBs were examined for the presence of antifungal compounds. The level of lusianthrin (2) in aseptic PLBs remained low (Fig. 5). On the other hand, the level in inoculated PLBs began to increase markedly after 3 months. The simultaneous occurrence of a large increase in the level of lusianthrin (2) and the formation of pelotons seems to indicate that the increased level of lusianthrin (2) inhibits mycelial growth and helps to enclose them within the plasma membrane to form pelotons. The level of lusianthrin (2) in PLBs 4 months after inoculation was twice the ED₁₀₀ (Fig. 5), yet hyphae and pelotons were found in the cells (Fig. 4A). This result suggests that the compound is localized in the cytoplasm of orchid cells and the concentration in the apoplast remains low to allow fungal growth. Since PLBs began to turn brown and die at this point in culture, it is possible that browning is due to autointoxication by the overproduction of lusianthrin (2). Therefore, the toxicity of lusianthrin (2) against the host *C. macranthos* var. *rebunense* was examined using PLBs. PLBs were transferred to Hyponex-Peptone medium (Shimura and Koda, 2004) that contained 0, 10, 30 or 100 μ M lusianthrin (2). Even at the highest concentration, browning of the PLBs was not observed until 4 months of culture (data not shown). This result suggests that the browning of PLBs cannot be attributed to autointoxication by the overproduction of lusianthrin (2).

Chrysin (1) that was present in the initial sample was not detected in the infected PLBs. The initial sample mostly consisted of plantlets that had already developed shoots. It is likely that chrysin (1) is an antifungal product of adult plants and is not formed in protocorms. We are now raising more than 500 micropropagated plants of this orchid in our Botanic Garden (Nagatani et al., 2006).

Studies to verify this speculation are now in progress using some of these plants.

3. Concluding remarks

Two antifungal compounds, lusianthrin (2) and chrysin (1), were isolated from the seedlings of *C. macranthos* var. *rebunense* that had developed shoots. The large increase in the level of lusianthrin (2) in PLBs after infection with the symbiotic fungus suggested that the compound is involved in regulating the symbiotic association during germination. In contrast, chrysin (1) was not detected in infected PLBs. Since orchid cells inevitably allow fungal infection during symbiotic germination, they have to be constitutively susceptible to fungal infection. It is likely that orchid plants equip multiple antifungal compounds and deploy them in specific developmental stages: lusianthrin (2) maintains the perilous symbiotic association for germination and chrysin (1) helps to protect adult plants as a phytoalexin. This is the first report on the antifungal compound that appears to be involved in symbiotic germination of orchid seeds.

4. Experimental

4.1. General

Silica gel (Wakogel C-300), Sephadex LH-20, and Sep-Pak C₁₈ cartridge (Waters) were used for column chromatography. TLC was conducted on Silica gel 70 F₂₅₄ Plate (Wako). A preparative column (Nova-Pak HR C₁₈, 100×25 mm, Waters) and an analytical column (Nova-Pak C₁₈, Radial-pak cartridge 100×8 mm, Waters) were used for HPLC. The elution profile was monitored by *A* at 210

nm. The ^1H (500 MHz) and ^{13}C (125 MHz) NMR spectra were obtained using a Bruker AMX500 spectrometer. CD_3OD or $\text{CDCl}_3/\text{CD}_3\text{OD}$ (1/1) was used as a solvent for NMR experiments. The proton signal of CHD_2OD (δ 3.30 ppm) was used as an internal standard in ^1H NMR experiments, whereas the carbon resonance of CD_3OD (δ 49.0 ppm) was used as an internal standard in the ^{13}C NMR experiments. The MS spectrum was recorded on a JEOL JMS-AX500 spectrometer.

4.2. *Plant material*

Since the collection of the orchid plants from the natural habitat is prohibited by law and their commercial value is extremely high, it is difficult to obtain sufficient plant material. For the past 7 years we have been studying the micropropagation of this orchid species through PLBs (Shimura and Koda 2004) and symbiotic germination with symbiotic fungi (Shimura and Koda 2005). For the subculture of micropropagated plantlets, normal and healthy plantlets were always selected, and plantlets that seemed to be inadequate for transplanting were dipped in EtOH and stored at -20°C . The plantlets obtained by both methods were mixed and subjected to the extraction of antifungal compounds.

4.3. *Isolation of chrysin and lusianthrin*

The plantlets (200 g fr. wt) were homogenized with a sufficient amount of EtOH to give a final EtOH- H_2O (4:1, v/v) extract and then passed through filter paper (Advantec, No.2). The filtrate was concentrated to 30 ml and the resultant aqueous residue was partitioned 3 times against equal volumes of *n*-hexane, EtOAc and *n*-BuOH, respectively. The EtOAc fraction (920 mg) was fractionated by a silica gel column (100×25 mm, EtOAc/toluene, stepwise). Strong antifungal activity was found in the EtOAc/toluene (1:4, v/v) fraction. This fraction (167 mg) was applied to a Sephadex LH-20 column (450×20 mm). Elution was

carried out with MeOH (300 ml) and then with MeOH-AcOH (999:1, 200 ml) at a flow rate of 1.5 ml min⁻¹. Antifungal activity was found in fractions eluted at 135-150 ml and 375-435 ml, and these fractions were designated A and B, respectively. Fraction A (13 mg) was subjected to TLC (hexane-EtOAc 1:1, *R_f* 0.45), and a final purification was carried out by silica gel column (85×3 mm, CHCl₃-EtOAc 15:1). Twenty-five 0.3-ml fractions were collected and fractions 7-11 contained 1.0 mg of the active compound, chrysin **1**, yellow powder. ¹H NMR (500 MHz, CD₃OD:CDCl₃=1:1): δ 7.94 (2H, *dd*, *J* = 1.6, 7.6 Hz, H-2 and H-6), 7.57 (1H, *tt*, *J* = 1.6, 7.6 Hz, H-4), 7.55 (2H, *br t*, *J* = 7.6 Hz, H-3 and H-5), 6.68 (1H, *s*, H-3), 6.49 (1H, *d*, *J* = 2.2 Hz, H-8), 6.28 (1H, *d*, *J* = 2.2 Hz, H-6); EIMS, 70eV, *m/z* (rel. int.): 254 [M]⁺ (100), 226 [M-CO]⁺ (23), 152 [M-C₈H₆]⁺ (17), 124 [M-C₉H₆O]⁺ (11). Fraction B (43 mg) was successively subjected to a Sep-Pak C₁₈ cartridge (Vac 20 cc, eluted with MeOH-H₂O, 4:1, v/v), prep. HPLC (MeOH-H₂O, 4:1, v/v, including 0.1% AcOH, 4ml min⁻¹, *R_t* 10.4 min), and analytical HPLC (MeCN-H₂O, 3:7, v/v, including 0.1% AcOH, 1 ml min⁻¹, *R_t* 30.2 min) to give lusianthrin **2** (1.2 mg, light yellow oil). ¹H NMR (500 MHz, CD₃OD): δ 9.49 (1H, *d*, *J* = 9.2 Hz, H-5), 7.53^a (1H, *d*, *J* = 9.0 Hz, H-9), 7.49^a (1H, *d*, *J* = 9.0 Hz, H-10), 7.12 (1H, *d*, *J* = 3.0, H-8), 7.06 (1H, *dd*, *J* = 3.0, 9.2 Hz, H-6), 6.84 (1H, *d*, *J* = 2.8 Hz, H-1), 6.67 (1H, *d*, *J* = 2.8 Hz, H-3), 3.87 (3H, *s*, CH₃O-); ¹³C NMR (125 MHz, CD₃OD): δ 158.7 (*s*, C-1), 158.0 (*s*, C-7), 155.4 (*s*, C-4), 135.9 (*s*, C-8a), 134.7 (*s*, C-10a), 130.5 (*d*, C-5), 128.4^b (*d*, C-9), 128.3^b (*d*, C-10), 125.9 (*s*, C-4b), 117.1 (*d*, C-8), 116.1 (*s*, C-4a), 112.1 (*d*, C-6), 103.3 (*d*, C-1), 102.1 (*d*, C-3), 55.7 (*q*, OCH₃); EIMS, 70eV, *m/z* (rel. int.): 240 [M]⁺ (100), 197 [M-C₂H₃O]⁺ (35). ^{a,b}Values are interchangeable with each other.

4.4. Assay for antifungal activity.

All cultures were carried out at 20°C in the dark. WO-97, a symbiotic fungus (Shimura and Koda, 2005), was used for the assay

of antifungal activity in the extracts. The fungus was cultured for 8 weeks on green pea medium, and inocula for assays were taken as 5-mm discs from the advancing edge of the colonies on agar plates. The sample to be tested was dissolved in MeOH and added to a test tube. After MeOH was removed under reduced pressure, green pea medium (10ml) was poured into the tube and autoclaved for 7 min. Before gelatinization, the medium was sonicated for 3 min to disperse water-insoluble compounds and then poured into a Petri dish. At the center of the dish, the fungal inoculum was placed upside-down so that the hyphae were in contact with the agar surface. Since WO-97 showed very slow growth, the diameter of the colony was measured 4 weeks after inoculation. Antifungal activity was expressed as inhibition of mycelial growth that was calculated by $[100 - (\text{average radius of colony on test medium}) / (\text{average radius of colony on basal medium}) \times 100]$.

In addition to WO-97, four kinds of fungi were used to assess the antifungal spectra of lusianthrin **2** and chrysin **1**; (1) *Rhizoctonia repens* R.r.1, a nonpathogenic fungus isolated from roots of *Spiranthes sinensis*; (2) *Rhizoctonia solani* AG2-2-IV, a pathogen of beet root-rot disease isolated from infected beet roots; (3) *Fusarium oxysporum* f.sp. *cepae*, a pathogen of onion bulb dry-rot disease isolated from infected onion bulbs; (4) *Phytophthora capsici* (ATCC 58208), a pathogen of gray blight disease. The Laboratory of Plant Pathology of our Graduate School preserves these fungal stocks on potato-sucrose-agar medium. Since these fungi show quite different growth rates, the diameter of the colony of each fungus was measured 7, 3, 6 and 3 days after inoculation, respectively. All experiments were carried out with 5 replicates.

4.5. Changes in the levels of antifungal compounds in PLBs after inoculation of the symbiotic fungus

WO-97 was inoculated at the center of a Petri dish that contained 10 ml green pea medium and had been cultured for 4

weeks. To facilitate fungal infection, PLBs that had developed roots were selected, and transferred to the Petri dish. The PLBs (ca. 4 g fr.wt) were harvested every month and subjected to extraction with 80% EtOH followed by EtOAc extraction after concentration. The EtOAc fraction was then purified by TLC (hexane-EtOAc 1:1, R_f 0.25). The amounts of lusianthrin **2** in these samples were measured by HPLC on an analytical column (MeCN-H₂O, 3:7, v/v, including 0.1% AcOH, 1 ml min⁻¹, A 254 nm, R_t 30.2 min). Since practical difficulties involved in obtaining a sufficient amount of uniform PLBs did not permit the replication of each experiment, the levels of aseptic control were determined 5 times with different lots. For light microscopy, the inoculated PLBs were fixed in a mixture of formaldehyde-AcOH-EtOH-H₂O (2:1:10:7), embedded in 5% agar and sectioned by a micro-slicer at 80 μ m. Hyphae were stained with 0.05% trypan blue in lactophenol (PhOH-glycerol-lactic acid-H₂O 1:1:1:1).

Acknowledgement

This work was supported in part by a Grant-in-Aid (Conservation study on designated national endangered plants with a model case of *Cypripedium macranthos* var. *rebunense*) from the Ministry of the Environment, Japan, and by a Grant-in-Aid (No. 16310157) from the Ministry of Education, Culture, Sports, Science and Technology of Japan. We are grateful to Ms. Mai Sadamoto for her assistance with maintaining the fungi, and to Prof. Shigeo Naito for providing the 3 representative pathogens. We also thank Dr. Eri Hukushi and Mr. Kenji Watanabe at the GC-MS & NMR Laboratory of the Faculty of Agriculture, Hokkaido University for the NMR and mass spectra measurements.

5. References

- Beyrle, H.F., Smith S.E., Peterson, R.L., Franco C.M.M., 1995. Colonization of *Orchis morio* protocorms by a mycorrhizal fungus: effects of nitrogen nutrition and glyphosate in modifying the responses. *Can. J. Bot.* 73, 1128-1140.
- Blakeman, J.P., Mokahel, M.A., Hadley, G., 1976. Effect of mycorrhizal infection on respiration and activity of some oxidase enzymes of orchid protocorms. *New Phytol.* 77, 697-704.
- Chen, L.J., Games, D.E., Jones, J., 2003. Isolation and identification of four flavonoid constituents from the seeds of *Oroxylum indicum* by high-speed counter-current chromatography. *J. Chromatogr. A* 988, 95-105.
- Fisch, M.H., Flick, B.H., Arditti, J., 1973. Structure and antifungal activity of hircinol, loroglossol and orchinol. *Phytochemistry* 12, 437-441.
- Hadley, G., 1970. Non-specificity of symbiotic infection in orchid mycorrhiza. *New Phytol.* 69, 1015-1023.
- Hadley, G., 1982. Orchid mycorrhiza. In: Arditti, J. (Ed.), *Orchid Biology: Reviews and Perspectives II*, Cornell University Press, Ithaca, NY, pp. 83-118.
- Kokubun, T., Harborne, J.B., 1995. Phytoalexin induction in the sapwood of plants of the Maloideae (Rosaceae): biphenyls or dibenzofurans. *Phytochemistry* 40, 1649-1654.
- Majumder, P.L., Lahiri, S., 1990. Lusianthrin and lusianthridin, two stilbenoids from the orchid *Lusia indivisa*. *Phytochemistry* 29, 621-624.
- Majumder, P.L., Sen, R.C., 1991. Bulbophyllanthrone, a phenanthraquinone from *Bulbophyllum odoratissimum*. *Phytochemistry* 30, 2092-2094.

- Medina, J.H., Paladini, A.C., Wolfman, C., Levi de Stein, M., Calvo, D., Diaz, L.E., Pena, C., 1990. Chrysin (5,7-di-OH-flavone), a naturally-occurring ligand for benzodiazepine receptors, with anticonvulsant properties. *Biochem. Pharmacol.* 40, 2227-31.
- Nagatani, K., Shimura, H., Matsuura, M., Koda, Y., 2006. Acclimation and potting of micro-propagated seedlings of *Cypripedium macranthos* var. *rebunense* obtained from aseptic culture and symbiotic germination, and maintenance of the potted plants. *Memoirs of the Graduate School of Agriculture Hokkaido University* 28, 121-131.
- Peterson, R.L., Bonfante, P., Faccio, A., Uetake, Y., 1996. The interface between fungal hyphae and orchid protocorm cells. *Can. J. Bot.* 74, 1861-1870.
- Peterson, R.L., Uetake, Y., Zelmer, C., 1998. Fungal symbioses with orchid protocorms. *Symbiosis* 25, 29-55.
- Shimura, H., Koda, Y., 2004. Micropropagation of *Cypripedium macranthos* var. *rebunense* through protocorm-like bodies derived from mature seeds. *Plant Cell Tiss. Org. Cult.* 78, 273-276.
- Shimura, H., Koda, Y., 2005. Enhanced symbiotic seed germination of *Cypripedium macranthos* var. *rebunense* following inoculation after cold treatment. *Physiol. Plant.* 123, 281-287.
- Ward, E.W.B., Unwin, C.H., Stoessl, A., 1975. Postinfectious inhibitors from plants. XV. Antifungal activity of the phytoalexin orcinol and related phenanthrenes and stilbenes. *Can. J. Bot.* 53, 964-971.

Legends for Figures.

Fig. 1. Antifungal activities of solvent fractions obtained from plantlets of *Cypripedium macranthos* var. *rebunense* against its symbiotic fungus, WO-97. Fractions (each equivalent to 3g fr. wt) were added to 10 ml of assay medium. WO-97 was inoculated onto the medium and cultured for 4 weeks.

Fig. 2. Antifungal spectrum of lusianthrin (2). The fungi used were; WO-97, a symbiotic fungus; *Rhizoctonia repens* R.r.1, a nonpathogenic fungus; *Rhizoctonia solani* AG2-2-IV, beet root-rot disease; *Fusarium oxysporum* f.sp. *cepae*, onion bulb dry-rot disease; *Phytophthora capsici*, gray blight disease. Numbers at the tops of the bars show the molar concentrations of the compounds. Bars indicate $\pm$ SD (n=5).

Fig. 3. Antifungal spectrum of chrysin (1). The fungi used were the same as in Fig. 2. Bars indicate $\pm$ SD (n=5).

Fig. 4. Microscopic observations of healthy (A) and brown (B) roots of protocorm-like bodies 4 months after inoculation. DP, digested peloton; M, monilioid cells; P, peloton.

Fig. 5. Time-course changes in the level of lusianthrin (2) in protocorm-like bodies after inoculation of a symbiotic fungus. Broken lines represent ED₅₀ and ED₁₀₀. SD (n=5) of the aseptic control was 1.35 n mol g fr.wt⁻¹.

EtOH extract