Title	Degradation of pentachlorophenol in contaminated soil suspensions by potassium monopersulfate catalyzed oxidation by a supramolecular complex between tetra(p-sulfophenyl)porphineiron(III) and hydroxypropyl-β-cyclodextrin.
Author(s)	Fukushima, Masami; Tatsumi, Kenji
Citation	Journal of Hazardous Materials, 144(1-2), 222-228 https://doi.org/10.1016/j.jhazmat.2006.10.013
Issue Date	2007-06-01
Doc URL	http://hdl.handle.net/2115/28125
Type	article (author version)
File Information	JHM144-1-2.pdf

[Instructions for use](#)

Revised manuscript: HAZMAT-D-06-00673

Degradation of pentachlorophenol in contaminated soil suspensions by potassium
monopersulfate catalyzed oxidation by a supramolecular complex between tetra
(*p*-sulfophenyl)porphineiron(III) and hydroxypropyl- β -cyclodextrin

Masami Fukushima^{a*} and Kenji Tatsumi^b

^a *Division of Solid Waste, Resources and Geoenvironmental Engineering, Graduate
School of Engineering, Hokkaido University, Sapporo 060-8628, Japan*

^b *National Institute of Advanced Industrial Science and Technology (AIST), 16-1
Onogawa, Tsukuba-West, Tsukuba 305-8569, Japan*

*Corresponding author. Tel./fax: +81-11-706-6304. E-mail address:
m-fukush@eng.hokudai.ac.jp (M. Fukushima)

Abstract

To enhance the catalytic oxidation of pentachlorophenol (PCP) in contaminated soil
suspensions using tetra(*p*-sulfophenyl)porphineiron(III) (Fe(III)-TPPS) as a catalyst and
potassium monopersulfate (KHSO₅) as the single-oxygen donor, the effect of added
hydroxypropyl- β -cyclodextrin (HP- β -CD) was examined. At pH 4 and 6, the percentage
of PCP disappearance increased substantially in the presence of HP- β -CD. In addition,
the self-degradation of Fe(III)-TPPS was significantly retarded in the presence of
HP- β -CD. This retarded self-degradation can be attributed to the stabilization of
Fe(III)-TPPS via the formation of a supramolecular complex with HP- β -CD. The kinetic

constant for the self-degradation of Fe(III)-TPPS in the presence of HP- β -CD at pH 6 was much smaller than that at pH 4, indicating that Fe(III)-TPPS is more stable at pH 6. Thus, the amount of Fe(III)-TPPS, KHSO₅ and HP- β -CD required to degrade PCP in contaminated soil suspensions was optimal at pH 6. When PCP-contaminated soil suspensions were treated under the optimized conditions, 12 – 18% and 24 – 28% of the PCP was mineralized to CO₂ in the absence and presence of HP- β -CD, respectively. These results show that the presence of HP- β -CD in the Fe(III)-TPPS/KHSO₅ catalytic system is effective in enhancing the degradation of PCP in contaminated soil suspensions.

Keywords: Pentachlorophenol; Soil remediation; Iron(III)-porphyrin; Hydroxypropyl- β -cyclodextrin; Supramolecular complex

1. Introduction

Pentachlorophenol (PCP) has, in the past, been utilized as a wood preservative and in herbicides. A recent study indicated that chlorophenols, such as PCP, are the main cause of detrimental health effects in humans in contaminated areas [1]. In addition, PCP can be converted into more toxic dimers, such as chlorinated dibenzo-*p*-dioxins and dibenzofurans, via a variety of oxidation processes in a soil environment [2, 3]. Thus, the degradation of PCP in contaminated soils could reduce the potential risk of pollution and related health issues. Technologies for the remediation of PCP-contaminated soils include chemical oxidation, washing the soil and biological degradation. The Fenton reaction is used for the chemical oxidation of PCP in contaminated soils [4, 5]. However, the disadvantage of the Fenton reaction is that large amounts of iron(II) and H₂O₂ are

1 required, and an iron(III) hydroxide sludge is produced during the reaction. To
2 overcome such problems, Liou et al. [6] developed an iron-resin catalyst for the
3 degradation of PCP by a Fenton-like system. In soil washing, PCP in the contaminated
4 soil is removed by extraction with organic solvents [7]. However, this may not be an
5 environmentally sound procedure. On the other hand, biological degradation of PCP
6 using microorganisms is known to be a practical process [8, 9]. However, the
7 introduction of non-indigenous microorganisms into contaminated soils is problematic
8 from an ecological point of view.

9 The use of a chemical catalyst to degrade PCP may circumvent some of the above
10 problems associated with bioremediation. It is known that iron(III)-porphyrins, which
11 are regarded as biomimetic models for active sites in lignase, are able to catalyze the
12 degradation of chlorophenols [10-13]. In general, the catalytic activities of
13 iron(III)-porphyrins are much less sensitive to variations in environmental conditions
14 than are microorganisms and even enzymes *in vitro*. Thus, the use of
15 iron(III)-porphyrins might make the maintenance of reaction conditions a more
16 straightforward procedure. However, iron(III)-porphyrins are unstable in the presence of
17 peroxides, such as H_2O_2 and KHSO_5 , because they are susceptible to self-degradation
18 [14, 15]. The self-degradation of iron(III)-porphyrins results in a decrease in the
19 oxidation efficiency of the organic substrate. We recently reported that the addition of
20 hydroxypropyl- β -cyclodextrin (HP- β -CD) was effective in stabilizing
21 iron(III)-porphyrin in the presence of KHSO_5 , due to the formation of a supramolecular
22 complex between HP- β -CD and iron(III)-porphyrin [16]. As shown in Fig. 1, the
23 supramolecular complex is formed by interactions between the sulfophenyl groups in
24 Fe(III)-TPPS and the hydrophobic cavity of HP- β -CD.

1 A few applications of the catalytic systems using iron(III)-porphyrins for the
2 remediation of PCP-contaminated soils have been reported [17, 18]. However, PCP
3 degradation (mineralization to CO₂) was achieved up to 10% [17]. Fe(III)-TPPS is
4 known to be a non-toxic biomimetic catalyst, and KHSO₅, commercially sold as
5 “Oxone”, is known to be a stable and environmentally safe oxidant. In addition,
6 HP-β-CD, a biologically derived material, has been used for the extraction of
7 chlorophenols from contaminated soils [19-21]. These facts suggest that combining the
8 Fe(III)-TPPS/KHSO₅ catalytic system and HP-β-CD could be a clean and safe process
9 for the remediation of PCP-contaminated soils. The purpose of the present study was to
10 examine the effect of added HP-β-CD on the degradation of PCP in the
11 Fe(III)-TPPS/KHSO₅ catalytic system. In order to apply the above system to the
12 treatment of PCP-contaminated soils, the optimal pH and dosage for the reaction were
13 determined. In addition, the mineralization of PCP to CO₂ was examined under optimal
14 conditions.

15

16 **2. Materials and Methods**

17 *2.1. Reagents and Materials*

18 Tetra(*p*-sulfophenyl)porphineiron(III) (Fe(III)-TPPS) was prepared according to a
19 method described in a previous report [22]. KHSO₅ was obtained as a triple salt,
20 2KHSO₅·KHSO₄·K₂SO₄ (Merck). PCP (99.0% purity) was purchased from Aldrich, and
21 a stock solution (0.01 M) was prepared by dissolving it in acetonitrile. ¹⁴C-labeled PCP
22 was purchased from American Radiolabeled Chemicals Inc. (0.1 mCi ml⁻¹ in ethanol,
23 specific activity 80 mCi mmol⁻¹). HP-β-CD (1.0 molar substitution, Mw 1540) was
24 purchased from Aldrich.

Kaolin was purchased from Kukita Yakuhin Kogyo (Tokyo) and was used without further treatment. Commercial Kanuma- and Red-soils for gardening were used in the present study. After air-drying, these soils were ground in a mortar with a pestle and then sieved through a stainless steel sieve (0.22 mm). A particle size below 0.22 mm was employed in the tests. The pH values, bet-N₂ specific surface areas and elemental compositions of the soil samples are summarized in Table 1. The specific surface area for Kanuma-soil was much larger than those for other samples. This can be attributed to the fact that Kanuma-soil includes large amounts of allophane moieties [23].

2.2. Batch catalytic oxidation of PCP in soil suspensions

A 0.2 g of soil sample was placed in a 10-ml glass tube. A 2 ml aliquot of 0.02 M NaH₂PO₄/Na₂HPO₄/citrate buffer at pH 4 or 6 was pipetted into the glass tube, and the solution was spiked with a 10 µl aliquot of 0.01 M PCP in acetonitrile. The suspension was then shaken for 24 h to allow the PCP to reach equilibrium with the soil. A 0 – 100 µl aliquot of aqueous Fe(III)-TPPS (200 µM) and a 0 – 50 µl aliquot of aqueous HP-β-CD (100 mg ml⁻¹) were added to the suspension, and a 0 – 50 µl of aqueous 0.01 M KHSO₅ was then added. The glass tube was then allowed to shake at room temperature (23 – 25°C). After 1, 3 or 5 days of shaking, 1 ml of 2-propanol was added and the suspension was allowed to shake for a further 2 h. After centrifugation of the soil suspension (2500 rpm for 3 min), the supernatant was filtered through a DISMIC[®] filter (diameter 13 mm, pore size 0.45 µm, hydrophilic PTFE type, ADVANTEC). It was confirmed that no adsorption of PCP to the filter occurred, when an aqueous solution of PCP (50 µM) in the absence of soil was passed through the filter. A 20 µl aliquot of the filtrate was injected into a JASCO PU-980 type HPLC pumping system.

The mobile phase consisted of a mixture of 0.08% aqueous H_3PO_4 and methanol (20/80 = v/v), and the flow rate was set at 1 ml min^{-1} . A 5C18-MS Cosmosil packed column (4.6 mm i.d. $\times$ 250 mm, Nacalai Tesque) was used as the solid phase, and the column temperature was maintained at 50°C . PCP was determined by Uv absorption at a wavelength of 220 nm. The efficiency of extraction of PCP from the contaminated soil suspensions was evaluated (Table 2). In the absence of both Fe(III)-TPPS and KHSO_5 (“PCP + buffer” in Table 2), 91 – 101% of the PCP was extracted from the soil suspensions at pH 4 and 6. All runs were conducted in triplicate.

2.3. Mineralization of ^{14}C -labeled PCP

A 0.5 ml aliquot of ^{14}C -labeled PCP in ethanol was evaporated in a stream of dry N_2 . The residue was then dissolved in 1 ml of acetonitrile (final concentration of PCP 625 μM). A 19.2 ml aliquot of buffer solution at pH 6 and a 0.8 ml aliquot of ^{14}C -labeled PCP in acetonitrile were mixed with stirring (final concentration of PCP 25 μM). A 0.1 g sample of soil was placed in a 20-ml Erlenmeyer flask. A 1 ml aliquot of the buffer solution, including ^{14}C -labeled PCP, was then added and 25 μl of aqueous Fe(III)-TPPS (400 μM) and 0 or 15 μl of aqueous HP- β -CD (100 mg ml^{-1}) were then added. After adding 25 μl of aqueous KHSO_5 (0.01 M), the flask was fitted with a stopper, which contained a polyethylene center well containing a glass wool plug impregnated with 0.4 ml of aqueous 2 M NaOH as a $^{14}\text{CO}_2$ trap. After a 5 day reaction period, 1 ml of 2-propanol and 0.4 ml of 1.8 M aqueous H_2SO_4 were injected via a syringe from the top of stopper to release the $^{14}\text{CO}_2$ from the soil suspension. After shaking for 1 day, a 2 ml aliquot of the soil suspension was centrifuged (6200 rpm, 1 min), and the supernatant was filtered thorough a DISMIC[®] filter. A 1 ml aliquot of the filtrate was pipetted into a

1 glass vial including a mixture of methanol (1 ml) and scintillation cocktail (9 ml). The
2 glass wool in the trap was transferred to the glass vial including 1.6 ml of water, and the
3 trap was then washed with 0.4 ml of 2 M NaOH, 0.8 ml of water and 1 ml of 2-propanol.
4 A 1 ml aliquot of this mixture was pipetted into a glass vial including methanol and
5 scintillation cocktail. The radioactivities (dpm) in the vials were determined using an LS
6 6000 series liquid scintillation counter (Beckman Instruments, Inc). To determine the
7 radioactivity before the reaction (R_{before}), a 1 ml aliquot of buffer solution including
8 ^{14}C -labeled PCP was added to an Erlenmeyer flask containing 0.1 g of soil. After
9 shaking for 5 days, the same procedures, as described above, were carried out. Each
10 experimental run was conducted in duplicate. The percentage of ^{14}C -labeled PCP
11 converted to $^{14}\text{CO}_2$ ($\%^{14}\text{CO}_2$) was calculated using the equation below:

$$12 \quad (\%^{14}\text{CO}_2) = (R_{\text{before}} - R_{\text{sus}})/R_{\text{before}} \times 100$$

13 where R_{sus} represents the radioactivities after the reaction in the soil suspension. In
14 addition, the recoveries of ^{14}C were calculated by dividing the sum of the radioactivities
15 in the trap and soil suspension by the radioactivity before the reaction. The recoveries in
16 all experimental runs ($n = 12$) were $92 \pm 7\%$.

17

18 *2.4. Uv-vis absorption spectra and kinetic measurement*

19 Uv-vis absorption spectra of a buffer solution at pH 4 or 6, containing Fe(III)-TPPS (5
20 μM) and HP- β -CD, were obtained on a Jasco V-550 type spectrophotometer (Japan
21 Spectroscopic Co.) with a quartz cell (1×1 cm) at 25°C . The kinetics of
22 self-degradation were monitored by the decolorization of Fe(III)-TPPS at 25°C . A 37.5
23 μl aliquot of aqueous KHSO_5 (0.01 M) was added to 3 ml of an aqueous solution in a 1
24 $\times 1$ cm quartz cell, which contained Fe(III)-TPPS (5 μM) and HP- β -CD (1.0 mg ml^{-1}) at

pH 4 or 6, with stirring. The absorbance of Fe(III)-TPPS before adding KHSO_5 (A_0) was monitored at 394 nm in the absence of HP- β -CD or at 419 nm in the presence of HP- β -CD. After adding KHSO_5 , the absorbance at arbitrary times (A_t) was determined at 0.2 s intervals up to 180 s using the kinetic mode of the spectrophotometer. No blank decolorization of Fe(III)-TPPS by only light at 394 nm or 419 nm was observed, even after 5 min of irradiation.

3. Results and Discussion

3.1. Control experiments

The mechanism of oxidation of PCP by Fe(III)-TPPS appears to proceed via a peroxide shunt, as described in a previous report [12]. A single-oxygen donor, such as KHSO_5 , is required to produce the active oxidants from Fe(III)-TPPS (ferryl-porphyrin cation radical and ferryl-oxo species). Such oxidants can be reduced to Fe(III)-TPPS by the oxidation of PCP to a pentachlorophenoxy radical ($\text{PCP}^\cdot$). The $\text{PCP}^\cdot$ is further oxidized, giving rise to a variety of byproducts are produced [10, 11, 13]. Prior to applying the Fe(III)-TPPS/ KHSO_5 catalytic system to the degradation of PCP in contaminated soil suspensions, the influence of some matrices in the soil suspensions on the disappearance of PCP was examined.

As shown in Table 1, the soil samples contained small amounts of manganese. It is known that manganese is present as manganese dioxide in soils, and that it is capable of oxidizing xenobiotics such as PCP [24]. Thus a control experiment to determine, whether the PCP in the soil suspension is degraded in the absence of both Fe(III)-TPPS and KHSO_5 or not, was carried out.

The recoveries of PCP from the soil suspensions are summarized in Table 2. In the

1 presence of Fe(III)-TPPS only (“PCP + buffer + Fe(III)-TPPS” in Table 2), the
2 recoveries were in the range of 92 – 99% at pH 4 and 6, indicating no influence of soil
3 components and Fe(III)-TPPS. However, in the presence of KHSO₅ only (“PCP + buffer
4 + KHSO₅” in Table 2), a significant lower recovery (59%) was observed in the Red-soil
5 at pH 4. As shown in Table 1, the contents of organic carbon and iron in the Red-soil
6 were much larger than those in the other soils. Paciolla et al. [25, 26] proposed that
7 ferryl-oxo species can be formed by the hydrogen peroxide catalysed oxidation by an
8 iron(III)-soil organic matter complex. Thus, it is likely that the lower recovery for the
9 Red-soil in the presence of KHSO₅ only at pH 4 may be due to the formation of the
10 ferryl-oxo species via a reaction between KHSO₅ and iron in the soil.

11

12 *3.2. Influence of pH on the disappearance of PCP*

13 In the presence of both Fe(III)-TPPS and KHSO₅, the influence of pH and HP-β-CD on
14 the percentage of PCP disappearance was investigated (Figure 2). It had previously been
15 reported that the percentage of PCP disappearance increased with increasing pH up to
16 pH 7 in an Fe(III)-TPPS/KHSO₅ catalytic system [11, 12]. As expected from the
17 previous reports, the percentage of PCP disappearance at pH 6 was larger than that at
18 pH 4 for all soil samples. In addition, the disappearance of PCP was remarkably
19 enhanced at pH 4 and 6 when HP-β-CD was added. In the presence of HP-β-CD, the
20 percentage of PCP disappearance at pH 6 was also larger than that at pH 4 for all soils.

21

22 *3.3. Influence of pH on the stability of Fe(III)-TPPS*

23 The deactivation of Fe(III)-TPPS is due to self-degradation via oxidation by KHSO₅ [14,
24 15]. The kinetics of the degradation of Fe(III)-TPPS were monitored at pH 4 and 6 in

1 the absence and presence of HP- β -CD (Fig. 3). The kinetic constants for the
2 self-degradation of Fe(III)-TPPS, calculated from curve-fitting to the data points in Fig.
3 3, are summarized in Table 3. The kinetic constant in the absence of HP- β -CD (k_0) at pH
4 6 was 10-times smaller than that at pH 4. In addition, the kinetic constant in the
5 presence of HP- β -CD ($k_{\text{HP-}\beta\text{-CD}}$) at pH 6 was 17-times smaller than that at pH 4. These
6 results indicate that the self-degradation of Fe(III)-TPPS is significantly retarded at pH
7 6 in the absence and presence of HP- β -CD. As shown in Table 3, the ratio of k_0 to
8 $k_{\text{HP-}\beta\text{-CD}}$ ($k_0/k_{\text{HP-}\beta\text{-CD}}$) at pH 6 was much larger than that at pH 4. This shows that the
9 retardation of Fe(III)-TPPS self-degradation by added HP- β -CD at pH 6 is greater than
10 that at pH 4.

11 The stabilization of Fe(III)-TPPS in the presence of HP- β -CD can be attributed to the
12 formation of a supramolecular complex [16]. Figure 4 shows the influence of pH and
13 HP- β -CD on the uv-vis absorption spectrum of Fe(III)-TPPS. At pH 6, the Soret band
14 for Fe(III)-TPPS at 394 nm in the absence of HP- β -CD was clearly shifted to 419 nm in
15 the presence of HP- β -CD (1 mg ml⁻¹), indicating the formation of a supramolecular
16 complex. However, at pH 4, a red-shift in the Soret band in the presence of HP- β -CD
17 was not clearly observed, but the band was broadened. The broadening of the peak at
18 pH 4 may be due to the contribution of both unbound and bound species of
19 Fe(III)-TPPS to HP- β -CD. To evaluate the ability of Fe(III)-TPPS to bind HP- β -CD, the
20 formation constants (K_f) were determined by a spectroscopic titration method, as
21 described in a previous report [16]. As shown in Table 3, the log K_f at pH 6 (5.1) was
22 larger than that at pH 4 (4.6). These results lead to the conclusion that, at pH 6, the
23 addition of HP- β -CD is more effective in stabilizing Fe(III)-TPPS than at pH 4.
24 Therefore, the conditions in the Fe(III)-TPPS/HP- β -CD/KHSO₅ catalytic system for

degrading PCP in contaminated soil suspensions were optimal at pH 6. As indicated in Table 1, the pH values of soil suspensions were around 6, except for the case of kaolin, which can be regarded as a control soil sample. It has been reported that pH values of soils are in the range of 3 – 7 [27]. Thus, pH control may be required, if the pH of the soil is in the acidic region.

3.4. Optimization of dosage

Figure 5 shows the effect of the concentration of Fe(III)-TPPS in the soil suspension on the percent PCP disappearance. For the Kaolin and Kanuma-soils, the percent PCP disappearance increased with increasing concentration of Fe(III)-TPPS up to 5 μM .

However, for the Red-soil, 10 μM of Fe(III)-TPPS was required to reach a plateau.

Figure 6 shows the influence of the concentration of KHSO_5 in a soil suspension on the percent PCP disappearance. Although the percent PCP disappearance increased with increasing concentration of KHSO_5 in all soils, a higher concentration of KHSO_5 was required in the Red-soil to reach the percent PCP disappearance comparable to the cases of Kaolin and Kanuma-soil. As shown in Table 1, organic carbon in the Red-soil (1.58%) was much higher than those in other soils. Because soil organic matter related to humic substances includes a variety of phenolic compounds [28], its presence may lead to the retardation of the catalytic oxidation of PCP. Thus, the larger dosages of Fe(III)-TPPS and KHSO_5 may be required in the case of the Red-soil.

Figure 7 shows the influence of the concentration of HP- β -CD in the soil suspension on the percent PCP disappearance. In all soils, the percent PCP disappearance increased with increasing concentration of HP- β -CD up to 1.0 mg ml^{-1} and then reached a plateau. In the present study, because the dosages should be reduced as much as possible, the

1 concentrations of Fe(III)-TPPS, KHSO₅ and HP-β-CD in the soil suspensions were set
2 to 10 μM, 250 μM and 1.5 mg l⁻¹, respectively. Although the influence of the reaction
3 period (1, 3 and 5 days) was investigated under optimal dosages, the percent PCP
4 disappearance did not vary significantly in any of the soils (89 – 96%).

6 3.5. Mineralization of PCP to CO₂

7 In the Fe(III)-TPPS/HP-β-CD/KHSO₅ catalytic system, tetrachloroquinone,
8 nonachlorodiphenyl ether and octachlorodibenzo-*p*-dioxin were detected as oxidation
9 products in the initial stage of the reaction (10 min) [16]. However, the levels of these
10 compounds decreased with increasing reaction time and the numbers of chlorine atoms
11 released from the PCP were 2.8 – 3.1, suggesting the further oxidation of the byproducts
12 [16]. However, because soil samples include large amounts of chloride ions,
13 dechlorination could not be determined in the present study. In remediation technologies
14 for soil contaminated with xenobiotics, mineralization of the xenobiotics is highly
15 desirable to reduce the potential risks of pollution. Therefore, the mineralization of PCP
16 to CO₂ was the major focus in the present study. Figure 8 shows the effect of HP-β-CD
17 on the percentage of mineralization of ¹⁴C-labeled PCP to ¹⁴CO₂ for a 5 day reaction
18 period. In all soils, the percentages of ¹⁴CO₂ in the presence of HP-β-CD (24 – 28%)
19 were larger than those in the absence of HP-β-CD (12 – 18%). These results
20 demonstrate that added HP-β-CD is effective in enhancing PCP degradation in
21 contaminated soil suspensions.

23 4. Conclusions

24 The addition of HP-β-CD to the Fe(III)-TPPS / KHSO₅ catalytic system was found to be

useful for enhancing the degradation of PCP in contaminated soil suspensions. Although Fe(III)-TPPS was stabilized by forming a supramolecular complex at pH 4 and 6, a pH of 6 was selected as optimal for the degradation conditions because the self-degradation of Fe(III)-TPPS was retarded to a greater extent. The combined processes of PCP extraction with some CDs and the degradation of PCP using a TiO₂ photocatalyst [20] and electrochemical method [21] have recently been reported. Thus, the Fe(III)-TPPS / HP-β-CD / KHSO₅ catalytic system, examined in the present study, may also be useful for the treatment of extracts of PCP-contaminated soils with aqueous HP-β-CD.

Acknowledgment

This work was supported by Grants-in-Aid for Scientific Research in Japan Society for the Promotion of Science (18201013). Authors wish to thank to Dr. H. Tamura (Graduate School of Engineering, Hokkaido University) for useful advice concerning the measurement of the bet-N₂ specific surface area.

References

- [1] T. Vartiainen, P. Lampi, J.T. Tuomisto and J. Tuomisto, Polychlorodibenzo-*p*-dioxin and polychlorodibenzofuran concentrations in human fat samples in a village after pollution of drinking water with chlorophenols, *Chemosphere* 30 (1995) 1429-1438.
- [2] M.M. Laine, J. Ahtianinen, N. Wågman, L.G. Öberg, K.S. Jørgensen, Fate and toxicity of chlorophenols, polychlorinated dibenzo-*p*-dioxins and dibenzofurans during composting of contaminated sawmill soil, *Environ. Sci. Technol.* 31 (1997) 3244-3250.
- [3] E.J. Hoekstra, H. de Weerd, E.W.B. de Leer, U.A.T. Btinkman, Natural formation of chlorinated phenols, dibenzo-*p*-dioxins, and dibenzofurans in soil of a Douglas fir forest,

1 Environ. Sci. Technol. 33 (1999) 2543-2549.

2 [4] R.J. Watts, M.D. Udell, R.M. Monsen, Use of iron minerals in optimizing the
3 peroxide treatment of contaminated soils, Wat. Environ. Res. 65 (1993) 839-844.

4 [5] J.X. Ravikumar, M.D. Guroul, Chemical oxidation of chlorinated organics by
5 hydrogen peroxide in the presence of sand, Environ. Sci. Technol. 28 (1994) 394-400.

6 [6] R.-M. Liou, S.-H. Chen, M.-Y. Hung, C.-S. Hsu, Catalytic oxidation of
7 pentachlorophenol in contaminated soil suspensions by Fe^{3+} -resin H_2O_2 , Chemosphere
8 55 (2004) 1271-1280.

9 [7] A.P. Khodadoust, J.A. Wagner, M.T. Suidan, S.I. Safferman, Solvent washing of
10 PCP contaminated soils with anaerobic treatment of wash fluids, Wat. Environ. Res. 66
11 (1994) 692-697.

12 [8] R.T. Lamar, J.W. Evans, J.A. Glaser, Solid-phase treatment of a
13 pentachlorophenol-contaminated soil using lignin-degrading fungi, Environ. Sci.
14 Technol. 27 (1993) 2566-2571.

15 [9] R.T. Lamar, M.W. Davis, D.M. Dietrich, J.A. Glaser, Treatment of a
16 pentachlorophenol- and creosote-contaminated soil using the lignin-degrading fungus
17 *Phanerochaete sordida*: a field demonstration, Soil Biol. Biochem. 26 (1994)
18 1603-1611.

19 [10] R.M. Shukla, A. Robert, B. Meunier, Kinetics investigations of oxidative
20 degradation of aromatic pollutant 2,4,6-trichlorophenol by an iron-porphyrin complex, a
21 model of lignase, J. Mol. Catal. A 113 (1996) 45-49.

22 [11] M. Fukushima, H. Ichikawa, M. Kawasaki, A. Sawada, K. Morimoto, K. Tatsumi,
23 Effects of humic substances on the pattern of oxidation products of pentachlorophenol
24 induced by a biomimetic catalytic system using tetra(*p*-sulfophenyl)porphineiron(III)

1 and KHSO_5 , Environ. Sci. Technol. 37 (2003) 1031-1036.

2 [12] M. Fukushima, A. Sawada, M. Kawasaki, H. Ichikawa, K. Morimoto, K. Tatsumi,
 3 M. Aoyama, Influence of humic substances on the removal of pentachlorophenol by a
 4 biomimetic catalytic system with a water-soluble iron(III)-porphyrin complex, Environ.
 5 Sci. Technol. 37 (2003) 386-394.

6 [13] S. Rismayani, M. Fukushima, A. Sawada, H. Ichikawa, K. Tatsumi, Effects of peat
 7 humic acids on the catalytic oxidation of pentachlorophenol using metalloporphyrins
 8 and metallophthalocyanines. J. Mol. Catal. A 217 (2004) 13-19.

9 [14] J.R.L. Smith, Supported metalloporphyrin models for heme-catalyzed oxidations,
 10 in: R.A. Sheldon (Ed.), Metalloporphyrins in Catalytic Oxidations, Dekker, New York,
 11 1994, pp. 325-327.

12 [15] M.J. Nappa, C.A. Tolman, Steric and electronic control of iron porphyrin catalyzed
 13 hydrocarbon oxidations, Inorg. Chem. 24 (1985) 4711-4719.

14 [16] M. Fukushima, K. Tatsumi, Effect of hydroxypropyl- β -cyclodextrin on the
 15 degradation of pentachlorophenol by potassium monopersulfate catalyzed with
 16 iron(III)-porphyrin complex, Environ. Sci. Technol. 39 (2005) 9337-9342.

17 [17] S.-T. Chen, D.K. Stevens, G. Kang, Pentachlorophenol and crystal violet
 18 degradation in water and soils using heme and hydrogen peroxide, Wat. Res. 33 (1999)
 19 3657-3665.

20 [18] M. Fukushima, A. Sawada, M. Kawasaki, K. Tatsumi, Removal of
 21 pentachlorophenol from contaminated soil by iron(III)-porphyrin complexes and
 22 potassium monopersulfate, Toxicol. Environ. Chem. 85 (2003) 39-49.

23 [19] K. Hanna, C. de Brauer, P. Germain, Solubilization of the neutral and charged form
 24 of 2,4,6-trichlorophenol by β -cyclodextrin, methyl- β -cyclodextrin and

1 hydroxypropyl- β -cyclodextrin in water, J. Haz. Mat. B100 (2003) 109-116.

2 [20] K. Hanna, C. de Brauer, P. Germain, J.M. Chovelon, C. Ferronato, Degradation of
3 pentachlorophenol in cyclodextrin extraction effluent using a photocatalytic process, Sci.
4 Total Environ. 332 (2004) 51-60.

5 [21] K. Hanna, S. Chiron, M. A. Oturan, Coupling enhanced water solubilization with
6 cyclodextrin to indirect electrochemical treatment for pentachlorophenol contaminated
7 soil remediation, Wat. Res. 39 (2005) 2763-2773.

8 [22] M. Kawasaki, A. Kuriss, M. Fukushima, A. Sawada, K. Tatsumi, Effects of pH and
9 organic co-solvents on the oxidation of naphthalene with peroxosulfate catalyzed by
10 iron(III) tetrakis(*p*-sulfonatophenyl)porphyrin, J. Por. Phthal. 7 (2003) 645-650.

11 [23] Y. Adachi, S. Koga, M. Kobayashi, M. Inada, Study of colloidal stability of
12 allophane dispersion by dynamic light scanning, Colloid Surface A 265 (2005) 149-154.

13 [24] R. A. Petrie, P.R. Grossl, R.C. Sims, Oxidation of pentachlorophenol in manganese
14 oxide suspensions under controlled Eh and pH environments, Environ. Sci. Technol. 36
15 (2002) 3744-3748.

16 [25] M.D. Paciolla, G. Davies, S.A. Jansen, Generation of hydroxyl radicals from
17 metal-loaded humic acids, Environ. Sci. Technol. 33 (1999) 1814-1818.

18 [26] M.D. Paciolla, S. Kolla, S.A. Jansen, The reduction of dissolved iron species by
19 humic acid and subsequent production of reactive oxygen species, Adv. Environ. Res. 7
20 (2002) 169-178.

21 [27] G.W. Thomas, Soil pH and soil acidity, in: D.L. Sparks, A.L. Page, P.A. Helmke,
22 R.H. Loeppert, P.N. Soltanpour, M.A. Tabatabai, C.T. Johnston, M.E. Sumner (Eds.),
23 Methods of soil analysis part 3: Chemical methods, Soil Science Society of America,
24 Madison, 1996, pp. 475-490.

- 1 [28] K.H. Tan, Humic Matter in Soil and the Environment. Principles and Controversies,
- 2 Dekker, New York, 2003, pp.75-126.

Figure Captions

Fig. 1. Chemical structures of Fe(III)-TPPS and HP- β -CD, and the formation of a supramolecular complex.

Fig. 2. Influence of pH and HP- β -CD on the percentage of PCP disappearance. Soils 0.2 g, [PCP] 50 μ M, [Fe(III)-TPPS] 10 μ M, [KHSO₅] 250 μ M, [HP- β -CD] 2.5 mg ml⁻¹, reaction period of 1 day.

Fig. 3. Influence of pH and HP- β -CD on the kinetics of the self-degradation of Fe(III)-TPPS. [Fe(III)-TPPS] 5 μ M, [KHSO₅] 125 μ M, [HP- β -CD] 1.0 mg ml⁻¹.

Fig. 4. Influence of pH and HP- β -CD on the uv-vis absorption spectrum of Fe(III)-TPPS (5 μ M).

Fig. 5. Influence of Fe(III)-TPPS concentration on percent PCP disappearance. Soils 0.2 g, [PCP] 50 μ M, [KHSO₅] 250 μ M, [HP- β -CD] 2.5 mg ml⁻¹, reaction period of 1 day, ■ kaolin, ● Kanuma-soil, ▲ Red-soil.

Fig. 6. Influence of KHSO₅ concentration on percent PCP disappearance. Soils 0.2 g, [PCP] 50 μ M, [Fe(III)-TPPS] 10 μ M, [HP- β -CD] 2.5 mg ml⁻¹, reaction period of 1 day, ■ kaolin, ● Kanuma-soil, ▲ Red-soil.

Fig. 7. Influence of HP- β -CD concentration on percent PCP disappearance. Soils 0.2 g, [PCP] 50 μ M, [Fe(III)-TPPS] 10 μ M, [KHSO₅] 250 μ M, reaction period of 1 day, ■ kaolin, ● Kanuma-soil, ▲ Red-soil.

Fig. 8. Effects of HP- β -CD on percent ¹⁴C-labeled PCP mineralization to ¹⁴CO₂. Soils 0.1 g, [PCP] 25 μ M, [Fe(III)-TPPS] 10 μ M, [KHSO₅] 250 μ M, [HP- β -CD] 1.5 mg ml⁻¹, reaction period of 5 days.

Fig. 1
(HAZMAT-D-06-00673)

Fig. 2
(HAZMAT-D-06-00673)

Fig. 3
(HAZMAT-D-06-00673)

Fig. 4
(HAZMAT-D-06-00673)

Fig. 5

(HAZMAT-D-06-00673)

Fig. 6
(HAZMAT-D-06-00673)

Fig. 7
(HAZMAT-D-06-00673)

Fig. 8

(HAZMAT-D-06-00673)

Table 1

The pH values, bet-N₂ specific surface areas and elemental compositions for soil samples.

Soil samples	Kaolin	Kanuma-soil	Red-soil
pH ^a	7.71	6.27	6.01
SSA (m ² g ⁻¹) ^b	2.72	106	26.3
OC (wt %) ^c	nd ^d	0.16	1.58
Al (wt %)	8.6	10.1	9.3
Si (wt %)	35.3	16.0	12.8
Ca (wt %)	0.09	0.77	0.36
Fe (wt %)	0.11	0.97	4.69
Mn (wt %)	0.002	0.020	0.079

^a The pH of the soil slurry was determined for a mixture of soil and water at a ratio of 1 : 2.5 (soil : water = w/w) [18].

^b Bet-N₂ specific surface area.

^c Organic carbon

^d Not detected

Table 2

Extraction efficiencies (%) of PCP from contaminated soil suspensions for three controls.

Patterns of control	Kaolin	Kanuma-soil	Red-soil
<i>pH 4</i>			
PCP + buffer	94 ± 5	101 ± 6	99 ± 3
PCP + buffer + Fe(III)-TPPS	97 ± 4	97 ± 5	99 ± 6
PCP + buffer + KHSO ₅	88 ± 7	99 ± 4	59 ± 3
<i>pH 6</i>			
PCP + buffer	91 ± 3	97 ± 7	96 ± 2
PCP + buffer + Fe(III)-TPPS	92 ± 4	96 ± 2	97 ± 3
PCP + buffer + KHSO ₅	94 ± 2	93 ± 4	89 ± 4

Table 3

Kinetic constants for the self-degradation of Fe(III)-TPPS (k_0 and $k_{\text{HP-}\beta\text{-CD}}$) and the formation constants of the supramolecular complex (K_f) at pH 4 and 6.

pH	k_0 (s^{-1})	$k_{\text{HP-}\beta\text{-CD}}$ (s^{-1})	$k_0/k_{\text{HP-}\beta\text{-CD}}$	$\log K_f$
4	1.3	7.3×10^{-2}	18	4.6
6	0.16	4.3×10^{-3}	36	5.1