



Title	A preliminary study of the genetic diversity of Xinjiang Tarim red deer ( <i>Cervus elaphus yarkandensis</i> ) using the microsatellite DNA method
Author(s)	MAHMUT, Halik; GANZORIG, Sumiya; ONUMA, Manabu; MASUDA, Ryuichi; SUZUKI, Masatsugu; OHTAISHI, Noriyuki
Citation	Japanese Journal of Veterinary Research, 49(3), 231-237
Issue Date	2001-11-30
DOI	10.14943/jjvr.49.3.231
Doc URL	<a href="http://hdl.handle.net/2115/2917">http://hdl.handle.net/2115/2917</a>
Type	bulletin (article)
File Information	KJ00002400390.pdf



[Instructions for use](#)

A preliminary study of the genetic diversity of Xinjiang  
Tarim red deer (*Cervus elaphus yarkandensis*) using the  
microsatellite DNA method

Halik Mahmut<sup>1,2\*</sup>, Sumiya Ganzorig<sup>3</sup>, Manabu Onuma<sup>2</sup>, Ryuichi  
Masuda<sup>4</sup>, Masatsugu Suzuki<sup>2</sup> and Noriyuki Ohtaishi<sup>2</sup>

(Accepted for publication : October 30, 2001)

Abstract

To evaluate the genetic diversity of the Xinjiang Tarim red deer (*Cervus elaphus yarkandensis*) population, we analyzed the frequencies of microsatellite alleles. Samples were collected from 3 isolated populations in Xaya, Lopnur and Qarqan of Xinjiang. Although 10 microsatellite loci were examined, alleles of 133 to 190 base-pairs were detected for only 3 loci: BM5004, BM4208 and BM888. The average observed multilocus heterozygosity was  $0.08 \pm 0.04$  for the Xaya, 0 for the Lopnur, and  $0.17 \pm 0.08$  for the Qarqan population. The average heterozygosity of all populations was  $0.08 \pm 0.02$ . The observed heterozygosities were significantly lower than the expected values. The present results suggest that the bottleneck effect has occurred in the populations of the Xinjiang Tarim red deer.

Key words : *Cervus elaphus yarkandensis*, microsatellite DNA, Tarim deer, Xinjiang

---

<sup>1</sup> Department of Biology, Xinjiang University, Urumqi, Xinjiang 830046, China

<sup>2</sup> Laboratory of Wildlife Biology, Graduate School of Veterinary Medicine, Hokkaido University, Sapporo 060-0818, Japan

<sup>3</sup> Laboratory of Parasitology, Graduate School of Veterinary Medicine, Hokkaido University, Sapporo 060-0818, Japan

<sup>4</sup> Laboratory of Genetic Diversity, Center for Advanced Science and Technology, Hokkaido University, Sapporo 060-0810, Japan

\*Corresponding author :

Halik Mahmut,

Laboratory of Wildlife Biology, Graduate School of Veterinary Medicine,

Hokkaido University, Sapporo 060-0818, Japan

TEL +81-11-706-7188

FAX +81-11-706-5569

E-mail : mahmut@vetmed.hokudai.ac.jp

## Introduction

The Tarim red deer (*Cervus elaphus yarkandensis*) inhabits the river valleys of the Tarim, Konqi and Qarqan rivers in Xinjiang, China (Fig. 1). Due to its limited distribution and population size, the Tarim deer has been included in the Red List (1994) of the International Union for the Conservation of Nature and Natural Resources (IUCN) in the category of an endangered species and classified as a second class protected animal in China. Currently, the habitat of the Tarim deer is an area of intensive human activity and the deer population become isolated into three areas, namely Xaya, Lopnur and Qarqan. Luo and Gu<sup>10)</sup> reported that the Tarim deer population is on the verge of extinction and that the number of individuals is decreasing in each area. So far, the protective measures that have been implemented regarding the Tarim deer have not been sufficiently effective, and the ecology and genetic features of this species remain to be elucidated.

Detailed knowledge of the reproduction and the genealogical relationships among individuals is essential to developing effective measures to protect endangered animals. An especially serious problem is caused by popu-




Fig. 1. Locations (1. Alar, 2. Xaya, 3. Bugur, 4. Lopnur, 5. Tikanlik, 6. Aktaz) of the sampled populations of the Tarim red deer (*Cervus elaphus yarkandensis*) in Tarim basin, Xinjiang, China.

lation bottlenecks: a severe reduction of population size, which reduces the diversity of a population and its genetic variability. This makes animals especially vulnerable to changes in their environment and to diseases.

Microsatellite analysis makes it possible to determine the genotypes of individuals by examining the variation of the number of repetitions of sequences that are several nucleotides long. The mutation rate of a given microsatellite locus is considered to be  $10^{-4} \sim 10^{-5}$  per generation<sup>4)</sup>. Microsatellite DNA is used as a marker of DNA polymorphism that is an effective hereditary index in ecological, evolutionary and genetic studies<sup>3,5,7)</sup>. Microsatellite analysis has been used for the study of genetic structures and reproductive success in deer species<sup>11,13,14,20)</sup>. In the present study, we obtained preliminary information about the genetic diversity in the Tarim deer populations using microsatellite polymorphisms.

## Materials and Methods

### Sample collection

The samples for the DNA analysis were collected during a field survey in September 2000 in the Tarim basin, Xinjiang. The deer hair samples were collected from both wild and captive Tarim deer from 3 isolated populations (Fig. 1), namely, those in Xaya (9 animals), Lopnur (5 animals) and Qarqan (4 animals).

### DNA extraction

DNA was extracted from the hairs using the chelex protocol<sup>21)</sup>.

### PCR

In the present study, 10 sets of primers were used as diagnostic markers (Table 1). All of these sets were designed based on sequence information from the genomes of bovine and ovine species and applied to genetic studies of

Table 1 . PCR primer sequences for microsatellite DNA analysis used in the present study

Microsatellite locus	Primer sequence (5' – 3')	Reference
BM203	GGGTGTGACATTTTGTTCCTC CTGCTCGCCACTAGTCCTTC	Bishop et al. (1994) <sup>11</sup>
BM848	TGGTTGGAAGGAAAACCTTGG CCTCTGCTCCTCAAGACAC	Bishop et al. (1994) <sup>11</sup>
BM888	AGGCCATATAGGAGGCAAGCTT CTCGGTCAGCTCAAACGAG	Bishop et al. (1994) <sup>11</sup>
BM4107	AGCCCCTGCTATTGTGTGAG ATAGGCTTTGCATTGTTTCAGG	Bishop et al. (1994) <sup>11</sup>
BM4208	TCAGTACACTGGCCACCATG CACTGCATGCTTTTCCAAAC	Bishop et al. (1994) <sup>11</sup>
BM5004	TCTGGAGTGAATGTTTCTGAGG TTGTGATGAGCACCTGAAGG	Bishop et al. (1994) <sup>11</sup>
BMC1009	GCACCAGCAGAGAGGACATT ACCGGCTATTGTCCATCTTG	Bishop et al. (1994) <sup>11</sup>
ETH152	TACTCGTAGGGCAGGCTGCCTG GAGACCTCAGGGTTGGTGATCAG	Steffen et al. (1993) <sup>181</sup>
OarFCB193	TTCATCTCAGACTGGGATTCAGAAAGGC GCTTGAAATAACCTCCTGCATCCC	Buchanan and Crawford (1993) <sup>21</sup>
VH110	CTCTAGAGGATCACAGAGAGTCGG GCAGAAACATTTTTTTTCTCAATATAGTTTCCC	Hanrahan et al. (1993) <sup>81</sup>

Cervidae<sup>9,14,16,17,19</sup>. The sets used were : BM203, BM848, BM888, BM4107, BM4208, BM5004, BMC1009, ETH152, OarFCB193, and VH110. The polymerase chain reaction (PCR) amplification was performed using Gene Amp PCR System 9700 (PE Byosystems). The PCR was performed in a total volume of 50 µl, using the primer pair and 20 ng of genomic DNA. The reaction mixture contained 0.8 µM [F]-dCTP and 5 units of TaKaRa Taq DNA polymerase (Takara, Japan) in buffer consisting of 10mM Tris-HCl (pH8.3), 50mM KCl, and 1.5 mM MgCl<sub>2</sub>. TAMRA-dCTP (yellow, 100µM) was obtained from ABI/PEC (product number 402793). PCR was performed using 35 cycles of a series of incubations at 94 °C (denaturation) for 1 min, 45–50°C (annealing) for 1 min, and 72°C (extension) for 1min. The annealing temperature was reduced to 45°C for OarFCB 193, 46°C for BM848 and VH110, 47 °C for BM 203, BM888, BM5004 and ETH152, and 50°C

for BM4107, BM4208 and BMC1009 marker amplification. A final extension step was performed at 72°C for 10 min.

#### Genotyping

An aliquot (0.5 µl) of the PCR products was mixed with 2.5 µl of deionized formamide, 0.5 µl of Blue Dextran/ethylenediaminetetraacetic acid (EDTA), and 0.5 µl of GeneScan 1000 Red Dye (ROX) marker. The mixture was then loaded on a 6% denaturing polyacrylamide gel (ABI PRISM Geluxe 377–36 WTR) and electrophoresed for 2 hr using a DNA sequencer ABI-PRISM 377.

#### Data analysis

Genetic polymorphism was expressed as the number of alleles per locus, while the observed heterozygosity (Ho) and expected heterozygosity (He ; based on the Hardy-Weinberg assumption) were calculated using the

GENEPOP package, version 3.1b<sup>15</sup>). This program was also used to test for deviations from Hardy-Weinberg equilibrium within a population at a given locus and over all loci.

### Results

Among the 10 microsatellite loci examined, clear bands of 133 to 190 base-pairs representing alleles were detected for only 3 loci, namely BM5004, BM4208 and BM888. At the BM5004 locus, 2 alleles were found in all 3 populations. At the BM4208 locus, 4 alleles were found in Xaya, 3 in Lopnur and 1 in Qarqan. At the BM888 locus, 2 alleles were found in Xaya and single bands were observed

in Lopnur and in Qarqan. At least 1 common allele was found for each locus in all 3 populations of Tarim deer. Table 2 shows the allele frequencies and observed heterozygocities of the 3 loci. The average observed multilocus heterozygosity was  $0.08 \pm 0.04$  for the Xaya, 0 for the Lopnur,  $0.17 \pm 0.08$  for the Qarqan population.

The average heterozygosity of all populations was  $0.08 \pm 0.02$ .

For the other 7 loci examined (BM203, BM848, BM4107, BMC1009, ETH152, OarFCB 193 and VH110), no allele were identified, probably due to mismatching of primer sequences or due to null alleles.

Table 2 . Microsatellite variation in the Tarim red deer (*Cervus elaphus yarkandensis*) populations

	Locus	Xaya	Lopnur	Qarqan	Total	
BM5004	No. of individuals	9	5	4	18	
	Allele <sup>a)</sup> & frequency	133	0.833	0.400	0.375	0.611
		137	0.167	0.600	0.625	0.389
		ho <sup>b)</sup>	0.111	0	0.250	0.111
		he <sup>c)</sup>	0.278	0.480	0.469	0.475
BM4208	No. of individuals	7	5	4	16	
	Allele <sup>a)</sup> & frequency	144	0	0.200	0.250	0.125
		148	0.571	0.200	0	0.313
		150	0.357	0.600	0.750	0.531
		162	0.071	0	0	0.031
	ho <sup>b)</sup>	0.143	0	0	0.063	
he <sup>c)</sup>	0.541	0.320	0.375	0.604		
BM888	No. of individuals	9	5	4	18	
	Allele <sup>a)</sup> & frequency	200	1	1	0.875	0.972
		198	0	0	0.125	0.028
		ho <sup>b)</sup>	0	0	0.250	0.056
		he <sup>c)</sup>	0	0	0.219	0.054
Average over 3 loci $\pm$ SE	No. of individuals	8.33 $\pm$ 0.67	5.00 $\pm$ 0.00	4.00 $\pm$ 0.00	17.3 $\pm$ 0.7	
A	0.38 $\pm$ 0.14	0.38 $\pm$ 0.12	0.38 $\pm$ 0.12	0.38 $\pm$ 0.12		
Ho <sup>b)</sup>	0.08 $\pm$ 0.04	0	0.17 $\pm$ 0.08	0.08 $\pm$ 0.02		
He <sup>c)</sup>	0.27 $\pm$ 0.16	0.27 $\pm$ 0.14	0.35 $\pm$ 0.07	0.38 $\pm$ 0.17		

<sup>a)</sup>Molecular sizes (bases) refer to allele name.

<sup>b)</sup>ho and Ho : observed heterozygosity.

<sup>c)</sup>he and He : expected heterozygosity.

## Discussion

The Tarim red deer lives in a desert landscape and its morphology differs from that of other red deer. Due to the poor condition of its habitat and the influence of recent agricultural expansion, very few Tarim deer have survived. Gao and Gu<sup>6)</sup> reported that only 15,000 of these deer were surviving in the Tarim basin. Recent studies showed that the population of Tarim deer in Qarqan does not number more than 50<sup>10)</sup>, while there are around 200 deer in Lopnur and Xaya. The habitat fragmentation in the Tarim basin is thought to influence the effective population size of this deer. However, little is known about the genetic status of the local populations. Studies of the genetic variations within a population of the Japanese sika deer<sup>12, 20)</sup> or the North American wapiti<sup>14)</sup> revealed that microsatellite DNA analysis is very useful for detecting population bottlenecks, inbreeding and other genetic parameters of deer populations. In the present study, we tried to find polymorphic loci that would be useful for monitoring genetic diversity of deer populations. Only 3 of 10 primer sets used provided clear results. The other 4 sets were not informative although they had been informative in other deer species<sup>11, 13, 14, 20)</sup>. The low level of amplification obtained in the present research may have been due to mismatching of primer sequences or to null alleles.

The limited number of samples used in the present study was not sufficient to warrant broader conclusions about the level of inbreeding or genetic bottleneck. The results showed that the observed heterozygosity was lower than that expected, with the exception of BM888 at Qarqan. This reduced heterozygosity suggests that the population of Tarim deer in Xinjiang has passed through bottleneck(s), and, as mentioned above, that the

population size of Tarim deer has been reduced.

The mean heterozygosity of the Tarim red deer population was  $H_o = 0.08 \pm 0.02$ . By comparison, the wapiti have a higher observed mean heterozygosity,  $H_o = 0.552 \pm 0.039$ , at the BM5004, BM4208 and BM888 loci<sup>19)</sup>. The low  $H_o$  of Tarim red deer may result from inbreeding due to population subdivision and shrinkage. In order to more precisely evaluate the genetic diversity of the Tarim red deer, we will have to obtain more microsatellite data.

Genetic research provides information that is important for establishing conservation plans for endangered species. It is essential to accumulate further genetic data about Tarim red deer in order to make an effective conservation plan.

## Acknowledgments

This study was made possible by the help of Xinjiang University and the Department of Education, Science and Forestry of Xinjiang. We thank the staff of the Laboratory of Wildlife Biology, Graduate School of Veterinary Medicine, Hokkaido University. We are also grateful to Prof. Ablimit Abdukadir (Xinjiang Institute of Ecology and Geography, Chinese Academy of Sciences) and Mr. Xiaoming Jiang (Regiment 33, Division 2 of Agriculture Xinjiang Production and Construction Corporation) for their invaluable advice. This work was partially supported by Grants-in-Aid for Scientific Research from the Ministry of Education, Science, Sports and Culture of Japan (No:08041131).

## References

- 1) Bishop, M. D., Kappes, S. M., Keele, J. W., Stone, R. T., Sunden, S. L. F., Hawkins, G. A., Solonas-Toldo, S., Fries, R., Grosz, M. D., Yoo, J. and Beattie, C. W. 1994. A genetic linkage map of cattle.

- Genet.*, 136 : 619-639.
- 2) Buchanan, F. C. and Crawford, A. M. 1993. Ovine microsatellites at the OarFCB11, OarFCB128, OarFCB193, OarFCB226 and OarFCB304 loci. *Anim. Genet.*, 24 : 145.
  - 3) Craighead, L., Paetkau, D., Reynolds, H. V., Vyse, E. R. and Strobeck, C. 1995. Microsatellite analysis of paternity and reproduction in Arctic grizzly bears. *J. Heredity*, 86 : 255-261.
  - 4) Dallas, J. F. 1992. Estimation of microsatellite mutation rates in recombinant inbred strains of mouse. *Mamm. Genome*, 5 : 32-38.
  - 5) Ellegren, H. 1992. Polymerase-chain reaction (PCR) analysis of microsatellites. A new approach to studies of genetic relationships in birds. *The Auk*, 109 : 886-895.
  - 6) Gao, X. and Gu, J. 1985. Red Deer in Xinjiang. *Chinese Wildlife*, 2 : 24-26. (in Chinese)
  - 7) Girman, D. J., Mills, M. G. L., Feffen, E. and Wayne, R. K. 1997. A molecular genetic analysis of social structure, dispersal, and interpack relationship of the African wild dog (*Lycaon pictus*). *Behav. Ecol. Sociobiol.*, 40:187-198.
  - 8) Hanrahan, V., Ede, A. J., Pierson, C. A. and Hill, D. F. 1993. Ovine microsatellites at the OarVH98, OarVH110, OarVH116, OarVH117 and OarVH130 loci. *Anim. Genet.*, 24 : 223.
  - 9) Kuhn, R., Anastassiadis, C. and Pirchner, F. 1996. Transfer of bovine microsatellites to the cervine (*Cervus elaphus*). *Anim. Genet.*, 27 : 199-201.
  - 10) Luo, N. and Gu, J. 1993. Status of Yarkand deer and strategy for its conservation and utilization. *Acta Zool. Arid Inland*, 1 : 38-41. (in Chinese)
  - 11) Miyazaki, K., Yamauchi, K., Hamasaki, S., Kikusui, T., Takeuchi, Y. and Mori, Y. 2001. Identification of individual sika deer (*Cervus nippon*) by fecal DNA analysis. *Jpn. J. Zoo Wildl. Med.*, 6(1): 1-6. (in Japanese with English abstract)
  - 12) Nagata, J., Masuda, R., Kaji, K., Ochiai, K., Asada, M. and Yoshida, M. C. 1998. Microsatellite DNA variations of the sika deer, *Cervus nippon*, in Hokkaido and Chiba. *Mamm. Study*, 23 : 95-102.
  - 13) Okada, A. and Tamate, H. B. 2000. Pedigree analysis of the sika deer (*Cervus nippon*) using microsatellite markers. *Zool. Sci.*, 17:335-340.
  - 14) Polziehn, R. O., Hamr, J., Mallory, F. F. and Strobeck, C. 2000. Microsatellite analysis of North American wapiti (*Cervus elaphus*) populations. *Mol. Ecol.*, 9 : 1561-1576.
  - 15) Raymond, M. and Rousset, F. 1995. GENEPOP (version 1.2) : population genetic software for exact tests and ecumenicism. *J. Heredity*, 86 : 248-249.
  - 16) Roed, Kh. 1998. Microsatellite variation in Scandinavian Cervidae using primers derived from Bovidae. *Hereditas*, 129 : 19-25.
  - 17) Slate, J., Coltman, D. W., Goodman, S. J., MacLean, I., Pemberton, J. M. and Williams, J. L. 1998. Bovine microsatellite loci are highly conserved in red deer (*Cervus elaphus*), sika deer (*Cervus nippon*) and Soay sheep (*Ovis aries*). *Anim. Genet.*, 29 : 307-315.
  - 18) Steffen P., Eggen A., Dietz A. B., Womack J. E., Strazinger G. and Fries R. 1993. Isolation and mapping of polymorphic microsatellites in cattle. *Anim. Genet.*, 24 : 121-124.
  - 19) Talbot, J., Haigh, J. and Plante, Y. 1996. A parentage evaluation test in North American elk (Wapiti) using microsat-

- ellites of ovine and bovine origin. *Anim. Genet.*, 27 : 117-119.
- 20) Tamate, H. B., Okada, A., Minami, M., Ohnishi, N., Higuchi, H. and Takatsuki, S. 2000. Genetic variations revealed by microsatellite markers in a small population of the sika deer (*Cervus nippon*) on Kinkazan Island, Northern Japan. *Zool. Sci.*, 17:47-53.
- 21) Walsh P. S., Metzger, D. A. and Higuchi, R. 1991. Chelex 100 as a medium for simple extraction of DNA for PCR-based typing from forensic material. *Bio. Techniques*, 10 : 506-513.