



Title	BIOTIN-LABELED ANTIGEN SANDWICH ENZYME-LINKED IMMUNOSORBENT ASSAY (BLA-S-ELISA) FOR DETECTION OF HEMORRHAGIC FEVER WITH RENAL SYNDROME (HFRS) ANTIBODY IN HUMAN AND A VARIETY OF ANIMAL SERA
Author(s)	YAHARA, Yoshihiro
Citation	Japanese Journal of Veterinary Research, 34(2), 165-165
Issue Date	1986-04-30
Doc URL	http://hdl.handle.net/2115/3012
Type	bulletin (article)
File Information	KJ00002374419.pdf



[Instructions for use](#)

BIOTIN-LABELED ANTIGEN SANDWICH ENZYME-LINKED
IMMUNOSORBENT ASSAY (BLA-S-ELISA) FOR DETECTION
OF HEMORRHAGIC FEVER WITH RENAL SYNDROME (HFRS)
ANTIBODY IN HUMAN AND A VARIETY OF ANIMAL SERA

Yoshihiro YAHARA

*Department of veterinary public Health
Faculty of Veterinary Medicine
Hokkaido University, Sapporo 060, Japan*

Biotin-labeled antigen sandwich (BLA-S) ELISA was developed to detect hemorrhagic fever with renal syndrome (HFRS) antibody in sera from humans and a variety of animals, regardless of species- and class-specificities of immunoglobulins. The assay was evaluated for its practicability in sero-epidemiological surveys. HFRS antibody was detected by being sandwiched between the antigens, and the avidin biotin system was employed to enhance sensitivity and specificity.

The results were summarized as follows:

- 1) BLA-S-ELISA could detect HFRS antibodies in the sera from HFRS infected humans and from all species of animals tested (rabbit, laboratory rat, urban rat, mouse, guinea pig, mongolian gerbil, *Apodemus (Ap.) agrarius*, *Clethrionomys (Cl.) rufocanus bedfordiae* and *Ap. argenteus*). BLA-S-ELISA showed almost the same sensitivity as the immune adherence hemagglutination (IAHA) test. Antibody titers in the sera from normal animals were less than 1:5 in all species tested.
- 2) BLA-S-ELISA antibody titers were extremely low (less than 1:10) in the sera from urban rats from an HFRS non-endemic area. In the sera of urban rats from an endemic focus, antibody titers significantly correlated to each other between BLA-S-ELISA and IFA, and also between IAHA and ELISA. BLA-S-ELISA could detect a low level of antibody in the sera judged as negative by IFA, IAHA and ELISA.
- 3) Five species of animals (rat, mongolian gerbil, *Ap. agrarius*, *Cl. rufocanus bedfordiae* and *Ap. argenteus*) were experimentally infected with HFRS viruses (SR-11, TB-314, KI-262, H-76-118 and MP-40). All species of animals produced a high level of HFRS antibody against each strain.
- 4) Cross BLA-S-ELISA was performed using sera of rats infected with each of these HFRS virus strains. The 5 strains could be classified into 3 types depending on the species of host animals: *Rattus*-borne type (SR-11, TB-314 and KI-262), *Apodemus*-borne type (H-76-118) and *Microtus*-borne type (MP-40). Thus, BLA-S-ELISA was found to be a more strain-specific method than other serological diagnostic methods.