



Title	The antioxidative function of eicosapentaenoic acid in a marine bacterium, <i>Shewanella marinintestina</i> IK-1
Author(s)	Nishida, Takanori; Morita, Naoki; Yano, Yutaka; Oriyasa, Yoshitake; Okuyama, Hidetoshi
Citation	FEBS Letters, 581(22), 4212-4216 https://doi.org/10.1016/j.febslet.2007.07.065
Issue Date	2007-09-04
Doc URL	http://hdl.handle.net/2115/30186
Type	article (author version)
File Information	FEBSL581-22.pdf



[Instructions for use](#)

The antioxidative function of eicosapentaenoic acid in a marine bacterium, *Shewanella*

marinintestina IK-1

Takanori Nishida^a, Naoki Morita^b, Yutaka Yano^c, Yoshitake Orikasa^a, and Hidetoshi Okuyama^{a,*}

^a *Graduate School of Environmental Earth Science, Hokkaido University, Kita-ku, Sapporo*

060-0810, Japan

^b *Research Institute of Genome-based Biofactory, National Institute of Advanced Industrial*

Science and Technology (AIST), Toyohira-ku, Sapporo 062-8517, Japan,

^c *National Research Institute of Fisheries Science, Fisheries Research Agency, Kanazawa-ku,*

Yokohama 236-8648, Japan

Abstract When the eicosapentaenoic acid (EPA)-deficient mutant strain IK-1Δ8 of the marine EPA-producing *Shewanella marinintestina* IK-1 was treated with various concentrations of hydrogen peroxide (H₂O₂), its colony-forming ability decreased more than that of the wild type. Protein carbonylation, induced by treating cells with 0.01 mM H₂O₂ under bacteriostatic conditions, was enhanced only in cells lacking EPA. The amount of cells recovered from the cultures was decreased more significantly by the presence of H₂O₂ for cells lacking EPA than for those producing EPA. Treatment of the cells with 0.1 mM H₂O₂

resulted in much lower intracellular concentrations of H₂O₂ being consistently detected in cells with EPA than in those without EPA. These results suggest that cellular EPA can directly protect cells against oxidative damage by shielding the entry of exogenously added H₂O₂ in *S. marinintestina* IK-1.

Keywords: Eicosapentaenoic acid; Hydrogen peroxide; Oxidative stress; Protein carbonyl;

Shewanella

*Corresponding author. Fax: +81 11 706 2347.

E-mail address: hoku@ees.hokudai.ac.jp (H. Okuyama)

Abbreviation: CFU, colony-forming unit; EPA, eicosapentaenoic acid; H₂O₂⁺, in the presence of H₂O₂; H₂O₂⁻, in the absence of H₂O₂; LB, Luria-Bertani; OD, optical density; PBS, phosphate-buffered saline.

1. Introduction

Previously we showed that eicosapentaenoic acid (EPA) has the potential to prevent the entry of H₂O₂ through the cell membrane in *Escherichia coli* strains genetically modified to

produce EPA. The *E. coli* strains DH5 α [1] and UM2 [2] transformant cells, which contained EPA, became resistant to oxidative stress by exogenous H₂O₂. When the *E. coli* strains were transformed with the EPA biosynthesis genes (*pfaA*, *pfaB*, *pfaC*, *pfaD*, and *pfaE*) [3] from *Shewanella pneumatophori* SCRC-2738 [4], the transformant produced EPA at levels of approximately 10% of total fatty acids. In the case of *E. coli* strain DH5 α , the colony-forming ability of EPA-containing cells treated with 3.0 mM H₂O₂ was similar to that of cells that were not exposed to H₂O₂. However, colony-forming ability was lost rapidly in cells without EPA under the same conditions. This was reflected in the difference in the degree of protein carbonylation between the strain with EPA and that without EPA. The treatment of both types of cells with H₂O₂ changed the amount of cells recovered from the cultures (estimated by the amount of fatty acid). The recovery of fatty acids was lower from cultures of *E. coli* cells without EPA than those with EPA, but it did not affect the fatty acid compositions. From these data, it has been suggested that cellular EPA is stable *in vivo* in the presence of H₂O₂ and may directly protect cells against oxidative damage. Using catalase-deficient *E. coli* UM2 that had been transformed with the same EPA biosynthesis genes, almost the same results were obtained [2], although the cells of *E. coli* UM2 were treated with 0.3 mM H₂O₂. In the case of *E. coli* UM2, cells were suspended in phosphate-buffered saline (PBS) and then treated with H₂O₂ under bacteriostatic conditions (medium-free and no agitation). Much lower intracellular concentrations of H₂O₂ were detected in cells with EPA than in those

lacking EPA. From these results, it was suggested that cellular EPA could directly protect cells against oxidative damage by preventing the entry of exogenously added H₂O₂. A similar antioxidative mechanism is thought to operate in bacteria that produce EPA naturally. However, no evidence for this was found in the literature.

Shewanella marinintestina IK-1 is an EPA-producing bacterium that was isolated from a squid body [5]. Although EPA is regarded as a modulator of membrane fluidity at low temperature also in this bacterium [5], we present here the results of an investigation into the possible role of EPA in protecting against oxidative stress (exogenous addition of H₂O₂) in bacteria that inherently produce EPA. The study compared the responses against exogenously added H₂O₂ of *S. marinintestina* IK-1 and its EPA-deficient mutant (strain IK-1Δ8).

2. Materials and methods

2.1. Bacterial cells and cultivation

S. marinintestina IK-1 [5] and its EPA-deficient mutant *S. marinintestina* IK-1Δ8 (see below) were used as the test organisms throughout the work. Both strains were precultured by agitation at 180 rpm in Luria–Bertani (LB) medium containing 3.0% (w/v) NaCl normally at 15 °C. Nongrowth cells of the both strains were prepared as follows. The cells were grown until the

culture had an optical density of 1.0 at 600 nm (OD₆₀₀) and were harvested by centrifugation at 3500 × g for 5 min at 4 °C. Collected cells were washed three times with PBS containing 3.0% (w/v) NaCl and suspended in the same buffer at an OD₆₀₀ of 1.0. The cell suspension was mixed with a solution of H₂O₂ (30% solution in water; Wako Chemicals, Osaka, Japan) normally at a final concentration of 0.01 mM and then incubated without agitation for 30 min at 15 °C. Cells were harvested at appropriate time intervals by centrifugation and used for further assay. The number of colony-forming units (CFUs) was measured as previously described [1].

2.2. Generation of EPA-deficient mutant of *S. marinintestina* IK-1

S. marinintestina IK-1 has an EPA biosynthesis gene cluster (N. Morita, Y. Yano, S. Ohgiya, and H. Okuyama, unpublished results), of which structure is very similar to that from *Shewanella pneumatophori* SCRC-2738. Namely, the EPA biosynthesis gene cluster of *S. marinintestina* IK-1 consisted of *pfaE*, *pfaA*, *pfaB*, *pfaC*, and *pfaD* in this order. To generate EPA-deficient mutants of *S. marinintestina* IK-1, insertional inactivation mutagenesis was performed targeting the *pfaD* gene. An insertional fragment of *pfaD* was amplified using primers of 5'-CGCTACCCAATGGCCCTTAC-3' and 5'-GCTGGCTCCAGCTTCAACAC-3' and cloned into the *EcoRV* site of mobilizable suicide pKNOCK-Cm [6]. The resulting *pfaD*: pKNOCK-Cm construct was introduced into *S. marinintestina* IK-1 from *E. coli* BW20767 (ATCC47084) by conjugal transfer as described by Fukuchi et al. [7]. Chloramphenicol-resistant exconjugants arose

from plasmid integration into the *S. marinintestina* IK-1 chromosome in a single crossover event yielding strain IK-1Δ8 with *pfaD* insertionally inactivated. The site of plasmid insertion was verified by PCR amplification of a portion of the *pfaD* gene using primers located upstream and downstream of the insertion site.

2.3. Analytical procedures

The carbonyl content in oxidatively modified cellular proteins was determined using cells treated with 0.01 mM H₂O₂, as previously described [8]. The degree of carbonylation was expressed as nmol of protein carbonyls per mg of protein. Protein was assayed by the Bradford method [9].

The intracellular and extracellular concentrations of H₂O₂ were measured with the method by González-Flecha and Demple [10,11]. The cells of *S. marinintestina* strains IK-1 and IK-1Δ8 that had been suspended in PBS at an OD₆₀₀ of 1.0, were treated with and without H₂O₂ and then separated as supernatants and cell pellets by centrifugation as previously described [2]. The extracellular concentration of H₂O₂ in the supernatants was measured using a titanium-based method [12]. The cell pellets were resuspended in the same buffer at the same concentration of cells and then allowed to stand for 30 min at 15 °C. This time is sufficient for the medium to reach the intracellular steady-state concentration of H₂O₂ on the basis of its free diffusion through the

cell membrane [10]. After centrifugation the intracellular concentration of H₂O₂ was measured by assaying the concentration in the supernatants.

Packed wet bacterial cells from 10 ml of cultures were subjected to methanolysis using 2 M methanolic HCl in the presence of 20 µg of heneicosanoic acid as an internal standard, as previously described [13]. The resulting fatty acid methyl esters were analyzed by gas-liquid chromatography, as previously described [13]. The amount of fatty acids was expressed as micrograms of their methyl esters per millilitre of culture.

To measure the dry cell weight the packed cells were freeze-dried and weighed. Total lipids from dry cells were extracted by the Bligh and Dyer method [14]. Phospholipids were separated by one-dimensional thin-layer chromatography of total lipids. Fatty acid composition and phosphorus contents of individual phospholipids were determined, as described previously [15].

The catalase activity of their cell-free extracts of *S. marinintestina* strains IK-1 and IK-1Δ8 was measured spectrophotometrically at 240 nm at room temperature, as described previously [16].

3. Results

3.1. Effects of H_2O_2 on growth of *S. marinintestina* strains IK-1 and IK-1 Δ 8

S. marinintestina strains IK-1 and IK-1 Δ 8 similarly grew in LB medium containing NaCl at 15 °C (data not shown). A similar finding has been observed for EPA-producing *Photobacterium profundum* SS9 and its EPA-deficient mutant [17]. To investigate their susceptibility to H_2O_2 , cells of these strains were subjected to bacteriostatic conditions, where the cells were suspended in PBS containing (H_2O_2+) or not containing (H_2O_2-) various concentrations of H_2O_2 and incubated for 30 min at 15 °C with no agitation. In Fig. 1 the number of CFUs of each strain under H_2O_2+ and H_2O_2- conditions was represented as a relative value (%) against the respective baseline under H_2O_2- conditions. CFUs for *S. marinintestina* strains IK-1 were much higher than those for strain IK-1 Δ 8 at any concentration of H_2O_2 (Fig. 1).

3.2. Effects of H_2O_2 on catalase activity and carbonyl contents in *S. marinintestina* strains IK-1 and IK-1 Δ 8

The catalase activity was almost the same in both *S. marinintestina* strains IK-1 and IK-1 Δ 8 and it was not affected by the presence of H_2O_2 (data not shown). It should be noted that the baseline catalase activity ($0.18 \text{ U mg protein}^{-1}$) of *S. marinintestina* IK-1 was remarkably low,

being only one tenth of that of *E. coli* and other bacteria such as *Bacillus subtilis* and *Vibrio parahaemolyticus* [16].

When bacterial cells are treated with H₂O₂, various proteins such as elongation factor G [18] or glyceraldehyde-3-phosphate dehydrogenase [19] are carbonylated, leading to a growth arrest. The baseline level of protein carbonyls in *S. marinintestina* strains IK-1 and IK-1Δ8 was approximately 80 nmol mg protein⁻¹ (Table 1). In *S. marinintestina* IK-1, levels of protein carbonyls did not change notably after incubation with 0.01 mM H₂O₂ for 30 min, while for strain IK-1Δ8, levels of protein carbonyl increased slightly from 83.1 ± 1.3 nmol mg protein⁻¹ to 94.1 ± 1.9 nmol mg protein⁻¹ for 30 min under H₂O₂+ conditions. However, no change was observed in either strain under H₂O₂- conditions.

3.3. Effects of H₂O₂ on amount of S. marinintestina strains IK-1 and IK-1Δ8 cells and their fatty acid composition

The baseline weight of dry cells from 1 ml of culture was 220 ± 30 μg and 190 ± 30 μg for *S. marinintestina* strains IK-1 and IK-1Δ8, respectively. When cells were treated with 0.01 mM H₂O₂ for 30 min a significant decrease in the dry cell weight per culture were observed only for strain IK-1Δ8 (Table 2). Little changes were observed in either strain under H₂O₂- conditions.

Similarly the amount of fatty acid extracted from H₂O₂-treated cells of *S. marinintestina* IK-1Δ8 was much lesser than that of *S. marinintestina* IK-1 (data not shown).

The major fatty acids in *S. marinintestina* IK-1 cells were dodecanoic acid (10.9 ± 1.1%), tetradecanoic acid (15.1 ± 0.3%), palmitoleic acid (22.4 ± 0.6%), and EPA (17.3 ± 0.4%). Under the same condition, no EPA was detected and levels of palmitoleic acid significantly increased to 49.0 ± 0.5% in cells of strain IK-1Δ8. Treating both strains with H₂O₂ had little effect on their fatty acid composition (data not shown).

S. marinintestina strains IK-1 and IK-1Δ8 had phosphatidylethanolamine (PE) and phosphatidylglycerol (PG), accounting for approximately 70 mol% and 25 mol%, respectively, of total lipids. In strain IK-1 all of EPA was detected in PE and PG.

3.4. Extracellular and intracellular concentrations of H₂O₂ in S. marinintestina strains IK-1 and IK-1Δ8

The extracellular and intracellular concentrations of H₂O₂ were also determined for *S. marinintestina* strains IK-1 and IK-1Δ8 using cells incubated under bacteriostatic conditions in the presence of 0.1 mM H₂O₂ and where no H₂O₂ was present. Cells were treated with 0.1 mM H₂O₂ in consideration of the detection limit of H₂O₂ by the titanium-based method [12]. The extracellular levels of H₂O₂ for both strains decreased slightly and gradually over time (Fig. 2A).

The rate of decrease was essentially the same for cells of *S. marinintestina* strains IK-1 and IK-1 Δ 8. Negligible concentrations of H₂O₂ were consistently detected in cells of both strains where no H₂O₂ was present (Fig. 2A). The intracellular concentration of H₂O₂ increased over time for both strains under H₂O₂ conditions. The H₂O₂ concentrations were 20.8 ± 4.2 μ M for strain IK-1 and 39.4 ± 3.1 μ M for strain IK-1 Δ 8 cells, 30 min after the addition of H₂O₂ (Fig. 2B). The intracellular concentration of H₂O₂ in cells of *S. marinintestina* strain IK-1 was approximately one-quarter of that in cells of strain IK-1 Δ 8.

4. Discussion

In this study, the antioxidative function of EPA was demonstrated in the marine bacterium *S. marinintestina* IK-1 that inherently produces EPA and in its EPA-deficient mutant (strain IK-1 Δ 8). *S. marinintestina* IK-1 produced EPA at approximately 17% of total fatty acids at 15 °C. *S. marinintestina* IK-1 completely lost the capacity to synthesize EPA after the insertional inactivation mutagenesis of the *pfaD* gene. The lack of EPA in strain IK-1 Δ 8 was maintained after repeated cultivations in a medium without antibiotics.

Previously we presented data based on *E. coli* recombinant systems showing that EPA in the cell membrane exerted an antioxidative effect on exogenously added H₂O₂. Although the molecular mechanism of this antioxidative function of EPA is unknown, it has been speculated

previously that the inherent molecular properties of EPA (as acyl constituents of phospholipids) allow it to form a more highly packed structure [20] than most other phospholipids so that it can inhibit the entry of H₂O₂ through the cell membrane. Considering that polyunsaturated fatty acids like EPA and docosahexaenoic acid are cellular molecules with the greatest susceptibility to oxidation [21], this shielding function of EPA against exogenous H₂O₂ is a unique mechanism. This mechanism operated against other oxidants such as *t*-butyl hydroperoxide (analog of hydrogen peroxide and not degraded by catalase) (unpublished result).

The ability of EPA to shield the cell membrane against exogenous H₂O₂ was also present in bacteria that naturally produced EPA. The CFU counting method (Fig. 1) showed that *S. marinintestina* IK-1 that produced EPA was more resistant to H₂O₂ than its EPA-deficient mutant strain IK-1Δ8. The higher resistance of *S. marinintestina* IK-1 to H₂O₂ was exemplified by the finding that lower levels of protein carbonyls were detected in *S. marinintestina* IK-1 (with EPA) than in strain IK-1Δ8 (without EPA) (Table 1), and that the recovered amount of cells (dry cell weight) and then simultaneously that of fatty acids was much higher from cultures of strain IK-1 than from cultures of strain IK-1Δ8 (Table 2). The latter observation suggests that more significant cell breakage occurred in strain IK-1Δ8 than in strain IK-1 and that EPA in cell membrane phospholipids function to prevent the entry of H₂O₂ from the external medium. The cell breakage was at a level of approximately 30% in strain IK-1Δ8 that had been treated with 0.01% H₂O₂

(Table 2), where CFU of this strain was much more significantly reduced (approximately 70%; Fig. 1). These results suggest that a viable but non-culturable state might have been induced in some of H₂O₂-treated strain IK-1Δ8 cells, as was the case of H₂O₂-treated *E. coli* [1]. Although the shielding mechanism of the cell membrane is largely unknown, we speculate the involvement of the structural hindrance of EPA-containing phospholipids, whose structure is more compact and prohibits the entry of H₂O₂ from outside.

Levels of protein carbonyls in nongrowth cells of *S. marinintestina* strains IK-1 and IK-1Δ8 before treatment with H₂O₂ were approximately 80 nmol mg protein⁻¹. Only slightly higher values were obtained for the mutant cells (Table 1). These values were significantly higher than those measured in catalase-deficient *E. coli* UM2 transformant cells, both with EPA and without EPA, in which levels of protein carbonyls determined under the same conditions were approximately 0.2 nmol mg protein⁻¹. The value was increased to approximately 0.5 nmol mg protein⁻¹ and 0.75 nmol mg protein⁻¹ for cells with EPA and those without EPA, respectively [2]. This was despite *E. coli* UM2 having no catalase or other notable H₂O₂-decomposing activity and *E. coli* UM2 transformants were treated with 0.3 mM H₂O₂, whereas *S. marinintestina* strains IK-1 and IK-1Δ8 were treated with 0.01 mM H₂O₂.

All of these observations suggest that the cellular proteins of *S. marinintestina* strains might be much more susceptible to oxidation than the cellular proteins of *E. coli*. It is known that

the marine bacteria *Vibrio rumoiensis* and *V. parahaemolyticus* are considerably more sensitive to exogenous H₂O₂ than *E. coli* and *B. subtilis* [22]. However, it is unknown whether such a characteristic is common in marine bacteria. To determine this would require biochemical and molecular testing of the susceptibility of proteins to oxidants.

Reference

- [1] Nishida, T., Oriyasa, Y., Ito, Y., Yu, R., Yamada, A., Watanabe, K. and Okuyama, H. (2006) *Escherichia coli* engineered to produce eicosapentaenoic acid becomes resistant against oxidative damages. FEBS Lett. 580, 2731–2735.
- [2] Nishida, T., Oriyasa, Y., Watanabe, K. and Okuyama, H. (2006) The cell membrane-shielding function of eicosapentaenoic acid for *Escherichia coli* against exogenously added hydrogen peroxide. FEBS Lett. 580, 6690–6694.
- [3] Okuyama, H., Oriyasa, Y., Nishida, T., Watanabe, K. and Morita, N. (2007) Bacterial genes responsible for the biosynthesis of eicosapentaenoic and docosahexaenoic acids and their heterologous expression. Appl. Environ. Microbiol. 73, 665–670.
- [4] Hirota, K., Nodasaka, Y., Oriyasa, Y., Okuyama, H. and Yumoto, I. (2005) *Shewanella pneumatophori* sp. nov., an eicosapentaenoic acid-producing marine bacterium isolated from

- the intestines of pacific mackerel (*Pneumatophorus japonicus*). Int. J. Syst. Evol. Microbiol. 55, 2355–2359.
- [5] Satomi, M., Oikawa, H. and Yano, Y. (2003) *Shewanella marinintestina* sp. nov., *Shewanella schlegeliana* sp. nov. and *Shewanella sairae* sp. nov., novel eicosapentaenoic-acid-producing marine bacteria isolated from sea-animal intestines. Int. J. Syst. Evol. Microbiol. 53, 491–499.
- [6] Alexeyev, MF. (1999) The pKNOCK series of broad-host-range mobilizable suicide vectors for gene knockout and targeted DNA insertion into the chromosome of Gram-negative bacteria. BioTechniques 26, 824–828.
- [7] Fukuchi, J., Sato, T., Kato, C., Ito, M. and Horikoshi, K. (2002) The host-vector system for deep-sea piezophilic bacteria, *Shewanella violacea* DSS12 and *Moritella japonica* DSK-1. JAMSTECR 46, 157–161. (in Japanese)
- [8] Levine, R., Garland, D., Oliver, C.N., Amichi, A., Climent, I., Lenz, A-G. , Ahn, B-W, Shaltiel, S. and Stadtman, R. (1990) Determination of carbonyl content in oxidatively modified proteins. Methods Enzymol. 186, 464–478.
- [9] Bradford, M.M. (1976) A rapid and sensitive method for quantitation of microgram quantities of protein utilizing the principle of protein-dye binding. Anal. Biochem. 14, 248–254.

- [10] González-Flecha, B. and Demple, B. (1994) Intracellular generation of superoxide as a by-product of *Vibrio harveyi* luciferase expressed in *Escherichia coli*. *J. Bacteriol.* 176, 2293–2299.
- [11] González-Flecha, B. and Demple, B. (1997) Homeostatic regulation of intracellular hydrogen peroxide concentration in aerobically growing *Escherichia coli*. *J. Bacteriol.* 179, 382–388.
- [12] Pobiner, H. (1961) Determination of hydrocarbon by conversion to hydrogen peroxide and measurement by titanium complexing. *Anal. Chem.* 33, 1423–1426.
- [13] Orikasa, Y., Nishida, T., Hase, A., Watanabe, K., Morita N. and Okuyama, H. (2006) A phosphopantetheinyl transferase gene essential for biosynthesis of n-3 polyunsaturated fatty acids from *Moritella marina* strain MP-1. *FEBS Lett.* 580, 4423–4429.
- [14] Bligh, E.G. and Dyer, W.J. (1959) A rapid method of total lipid extraction and purification. *Can. J. Biochem. Physiol.* 37, 911–917.
- [15] Okuyama, H., Itoh, H, Fukunaga, N. and Sasaki, S. (1984) Separation and properties of the inner and outer membranes of a psychrophilic bacterium, *Vibrio* sp. strain ABE-1. *Plant Cell Physiol.* 25,1255–1264
- [16] Yumoto, I., Yamazaki, K., Kawasaki, K., Ichise, N., Morita, N., Hoshino, T. and Okuyama, H. (1999) Isolation of *Vibrio* sp. S-1 exhibiting extraordinarily high catalase activity. *J. Ferment. Bioeng.* 85, 113–116.

- [17] Allen, E.E., Facciotti, D. and Bartlett, D.H. (1999) Monounsaturated but not polyunsaturated fatty acids are required for growth of the deep-sea bacterium *Photobacterium profundum* SS9 at high pressure and low temperature. *Appl. Environ. Microbiol.* 65: 1710–1720.
- [18] Tamarit, J., Cabisco, E. and Ros, J. (1998) Identification of the major oxidatively damaged proteins in *Escherichia coli* cells exposed to oxidative stress. *J. Biol. Chem.* 273, 3027–3032.
- [19] Weber, H., Engelmann, S., Becher, D. and Hecker, M. (2004) Oxidative stress triggers thiol oxidation in the glyceraldehyde-3-phosphate dehydrogenase of *Staphylococcus aureus*. *Mol. Microbiol.* 52, 133–140.
- [20] Rajamoorthi, K., Petrache, H.I., McIntosh, T.J. and Brown, M.F. (2005) Packing and viscoelasticity of polyunsaturated omega-3 and omega-6 lipid bilayers as seen by ²H NMR and X-ray diffraction. *J. Am. Chem. Soc.* 127, 1576–1588.
- [21] Halliwell, B. and Gutteridge, J.M.C. (1998) *Free radicals in biology and medicine*. 3rd ed. Oxford Univ. Press, Oxford.
- [22] Ichise, N., Morita, N., Hoshino, T., Kawasaki, K., Yumoto, I. and Okuyama, H. (1999) A mechanism of resistance to hydrogen peroxide in *Vibrio rumoiensis* S-1. *Appl. Environ. Microbiol.* 65, 73-79.

Table 1. Effects of H₂O₂ on the protein carbonyl contents of *Shewanella marinitestina* IK-1^a and its EPA-deficient mutant IK-1Δ8^a

	Protein carbonyl (nmol mg protein ⁻¹)		
	H ₂ O ₂ -		H ₂ O ₂ +
	0 ^b	30 ^b	30 ^b
IK-1	78.6 ± 0.9	82.2 ± 5.5	84.1 ± 5.0
IK-1Δ8	83.1 ± 1.3	84.6 ± 3.0	94.1 ± 1.9

^aCells were treated with 0.01 mM H₂O₂.

^bTime (min)

Table 2. Effects of H₂O₂ on the dry cell weight of *Shewanella marinitestina* IK-1 and its EPA-deficient mutant IK-1Δ8 that had been treated with and without H₂O₂ under bacteriostatic conditions

	Dry cell weight (μg ml cell culture ⁻¹)			
	IK-1		IK-1Δ1	
	H ₂ O ₂ -	H ₂ O ₂ +	H ₂ O ₂ -	H ₂ O ₂ +
0 time	220 ± 30 (100.0)		190 ± 30 (100.0)	
30 min	210 ± 0.0 (95.4)	210 ± 10 (94.5)	200 ± 20 (103.1)	140 ± 20 (71.2)

Figure legends

Fig. 1. Effects of H₂O₂ on the colony-forming ability of *Shewanella marinintestina* IK-1 and its EPA-deficient mutant IK-1Δ8. Cells of strain IK-1 (●) and strain IK-1Δ8 (▲) were suspended in phosphate-buffered saline. After incubated for 30 min in the presence of 0.01, 0.1, 1, and 10 mM H₂O₂ under nongrowth conditions, the number of CFUs was determined. The baseline value of each strain was that of cells treated without H₂O₂. The CFU of *S. marinintestina* strains IK-1 and IK-1Δ8 was $7.5 \pm 0.3 \times 10^9$ ml of culture⁻¹ and $7.2 \pm 0.3 \times 10^9$ ml of culture⁻¹, respectively, at each baseline.

Fig. 2. Changes of extracellular (A) and intracellular (B) concentrations of H₂O₂ in *Shewanella marinintestina* strains IK-1 and IK-1Δ8 cells that were treated with and without H₂O₂ under nongrowth conditions. Portions of cell suspensions were withdrawn at various time intervals and assayed to determine extracellular and intracellular H₂O₂ concentrations. Δ, strain IK-1Δ8 under H₂O₂⁻ conditions; ▲, IK-1Δ8 under H₂O₂⁺ conditions; ○, IK-1 under H₂O₂⁻ conditions; ●, IK-1

under H₂O₂+ conditions. Cells were incubated at 15 °C. The data indicated are means ± standard errors for three independent experiments.

Fig. 1.





Fig. 2.

A



B

