



Title	Childhood epidermolysis bullosa acquisita with autoantibodies against the noncollagenous 1 and 2 domains of type VII collagen : case report and review of the literature
Author(s)	Mayuzumi, M.; Akiyama, M.; Nishie, W.; Ukae, S.; Abe, M.; Sawamura, D.; Hashimoto, T.; Shimizu, H.
Citation	British Journal of Dermatology, 155(5), 1048-1052 https://doi.org/10.1111/j.1365-2133.2006.07443.x
Issue Date	2006-11
Doc URL	http://hdl.handle.net/2115/30232
Rights	The definitive version is available at www.blackwell-synergy.com
Type	article (author version)
File Information	BJD155-5.pdf



[Instructions for use](#)

British Journal of Dermatology

Manuscript No. BJD-2005-01478 Revised Version

Case Report

Childhood epidermolysis bullosa acquisita with autoantibodies against the non-collagenous 1 and 2 domains of type VII collagen: a case report and a review of the literature

M. MAYUZUMI^{1,2}, M. AKIYAMA¹, W. NISHIE^{1,2}, S. UKAE³, M. ABE¹,
D. SAWAMURA¹, T. HASHIMOTO⁴ and H. SHIMIZU¹

¹Department of Dermatology, Hokkaido University Graduate School of Medicine, Sapporo, Japan;

Departments of ²Dermatology and ³Pediatrics, Sapporo Social Insurance General Hospital, Sapporo, Japan;

⁴Department of Dermatology, Kurume University School of Medicine, Kurume, Japan

Short title: Childhood EBA

Key words: EBA, epitope, immunoblot, immunofluorescence, inflammatory phenotype

Correspondence

Masashi Akiyama

Department of Dermatology,

Hokkaido University Graduate School of Medicine,

N15 W7, Sapporo 060-8648, Japan

e-mail: akiyama@med.hokudai.ac.jp

Telephone: 81-11-716-1161 ext. 5962

Fax: 81-11-706-7820

Summary

Epidermolysis bullosa acquisita (EBA) is an acquired subepidermal bullous disease characterized by IgG autoantibodies to type VII collagen, a major component of anchoring fibrils. Most patients with EBA are adult and develop autoantibodies to the non-collagenous 1 (NC-1) domain of type VII collagen. In this report, we describe a 4-year-old Japanese boy presenting pruritic vesicles and tense blisters over his whole body. Immunofluorescence studies revealed linear IgG/C3 deposits along the dermal-epidermal junction of the patient's skin, and circulating IgG autoantibodies mapping to the dermal side of 1M NaCl-split skin. By immunoblotting analysis using dermal extracts as a substrate, the patient's IgG antibodies labeled a 290 kD protein corresponding to type VII collagen. Immunoblotting studies using recombinant proteins demonstrated that the patient's circulating autoantibodies recognized not only the NC-1 but also the non-collagenous 2 (NC-2) domain of type VII procollagen. Review of the previously reported cases and the present case suggested that EBA cases with autoantibodies to regions other than the NC-1 domain are all children younger than 10 years of age with clinical features of an inflammatory phenotype.

Introduction

Epidermolysis bullosa acquisita (EBA) is an autoimmune blistering skin disease which has circulating IgG autoantibodies to type VII collagen, a major component of anchoring fibrils¹, composed of three identical alpha chains. Each chain consists of a long central triple-helical collagenous domain flanked by a large amino-terminal 145 kD non-collagenous 1 (or NC1) domain and a smaller carboxyl-terminal 20 kD non-collagenous 2 (NC2) domain. Type VII procollagen molecules form anti-parallel dimers that are stabilized by disulfide bonding at the carboxyl terminus, and a portion of the NC2 domain is removed by specific proteolytic cleavage to yield the mature type VII collagen. Several dimers aggregate laterally to form the unique cross-banded structure, i.e. anchoring fibrils, which comprise anti-parallel dimers and contain NC1 domains at both ends, locating in the lamina densa and forming semi-circular loops visible by the electron microscope².

Two distinct phenotypes of EBA have been described; the classical non-inflammatory type and the inflammatory type. In a large number of the adults with clinically classical type of EBA, it has previously been recognized that autoantibodies are directed against epitopes within the NC1 domain, whereas no immunoreactivity with the NC2 or the triple helical domain was detected³. However, recently, in five children with the inflammatory subtype of EBA, novel variants with reactivity to the NC2 or/and the triple helical domains have been reported⁴⁻⁶. These findings indicated that some distinct immunopathological differences between childhood EBA and those occurred in adulthood could be present.

In this report, we describe a case of EBA in a 4-year-old Japanese boy and have reviewed previous reported cases of childhood EBA.

Case Report

A 4-year-old Japanese boy, generally in good health, developed painful oral blisters and erosions. A few weeks later, widespread pruritic vesicles and tense blisters appeared over almost his entire body, but especially on his scrotum and buccal mucosa. The blisters were seen both on erythematous bases and on normal skin, and showed an annular arrangement in some regions (Fig.1). His nails were not affected. He had no family history of any blistering disorders.

General laboratory examinations including full blood count, hepatic and renal function tests, urinalysis and C-reactive protein were within normal limits except for an increased number of eosinophils (13.5 %) in his peripheral blood. Anti-nuclear antibodies were negative. Anti-herpes simplex virus antibodies were not detected, either. Viral culture of vesicular fluid and a Tzanck smear test on a vesicular base were all negative.

Histopathological examination of skin biopsy specimens from the dorsal side of his foot revealed subepidermal blisters with an inflammatory infiltrate of lymphocytes, neutrophils, and eosinophils in the papillary and superficial reticular dermis. Especially, neutrophils and nuclear dusts were seen at the tips of the edematous dermal papillae (Fig.2).

On direct immunofluorescence (IF) studies, linear deposits of IgG and C3, but no IgA were seen along the dermal-epidermal junction of the patient's skin biopsy specimen. Indirect IF studies using a 1M NaCl-split normal human skin as a substrate showed IgG deposits on the dermal side of the artificial split at a titer of 1:160 (Fig.3). Circulating autoantibodies to the NC16a domain of 180 kD bullous pemphigoid antigen (BP180) were not detected using a BP 180 enzyme-linked immunosorbent assay (ELISA)

kit (MBL, Naka-ku, Nagoya, Japan). Immunoblot analysis demonstrated that the patient's sera reacted both with a 290 kD protein in dermal extracts and recombinant type VII collagen (Figs. 4a, b). In addition to the 290 kD band, many additional bands were seen in the immunoblot analysis with patient sera using dermal extract as a substrate. These additional bands were thought to reflect reactivity of patient's sera to background, degradation products because, in this immunoblot analysis, we used high concentration sera in order not to overlook 200 kD band in p200 pemphigoid. Epitope analyses with type VII pro-collagen recombinant fragments revealed that the patient's sera recognized both of the NC1 and NC2 domains (Fig.4c).

The boy was treated with oral prednisolone (1.1mg/kg per day) for 2 weeks, which inhibited new blister formation. When the dose of prednisolone was reduced to 0.4mg/kg per day, new blister formation was observed. Oral dapsone 1.5mg/kg per day was added, but on the next day, erythematous macules appeared on his upper trunk. Thus, dapsone was discontinued and the dose of prednisolone was increased to 0.7mg/kg per day. During the next 2 months, the patient kept taking oral prednisolone 0.5mg/kg per day, and the blisters steadily healed, leaving hyper- and hypo-pigmented macules and milia.

Discussion

EBA is a relatively uncommon disease with an incidence about 10 times less than that of bullous pemphigoid (BP). EBA was first defined with following characteristic features: trauma-induced bullae that heal with milia and scar mainly over the joints of the hands, elbows, feet and knees; nail dystrophy; typically adult onset; a negative family history of

epidermolysis bullosa; and exclusion of other bullous diseases on the basis of clinical and laboratory evidence. The definitive features now include histological and immunopathological findings, such as subepidermal blisters; circulating autoantibodies against skin basement membrane zone antigens, binding to the dermal side of NaCl-split skin; IgG/C3 deposits in the anchoring fibril zone; and the presence of anti-type VII collagen antibodies that are identified as a 290 kD band by immunoblot analysis on dermal extracts¹. Among clinically diagnosed EBA patients, antibodies can be detected by immunoblotting with dermal extracts in approximately 30% of the cases⁷, and by indirect IF using NaCl-split skin as a substrate in approximately 50% of the cases⁸.

Two distinct phenotypes of EBA have been described, i.e. the classical (non-inflammatory) type and inflammatory type. The classical type presents with marked skin fragility, blisters and erosions at sites of trauma, and healing with scarring and milia. The inflammatory type can mimic almost all other chronic bullous diseases, and its clinical differentiation from BP, cicatricial pemphigoid and linear IgA bullous dermatosis may be difficult^{4-6, 9-26}. In some patients, characteristics of both classical and inflammatory phenotypes of EBA have been observed⁹, but most patients with EBA in adulthood appear to suffer from the classical type of EBA.

Conversely, patients with childhood EBA (16-years-old or younger) are very rare. To our knowledge, there are 33 cases of childhood EBA including the present case reported in the literature^{4-6, 9-26} Among them, six cases were Japanese and the others were non-Japanese including at least five Caucasian and five African. These patients presented mainly with the inflammatory subtype. Some differences exist between EBA in adults and children. In the adult, the occurrence of mucosal involvement was seen in

approximately 50 % of patients, while in the children, mucosal lesions were seen in the vast majority of the reported cases^{9,18}. The adult form of EBA is known to be difficult to treat, requiring high doses of prednisolone, dapsons, immunosuppressive agents, plasmapheresis, etc. In contrast, dapsons and low dose prednisolone are usually effective in treatments for childhood EBA. The prognosis in childhood EBA seems to be much better than adult cases^{4-6,9-26}.

The major epitopes of circulating autoantibodies in adult patients with the classical type of EBA are known to be located within the NC1 domain, but neither in the NC2 nor in the triple helical domains³. However in five cases of childhood EBA (four Japanese children and one European child) reported in the literature, autoantibodies have been found to recognize epitopes on the NC2 and/or the triple helical domain over the last years (Table 1)⁴⁻⁶. These five patients and the present case were all young children less than 10 years of age, who all showed similar clinical features of the inflammatory subtype of EBA, and complete clearing or a relatively good response to treatments. Interestingly, two of these children also demonstrated autoantibody reactivity towards the 230kD BP antigen (BP230) and the NC16A domain of BP180⁴.

Schmidt *et al.*⁵ mentioned this phenomenon and referred it as “epitope spreading”. Epitope spreading sometimes occurs in other adult human autoimmune skin disorders, and has been observed in a few cases of EBA, specifically in young children⁵. The molecular events in epitope spreading remain to be elucidated, but it is supposed that the skin damage from autoimmune or inflammatory processes subsequently could induce autoimmunity to a sequestered or closely related antigen or epitope.

Five out of the six EBA patients with autoantibodies to the NC2 and/or

the triple helical domains are Japanese children. Tanaka *et al.*⁴ suggested the phenomenon may be unique to a particular ethnic group. To confirm this hypothesis, detailed epitope mappings in bullous dermatosis cases with similar clinical and pathological features will be required, especially in children from different ethnic groups.

Further studies on autoantibodies in larger series of patients with EBA are needed to clarify the correlation between the presence of certain autoantibodies against specific subdomains of type VII collagen and their clinical features, such as classical or inflammatory phenotype, the patient's age of onset, clinical courses, and the prognosis.

References

- 1) Woodley DT, Briggaman RA, O'Keefe EJ et al. Identification of the skin basement-membrane autoantigen in epidermolysis bullosa acquisita. *N Engl J Med* 1984; 310: 1007-13
- 2) Shimizu H, Ishiko A, Masunaga T et al. Most anchoring fibrils in human skin originate and terminate in the lamina densa. *Lab Invest* 1997; 76: 753-63
- 3) Lapiere J-C, Woodley DT, Parente MG et al. Epitope mapping of type VII collagen: identification of discrete peptide sequences recognized by sera from patients with acquired epidermolysis bullosa. *J Clin Invest* 1993; 92: 1831-9
- 4) Tanaka H, Ishida-Yamamoto A, Hashimoto T et al. A novel variant of acquired epidermolysis bullosa with autoantibodies against the central triple-helical domain of type VII collagen. *Lab Invest* 1997; 77: 623-32
- 5) Schmidt E, Hopfner B, Chen C et al. Childhood epidermolysis bullosa acquisita: a novel variant with reactivity to all three structural domains of type VII collagen. *Br J Dermatol* 2002; 147: 592-7
- 6) Fukumoto T, Umemura T, Higuchi M. et al. Childhood epidermolysis bullosa acquisita with autoantibodies against all 3 structural domains of type VII collagen. *J Am Acad Dermatol* 2004; 50(3): 480-2
- 7) Chen M, Chan LS, Cai X et al. Development of an ELISA for rapid detection of anti-type VII collagen autoantibodies in epidermolysis bullosa acquisita. *J Invest Dermatol* 1997; 108: 68-72
- 8) Gammon WR, Fine JD, Forbes M, Briggaman RA. Immunofluorescence on split skin for the detection and differentiation of basement membrane zone autoantibodies. *J Am Acad Dermatol* 1992; 27: 79-87
- 9) Stewart MI, Woodley DT, Briggaman RA. Epidermolysis bullosa

- acquisita and associated symptomatic esophageal webs. *Arch Dermatol* 1991; 127: 373-7
- 10) Borok M, Heng MCY, Ahmed AR et al. Epidermolysis bullosa acquisita in an 8-year-old girl. *Pediatric Dermatol* 1986; 3: 315-22
 - 11) Rubenstein R, Esterly NB, Fine JD. Childhood epidermolysis bullosa acquisita. *Arch Dermatol* 1987; 123: 772-6
 - 12) McCuaig CC, Chan LS, Woodley DT et al. Epidermolysis bullosa acquisita in childhood. *Arch Dermatol* 1989; 125: 944-9
 - 13) Espagne E, Prost C, Janssen F et al. Epidermolyse bulleuse acquise de l'enfant. A propos d'un cas. *Ann Dermatol Venereol* 1990; 117: 814-6
 - 14) Arpey CJ, Elewski BE, Moritz DK, Gammon WR. Childhood epidermolysis bullosa acquisita. Report of three cases and review of literature. *J Am Acad Dermatol* 1991; 24: 706-14
 - 15) Roger H, Machado P, Nicolas JF et al. Epidermolysis bullosa acquisita in a 3¹/₂-year-old girl. *J Am Acad Dermatol* 1992; 27: 858-62
 - 16) Kirtschig G, Wojnarowska F, Marsden RA et al. Acquired bullous disease of childhood: re-evaluation of diagnosis by indirect immunofluorescence examination on 1M NaCl split skin and immunoblotting. *Br J Dermatol* 1994; 130: 610-6
 - 17) Lacour JP, Bernard P, Rostain G et al. Childhood acquired epidermolysis bullosa. *Pediatric Dermatol* 1995; 12: 16-20
 - 18) Caux F, Kirtschig G, Lemarchand-Venencie et al. IgA-epidermolysis bullosa acquisita in a child resulting in blindness. *Br J Dermatol* 1997; 137: 270-5
 - 19) Callot-Mellot C, Bodemer C, Caux F et al. Epidermolysis bullosa acquisita in childhood. *Arch Dermatol* 1997; 133: 1122-6
 - 20) Feliciani C, Fasciocco D, Franchi A, Amerio P. Epidermolysis bullosa

- acquisita in a 10-year-old Alabian boy. *Br J Dermatol* 1997; 137: 307-8
- 21) Park SB, Cho KH, Youn JI et al. Epidermolysis bullosa acquisita in childhood- a case mimicking chronic bullous dermatosis of childhood. *Clin Exp Dermatol* 1997; 22: 220-2
- 22) Hayashi K, Ueda M, Sakai M et al. Epidermolysis bullosa acquisita in a 6-year-old Japanese boy. *Int J Dermatol* 1998; 37: 612-4
- 23) Edwards S, Wakelin SH, Wojnarowska F et al. Bullous pemphigoid and epidermolysis bullosa acquisita. *Pediatric Dermatol* 1998; 15: 184-90
- 24) Chorzelski T, Karczewska K, Dyduch A et al. Epidermolysis bullosa acquisita in a 4-year-old boy. *Pediatric Dermatol* 2000; 17: 157-8
- 25) Trigo-Guzman FX, Conti A, Aoki V et al. Epidermolysis bullosa acquisita in childhood. *J Dermatol* 2003; 30: 226-9.
- 26) Parodi A, Cozzani C, Verrini A et al. Epidermolysis bullosa acquisita in a child:an additional case. *Int J Dermatol* 2005; 44: 262-3

Figure legends

Fig 1. Clinical features. Disseminated, tense vesicles, bullae and erosions with crusts were seen over erythematous plaques on the face (a), arm (b), genital area (c) and the buttocks (d).

Fig 2. Histopathology. A subepidermal blister with an inflammatory infiltrate of lymphocytes together with eosinophils and neutrophils in the papillary dermis. (Original magnification $\times 125$).

Fig 3. (a) By direct IF staining, linear IgG deposits were seen along the dermal-epidermal junction in the patient's skin.
(b) Indirect IF using 1M NaCl-split human skin as a substrate revealed linear IgG deposits along the dermal side of the artificial split. (• dots: Roof side of the split skin).

Fig 4. (a) Immunoblot analysis of patient's circulating autoantibodies on normal human dermal extracts. *Arrows* indicate the 290kD EBA antigen and 200kD anti-p200 pemphigoid antigen. Control EBA serum (*lane 1*) and patient's serum (*lane 3*) reacted with EBA antigen, whereas control anti-p200 serum (*lane 2*) showed no 290kD band. (b) Immunoblot analysis of patient's circulating autoantibodies on recombinant type VII collagen. *Arrow* indicates the 290kD EBA antigen. Control EBA serum (*lane 1*) and patient's serum (*lane 3*) reacted with EBA antigen, whereas normal control serum (*lane 2*) showed no reactivity. (c) Immunoblot analyses using recombinant fusion proteins of NC1 and NC2 domains of type VII collagen. *Arrows* indicate respective positions of fusion proteins. Both control EBA serum (*lane 1*) and patient's serum (*lane 2*) reacted with NC1 and with

NC2.

Table 1. Cases with childhood EBA in which epitopes for autoantibodies were detected by immunoblot analysis

Case	Age/Race/Sex	DIF (BMZ)	IgG deposits in NaCl split skin IIF	Immunoblot on dermal extracts	Immunoblot on epidermal extracts	Recombinant protein immunoblot analysis				NC16a ELISA	Therapy	Reference
						NC1	TH	NC2	rBP180			
1	8/Japanese/F	IgG, IgA, C3	Dermal side	290kD (+)	230kD(+)	-	+	-	ND	-	PSL+DDS	4
2	1/Japanese/M	IgG, C3	Dermal side	290kD (+)	no reactivity	-	+	+	ND	+	Methylprednisolone+DDS	4
3	2/Japanese/F	IgG, C3	Dermal side	290kD (+)	no reactivity	-	+	+	ND	-	PSL	4
4	4/European/F	IgG, C3	Dermal side	290kD (+)	no reactivity	+	+	+	-	ND	PSL+DDS	5
5	5/Japanese/M	IgG, C3	Dermal side	290kD (+)	no reactivity	+	+	+	ND	-	Betamethasone	6
6	4/Japanese/M	IgG, C3	Dermal side	290kD (+)	ND	+	ND	+	ND	-	PSL	Present case

DIF: direct IF, IIF: indirect IF, BMZ: basement membrane zone, TH: triple helical collagenous domain, NC16a: NC16a domain of BP180, rBP180: recombinant BP180, ND: not done

PSL: prednisolone, DDS: diaminodiphenylsulfone







