



Title	INDUCTION OF PROTECTIVE IMMUNITY IN MICE AGAINST HETEROLOGOUS AND HOMOLOGOUS CHALLENGE INFECTION OF ANGIOSTRONGYLUS COSTARICENSIS AND A. SIAMENSIS BY LARVAL INFECTION AND BY ADULT WORM TRANSPLANTATION
Author(s)	KANEKO, Satoko
Citation	Japanese Journal of Veterinary Research, 36(2), 158-158
Issue Date	1988-05-20
Doc URL	http://hdl.handle.net/2115/3102
Type	bulletin (article)
File Information	KJ00002377086.pdf



[Instructions for use](#)

INDUCTION OF PROTECTIVE IMMUNITY IN MICE AGAINST
HETEROLOGOUS AND HOMOLOGOUS CHALLENGE INFECTION OF
ANGIOSTRONGYLUS COSTARICENSIS AND *A. SIAMENSIS* BY LARVAL
INFECTION AND BY ADULT WORM TRANSPLANTATION

Satoko KANEKO

*Department of Parasitology,
Faculty of Veterinary Medicine,
Hokkaido University, Sapporo 060, Japan*

Angiostrongylus costaricensis and *A. siamensis* are pathogenic in mice because of severe histopathological changes produced even in cases parasitized by a small number of worms. In a preliminary experiment to induce immunity, it was observed in ICR mice that treatment with 200 or 400 mg/kg of thiabendazole on days 6 to 10 PI (postinfection) was not very effective in removing the worms. However, the worms were completely removed when the mice were given 80 mg/kg of levamisole on days 4 to 6 PI.

After oral inoculation with 150L₃ (third stage larvae) of either *A. siamensis* or *A. costaricensis*, BDF₁ mice were treated with levamisole, and then challenged *per os* heterologously and homologously with 100L₃. Twelve days after the challenge infection, the reduction rate of worms recovered from immunized mice as compared with the non-immunized control mice were 95% and 98% in homologous infection, and 78% and 90% in heterologous challenge infections for *A. siamensis* and *A. costaricensis*, respectively. This showed that acquired resistance against the homologous species was stronger than that against the heterologous species.

Immunity by intraperitoneal transplantation of 10 adult worms resulted in 47% and 31% worm reduction rate in homologous, and 21% and 28% in heterologous challenge infections for *A. siamensis* and *A. costaricensis* respectively, as compared with the non-immunized control mice.

Therefore, the acquired resistance induced by oral inoculation of L₃ was shown to be stronger than that induced by intraperitoneal transplantation of the adult worms.