



Title	ESTABLISHMENT AND CHARACTERIZATION OF A THYMIDINE-KINASE DEFICIENT CELL LINE FROM JAPANESE QUAIL CELL LINE, QT35
Author(s)	NARITA, Takahiro
Citation	Japanese Journal of Veterinary Research, 37(2), 126-126
Issue Date	1989-06-20
Doc URL	http://hdl.handle.net/2115/3168
Type	bulletin (article)
File Information	KJ00002377271.pdf



[Instructions for use](#)

ESTABLISHMENT AND CHARACTERIZATION OF A THYMIDINE-KINASE
DEFICIENT CELL LINE FROM JAPANESE QUAIL CELL LINE, QT35.

Takahiro NARITA

*Department of Epizootiology
Faculty of Veterinary Medicine
Hokkaido University, Sapporo 060, Japan*

Thymidine-kinase deficient (TK⁻) cell lines of several mammalian cells have been established, and recombinant viruses that have foreign genes within the viral TK gene have been constructed by using the TK⁻ cell lines. In the present study, I established a TK⁻ cell line (QT-TK⁻ cells) from Japanese quail cell line QT35, and examined the biological characteristics of QT-TK⁻ cells for use in the selection of TK⁻ avian viruses. A TK⁻ mutant cell line, QT-TK⁻, which was resistant to 100 μ g/ml bromodeoxyuridine (BUdR) was established by multi-step selection with BUdR. The QT-TK⁻ cells could not grow in HAT medium. The uptake of ³H-thymidine, the TK activity in the cell extract, and the rate of cells that incorporated BUdR of QT-TK⁻ cells were significantly decreased to 0.3%, 0.5%, and 0.6% respectively, when compared with those of QT35 cells.

The thymidylate synthase activity, the karyotype, and the sensitivity to two avian viruses of QT-TK⁻ cells were similar to those of QT35 cells, indicating that the biological characteristics of these two cell lines were similar, except for the TK activity.

Fowlpoxvirus (FPV) strain H and strain FC126 of herpesvirus of turkeys (HVT) did not grow in QT-TK⁻ cells in the presence of 25~50 μ g/ml of BUdR, indicating that QT-TK⁻ cells were useful for the selection of TK⁻ viruses. Therefore, QT-TK⁻ cells were thought to be useful for the construction of recombinant avian viruses that have a foreign gene within the TK gene of the viral DNA.

Furthermore, the TK activity induced by FPV-H and HVT FC126 could be measured specifically in QT-TK⁻ cells. Thus, QT-TK⁻ cells may also be useful in analyzing the viral TKs of avian viruses.