



Title	Enantioselective amination of silylketene acetals with (N-arylsulfonylimino)phenyliodinanes catalyzed by chiral dirhodium(II) carboxylates: asymmetric synthesis of phenylglycine derivatives
Author(s)	Tanaka, Masahiko; Kurosaki, Yasunobu; Washio, Takuya; Anada, Masahiro; Hashimoto, Shunichi
Citation	Tetrahedron Letters, 48(50), 8799-8802 https://doi.org/10.1016/j.tetlet.2007.10.087
Issue Date	2007-12-10
Doc URL	http://hdl.handle.net/2115/32291
Type	article (author version)
File Information	TL0709-509-1.pdf



[Instructions for use](#)

Enantioselective amination of silylketene acetals with (*N*-arylsulfonylimino)phenyliodinanes catalyzed by chiral dirhodium(II) carboxylates: asymmetric synthesis of phenylglycine derivatives

Masahiko Tanaka, Yasunobu Kurosaki, Takuya Washio, Masahiro Anada and Shunichi Hashimoto*


Faculty of Pharmaceutical Sciences, Hokkaido University, Sapporo 060-0812, Japan

Abstract—The first catalytic enantioselective amination of silylketene acetals with (*N*-arylsulfonylimino)phenyliodinanes is described. The reaction of silylketene acetals derived from methyl phenylacetates with [*N*-(2-nitrophenylsulfonyl)imino]phenyliodinane (NsN=IPh) under the catalysis of dirhodium(II) tetrakis[*N*-tetrachlorophthaloyl-(*S*)-*tert*-leucinate], Rh₂(*S*-TCPTTL)₄, proceeds in benzene at room temperature to give *N*-(2-nitrophenylsulfonyl)phenylglycine derivatives in high yields and with enantioselectivities of up to 99% ee.


The catalytic asymmetric amination of silylketene acetals is one of the most straightforward methods for the preparation of enantioenriched α -amino esters.^{1,2} In 1999, Evans and Johnson described the first catalytic enantioselective amination of thioester silylketene acetals or silylketene aminals of acylpyrroles with azodicarboxylate derivatives, in which enantioselectivities up to 99% ee were achieved with the use of a Cu(OTf)₂-bis(oxazoline) catalyst.^{3,4} Kobayashi and co-workers later reported that a AgClO₄-BINAP system catalyzed the amination of the silylketene acetal of phenyl propionate with dibenzyl azodicarboxylate with good enantioselectivity (51% ee).⁵ In this context, the aziridination of silylketene acetals followed by ring opening of the aziridine intermediate provides a prototypical approach to catalytic asymmetric synthesis of α -amino esters.⁶ In 1994, Evans and co-workers reported the first copper(I)-catalyzed amination of silylketene acetals using [(*p*-tolylsulfonyl)imino]phenyliodinane (TsN=IPh, **2a**) as a nitrene precursor,⁷ and they concluded that this protocol does not represent a practical approach to the synthesis of α -amino esters. While high levels of enantiocontrol in aziridinations of alkenes have already been achieved using a variety of different chiral transition metal-catalysts,⁸ the catalytic enantioselective amination reaction of silylketene acetals with (*N*-

arylsulfonylimino)phenyliodinanes has, to the best of our knowledge, not been reported.⁹

Very recently, we demonstrated that the enantioselective amination of silyl enol ethers derived from acyclic ketones or enones with [(2-nitrophenylsulfonyl)imino]phenyliodinane (NsN=IPh, **2b**) catalyzed by chiral dirhodium(II) carboxylates provides *N*-(2-nitrophenylsulfonyl)- α -amino ketones with enantioselectivities of up to 95% ee.¹⁰ In this process, Rh₂(*S*-TFPTTL)₄ (**1a**),¹¹ characterized by the substitution of fluorine atoms for four hydrogen atoms on the phthalimido group in the parent dirhodium(II) complex, Rh₂(*S*-PTTL)₄ (**1c**),¹² proved to be the catalyst of choice in terms of product yield and enantioselectivity as well as catalytic activity. As a logical extension of our studies, we herein



* Corresponding author. Tel.: +81-117-063-236; fax: +81-117-064-981; e-mail: hsmt@pharm.hokudai.ac.jp.

Table 1. Enantioselective amination of silylketene acetals with *N*-(arylsulfonyl)imino]phenyliodinanes catalyzed by chiral dirhodium(II) carboxylates^a

Entry	Silylketene acetal		Iminoiodinane		Rh(II) catalyst	Solvent	Temp. (°C)	Time (h)	α-Amino ester			
	R ₃ Si	Z:E	Ar	Yield (%) ^b					Ee (%) ^c			
1	3a	Et ₃ Si	97:3	2b	2-NO ₂ C ₆ H ₄	Rh ₂ (<i>S</i> -TFPTTL) ₄ (1a)	CH ₂ Cl ₂	0	0.2	4b	95	88
2	3a	Et ₃ Si	97:3	2b	2-NO ₂ C ₆ H ₄	Rh ₂ (<i>S</i> -TCPTTL) ₄ (1b)	CH ₂ Cl ₂	0	0.2	4b	98	97
3	3a	Et ₃ Si	97:3	2b	2-NO ₂ C ₆ H ₄	Rh ₂ (<i>S</i> -TCPTTL) ₄ (1b)	CH ₂ Cl ₂	-20	2.5	4b	91	98
4	3a	Et ₃ Si	97:3	2b	2-NO ₂ C ₆ H ₄	Rh ₂ (<i>S</i> -PTTL) ₄ (1c)	CH ₂ Cl ₂	0	1	4b	80	64
5	3a	Et ₃ Si	97:3	2c	4-NO ₂ C ₆ H ₄	Rh ₂ (<i>S</i> -TCPTTL) ₄ (1b)	CH ₂ Cl ₂	0	0.2	4c	92	59 ^d
6	3a	Et ₃ Si	97:3	2d	2,4-(NO ₂) ₂ C ₆ H ₃	Rh ₂ (<i>S</i> -TCPTTL) ₄ (1b)	CH ₂ Cl ₂	0	2	4d	70	92 ^d
7	3a	Et ₃ Si	97:3	2a	4-MeC ₆ H ₄	Rh ₂ (<i>S</i> -TCPTTL) ₄ (1b)	CH ₂ Cl ₂	0	16	4a	40	88 ^e
8	3a	Et ₃ Si	97:3	2b	2-NO ₂ C ₆ H ₄	Rh ₂ (<i>S</i> -TCPTTL) ₄ (1b)	benzene	23	3	4b	98	99
9	3a	Et ₃ Si	97:3	2b	2-NO ₂ C ₆ H ₄	Rh ₂ (<i>S</i> -TCPTTL) ₄ (1b)	toluene	23	6	4b	91	99
10	3a	Et ₃ Si	23:77	2b	2-NO ₂ C ₆ H ₄	Rh ₂ (<i>S</i> -TCPTTL) ₄ (1b)	benzene	23	3	4b	18 ^e	99
11	3b	Me ₃ Si	97:3	2b	2-NO ₂ C ₆ H ₄	Rh ₂ (<i>S</i> -TCPTTL) ₄ (1b)	benzene	23	3	4b	88	95
12	3c	<i>t</i> -BuMe ₂ Si	96:4	2b	2-NO ₂ C ₆ H ₄	Rh ₂ (<i>S</i> -TCPTTL) ₄ (1b)	benzene	23	3	4b	63	97

^aThe following is a representative procedure (entry 8): **2b** (80.8 mg, 0.2 mmol) was added in one portion to a solution of **3a** (63.4 mg, 0.24 mmol) and Rh₂(*S*-TCPTTL)₄·2EtOAc (**1b**) (3.8 mg, 0.002 mmol, 1 mol %) in benzene (0.5 mL) at 23 °C. The mixture was concentrated in vacuo and chromatographed on silica gel to afford **4b** (68.7 mg, 98%) as a white solid.

^b Isolated yield.

^c Determined by HPLC [column: Daicel Chiralcel OD-H followed by Daicel Chiralpak IB, eluent: 5:1 hexane/*i*-PrOH, flow rate: 1.0 mL/min] unless otherwise stated.

^d Determined by HPLC [column: Daicel Chiralpak AD-H, eluent: 1:1 hexane/*i*-PrOH, flow rate: 1.0 mL/min].

^e Determined by HPLC [column: Daicel Chiralpak AD-H (2×250 mm), eluent: 9:1 hexane/*i*-PrOH, flow rate: 1.0 mL/min].

^f 60% of methyl phenylacetate was recovered.

report that dirhodium(II) tetrakis[*N*-tetrachlorophthaloyl-*(S)*-*tert*-leucinate], Rh₂(*S*-TCPTTL)₄ (**1b**),¹³ catalyzes enantioselective aminations of silylketene acetals derived from methyl phenylacetates with NsN=IPh (**2b**), to provide *N*-(2-nitrophenylsulfonyl)phenylglycine derivatives in high yields and with enantioselectivities of up to 99% ee.¹⁴ While a number of methods,¹⁵ such as the asymmetric Strecker reaction,^{16,17} the Sharpless asymmetric aminohydroxylation,¹⁸ and the catalytic enantioselective hydrogenation of α-aryl imino esters,¹⁹ have been used to synthesize optically active arylglycine derivatives,²⁰ this protocol provides a new, catalytic asymmetric approach to phenylglycine derivatives.

On the basis of our previous work,¹⁰ we initially explored the amination of triethylsilylketene acetal **3a** (*Z*:*E* = 97:3, 1.2 equiv) derived from methyl phenylacetate with NsN=IPh (**2b**) using 1 mol % of Rh₂(*S*-TFPTTL)₄. The reaction in CH₂Cl₂ proceeded smoothly at 0 °C to completion in less than 15 min, giving *N*-(2-nitrophenylsulfonyl)phenylglycine methyl ester (**4b**)²¹ in 95% yield after column chromatography on silica gel (Table 1, entry 1). The enantioselectivity of this reaction was determined to be 88% ee by HPLC analysis with a connected series of Daicel Chiralcel OD-H and Chiralpak IB columns. The preferred absolute stereochemistry of **4b** [[α]_D²⁴ -156.4 (*c* 1.36, CHCl₃)] was assigned as *R* by comparing the sign of optical rotation with (*R*)-**4b** [[α]_D²⁴ -172.3 (*c* 1.01, CHCl₃)], which was prepared from (*R*)-

phenylglycine methyl ester hydrochloride (NsCl, pyridine, CH₂Cl₂, 0 °C, 2 h, 87%).^{21,22} Gratifyingly, switching the catalyst from Rh₂(*S*-TFPTTL)₄ to Rh₂(*S*-TCPTTL)₄ greatly improved the enantioselectivity, providing **4b** in 98% yield with 97% ee (entry 2), although lowering the reaction temperature to -20 °C had little impact on the enantioselectivity (98% ee, entry 3). In this system, Rh₂(*S*-PTTL)₄ was also less effective in terms of both reactivity and enantioselectivity (entry 4). Using Rh₂(*S*-TCPTTL)₄ as the catalyst, we next evaluated the performance of other nitrene precursors, [(4-nitrophenylsulfonyl)imino]phenyliodinane (*p*NsN=IPh, **2c**), [(2,4-dinitrophenylsulfonyl)imino]phenyliodinane (DNsN=IPh, **2d**), and TsN=IPh (**2a**) (entries 5–7). This screening revealed that NsN=IPh (**2b**) was the optimal nitrene precursor for this transformation (entries 2 versus 5–7), although the reason for the advantage of **2b** is not clear at this time. A survey of solvents revealed that the use of benzene was the superior choice as nearly perfect enantiocontrol (99% ee) was achieved at room temperature (entry 8). The product **4b** was also formed with a similar high enantioselectivity in toluene (99% ee, entry 9), but the reaction time (6 h) was longer than that in benzene (3 h). It is noteworthy that the amination of silylketene acetal **3a** (*Z*:*E* = 23:77) with **2b** using Rh₂(*S*-TCPTTL)₄ produced **4b** with the same sense of asymmetric induction as above in only 18% yield with 99% ee, along with 60% recovery of methyl phenylacetate (entry 10). This observation indicates that only the *Z*-isomer of **3a** is responsible for the formation

Table 2. Enantioselective amination of silylketene acetals with **2b** catalyzed by $\text{Rh}_2(\text{S-TCPTTL})_4$ (**1b**)

3 (*Z:E* >95:5) + **2b** $\xrightarrow[\text{benzene, 23 } ^\circ\text{C}]{\text{Rh}_2(\text{S-TCPTTL})_4 \text{ (1b) (1 mol \%)}}$ **4**

Ns = 2-NO₂C₆H₄SO₂


Entry	Silylketene acetal		Time (h)	Phenylglycine derivatives		
	R			Yield (%) ^a	Ee (%) ^b	
1	3a	C ₆ H ₅	3	4b	98	99
2	3d	4-CF ₃ C ₆ H ₄	3	4e	95	97
3	3e	4-FC ₆ H ₄	3	4f	95	97
4	3f	4-ClC ₆ H ₄	2	4g	98	96
5	3g	4-BrC ₆ H ₄	3	4h	93	96
6	3h	4-MeC ₆ H ₄	6	4i	95	98
7	3i	4-MeOC ₆ H ₄	4	4j	98	80
8	3j	3-CF ₃ C ₆ H ₄	2	4k	92	97
9	3k	3-ClC ₆ H ₄	3	4l	94	98
10	3l	3-MeC ₆ H ₄	4	4m	93	98
11	3m	3-MeOC ₆ H ₄	3	4n	48	98

^a Isolated yield.


^b Determined by HPLC. See the supplementary data for details.

of **4b** in the present catalytic process. Therefore, it is not necessary to use geometrically pure silylketene acetals to achieve high enantioselectivity. We also examined the effect of the silicon substituents of the silylketene acetals **3a-c** on enantioselectivity. Variation in the silyl group revealed that the triethylsilyl functionality was optimal for this process (entries 8 versus 11 and 12).

Having demonstrated the effectiveness of the combination of $\text{Rh}_2(\text{S-TCPTTL})_4$ as the catalyst, NsN=IPh as the nitrene precursor and benzene as the solvent, the applicability of this catalytic system to a range of silylketene acetals was then investigated (Table 2). High yields and excellent enantioselectivities were consistently observed at room temperature with electron-withdrawing substituents such as trifluoromethyl and chlorine at the *para* or *meta* position on the benzene ring (96–98% ee, entries 2–5, 8 and 9). The reaction with silylketene acetals **3h** and **3l** bearing a methyl group at the *para* or *meta* position also afforded the corresponding phenylglycine derivatives **4i** and **4m** in high yields and very high enantioselectivities (98% ee, entries 6 and 10). However, the introduction of a methoxy group at the *para* or *meta* position had a detrimental effect on product yield or enantioselectivity. The use of *para*-methoxy-substituted silylketene acetal **3i** resulted in only modest enantioselection (80% ee, entry 7). Although a high enantioselectivity (98% ee) was achieved with *meta*-methoxy-substituted silylketene acetal **3m**, a marked decrease in product yield was observed (entry 11). Surprisingly, the introduction of methyl or chlorine substituents at the *ortho* position on the benzene ring gave aromatic C–H amination products **5a** and **5b** as the sole products instead of the expected phenylglycine derivatives (eq. (1)).²³



The *N*-(2-nitrophenylsulfonyl)phenylglycine derivatives are potentially useful for the synthesis of chiral ligands²⁴ and novel oligopeptide structures^{21,25} (Scheme 1). A single recrystallization of **4b** (99% ee) from EtOH produced an optically pure sample [*mp* = 103.0–104.0 °C, [α]_D²³ –172.0 (*c* 1.00, CHCl₃)] in 91% yield. Reduction of **4b** with DIBAL-H followed by treatment with triphosgene produced the *N*-Ns-protected oxazolidinone **7** [[α]_D²³ –629.7 (*c* 0.50, CHCl₃)] in 93% yield with no racemization.²² Removal of the Ns-group under standard Fukuyama conditions²⁶ furnished an Evans auxiliary (*R*)-**8**,²⁴ [[α]_D²³ –57.0 (*c* 1.01, CHCl₃), lit.,^{24b} [α]_D²² +58.0 (*c* 1.00, CHCl₃)] for (*S*)-**8** in 81% yield. In addition, methyl ester cleavage of **4b** with LiI in refluxing ethyl acetate²⁵ provided (*R*)-*N*-(2-nitrophenylsulfonyl)phenylglycine (**9**) [[α]_D²⁵ –181.5 (*c* 1.01, CHCl₃)] in 94% yield and in enantiomerically pure form.²²



Scheme 1. Reagents and conditions: (a) recrystallization from EtOH, 91%; (b) DIBAL-H, CH₂Cl₂, –78 °C to 0 °C, 0.5 h; (c) triphosgene, Et₃N, CH₂Cl₂, 0 °C, 1 h, 88% (2 steps); (d) PhSH, Cs₂CO₃, DMF, 23 °C, 1 h, 81%; (e) LiI, EtOAc, reflux, 6 h, 94%

In summary, we have developed the first catalytic enantioselective amination of silylketene acetals with (*N*-arylsulfonylimino)phenyliodanes. The reaction of *Z*-triethylsilylketene acetals derived from methyl phenylacetates with NsN=IPh is catalyzed by $\text{Rh}_2(\text{S-TCPTTL})_4$ at room temperature to give *N*-Ns-protected phenylglycine derivatives in high yields and with excellent enantioselectivities, although the efficiency of the present protocol depends on the substitution pattern on the benzene ring. Further studies to expand the range of substrates as well as mechanistic and stereochemical studies are currently underway.

Acknowledgments

This research was supported in part by a Grant-in-Aid for Scientific Research on Priority Areas "Advanced Molecular Transformations of Carbon Resources" from the Ministry of Education, Culture, Sports, Science and Technology, Japan and by Grant-in-Aid from Innovation Plaza Hokkaido in Japan Science and Technology Agency. We thank S. Oka, M. Kiuchi and T. Hirose of the Center for

Instrumental Analysis at Hokkaido University, for technical assistance in MS and elemental analysis.

Supplementary Material

Supplementary data associated with this article can be found, in the online version, at doi:10.1016/j.tetlet.2007.10.087.

References and notes

- For a book and reviews, see: (a) Genet, J.-P.; Greck, C.; Lavergne, D. In *Modern Amination Methods*; Ricci, A., Ed.; Wiley-VCH: Weinheim, 2000, Chapter 2, pp. 65–102; (b) Greck, C.; Drouillat, B.; Thomassigny, C. *Eur. J. Org. Chem.* **2004**, 1377–1385; (c) Erdik, E. *Tetrahedron* **2004**, *60*, 8747–8782; (d) Janey, J. M. *Angew. Chem. Int. Ed.* **2005**, *44*, 4292–4300.
- For general reviews on asymmetric synthesis of α -amino acid derivatives, see: (a) Duthaler, R. O. *Tetrahedron* **1994**, *50*, 1539–1650; (b) Ma, J.-A. *Angew. Chem. Int. Ed.* **2003**, *42*, 4290–4299.
- For a seminal work on catalytic enantioselective direct amination of *N*-acyloxazolidinones with azodicarboxylates, see: Evans, D. A.; Nelson, S. G. *J. Am. Chem. Soc.* **1997**, *119*, 6452–6453.
- Evans, D. A.; Johnson, D. S. *Org. Lett.* **1999**, *1*, 595–598.
- Yamashita, Y.; Ishitani, H.; Kobayashi, S. *Can. J. Chem.* **2000**, *78*, 666–672.
- (a) Cipollone, A.; Loreto, M. A.; Pellacani, L.; Tardella, P. A. *J. Org. Chem.* **1987**, *52*, 2584–2586; (b) Loreto, M. A.; Pellacani, L.; Tardella, P. A. *Tetrahedron Lett.* **1989**, *30*, 2975–2978.
- Evans, D. A.; Faul, M. M.; Bilodeau, M. T. *J. Am. Chem. Soc.* **1994**, *116*, 2742–2753.
- For reviews on transition metal-catalyzed enantioselective aziridination of alkenes with (arylsulfonylimino)phenyliodines and arylsulfonyl azides, see: (a) Jacobsen E. N. In *Comprehensive Asymmetric Catalysis*; Jacobsen, E. N., Pfaltz, A., Yamamoto, H., Eds.; Springer: Berlin, 1999; Vol. 2, Chapter 17, pp. 607–618; (b) Müller, P.; Fruit, C. *Chem. Rev.* **2003**, *103*, 2905–2919; (c) Katsuki, T. *Chem. Lett.* **2005**, *34*, 1304–1309.
- For catalytic enantioselective amination of silyl enol ethers using TsN=IPh as a nitrene precursor, see: (a) Adam, W.; Roschmann, K. J.; Saha-Möller, C. R. *Eur. J. Org. Chem.* **2000**, 557–561; (b) Liang, J.-L.; Yu, X.-Q.; Che, C.-M. *Chem. Commun.* **2002**, 124–125.
- Anada, M.; Tanaka, M.; Washio, T.; Yamawaki, M.; Abe, T.; Hashimoto, S. *Org. Lett.* in press.
- Tsutsui, H.; Yamaguchi, Y.; Kitagaki, S.; Nakamura, S.; Anada, M.; Hashimoto, S. *Tetrahedron: Asymmetry* **2003**, *14*, 817–821.
- Minami, K.; Saito, H.; Tsutsui, H.; Nambu, H.; Anada, M.; Hashimoto, S. *Adv. Synth. Catal.* **2005**, *347*, 1483–1487 and references cited therein.
- We previously disclosed that Rh₂(S-TCPTTL)₄ is an efficient catalyst for intermolecular benzylic C–H amination reaction, see: Yamawaki, M.; Tsutsui, H.; Kitagaki, S.; Anada, M.; Hashimoto, S. *Tetrahedron Lett.* **2002**, *43*, 9561–9564.
- Recently, Davies and Reddy reported highly enantioselective benzylic C–H aminations using dirhodium(II) tetrakis[*N*-tetrachlorophthaloyl-(*S*)-(1-adamantyl)glycinate], [Rh₂(S-TCPTAD)₄], as a catalyst, see: Reddy, R. P.; Davies, H. M. L. *Org. Lett.* **2006**, *8*, 5013–5016.
- For a review on asymmetric synthesis of arylglycines, see: Williams, R. M.; Hendric, J. A. *Chem. Rev.* **1992**, *92*, 889–917.
- For recent reviews on asymmetric Strecker reactions, see: (a) Yet, L. *Angew. Chem. Int. Ed.* **2001**, *40*, 875–877; (b) Gröger, H. *Chem. Rev.* **2003**, *103*, 2795–2827; (c) Spino, C. *Angew. Chem. Int. Ed.* **2004**, *43*, 1764–1766.
- For selected examples of catalytic enantioselective synthesis of arylglycine derivatives via the Strecker reaction, see: (a) Iyer, M. S.; Gigstad, K. M.; Namdev, N. D.; Lipton, M. *J. Am. Chem. Soc.* **1996**, *118*, 4910–4911; (b) Sigman, M. S.; Jacobsen, E. N. *J. Am. Chem. Soc.* **1998**, *120*, 5315–5316; (c) Ishitani, H.; Komiyama, S.; Kobayashi, S. *Angew. Chem. Int. Ed.* **1998**, *37*, 3186–3188; (d) Krueger, C. A.; Kuntz, K. W.; Dzierba, C. D.; Wirschun, W. G.; Gleason, J. D.; Snapper, M. L.; Hoveyda, A. H. *J. Am. Chem. Soc.* **1999**, *121*, 4284–4285; (e) Corey, E. J.; Grogan, M. J. *Org. Lett.* **1999**, *1*, 157–160; (f) Takamura, M.; Hamashima, Y.; Usuda, H.; Kanai, M.; Shibasaki, M. *Angew. Chem. Int. Ed.* **2000**, *39*, 1650–1652; (g) Funabashi, K.; Ratni, H.; Kanai, M.; Shibasaki, M. *J. Am. Chem. Soc.* **2001**, *123*, 10784–10785; (h) Vachal, P.; Jacobsen, E. N. *J. Am. Chem. Soc.* **2002**, *124*, 10012–10014; (i) Masumoto, S.; Usuda, H.; Suzuki, M.; Kanai, M.; Shibasaki, M. *J. Am. Chem. Soc.* **2003**, *125*, 5634–5635; (j) Kato, N.; Suzuki, M.; Kanai, M.; Shibasaki, M. *Tetrahedron Lett.* **2004**, *45*, 3147–3151; (k) Kato, N.; Suzuki, M.; Kanai, M.; Shibasaki, M. *Tetrahedron Lett.* **2004**, *45*, 3153–3155; (l) Huang, J.; Corey, E. J. *Org. Lett.* **2004**, *6*, 5027–5029; (m) Banphavichit, V.; Mansawat, W.; Bhanthumnavin, W.; Vilaivan, T. *Tetrahedron* **2004**, *60*, 10559–10568; (n) Nakamura, S.; Nakashima, H.; Sugimoto, H.; Shibata, N.; Toru, T. *Tetrahedron Lett.* **2006**, *47*, 7599–7602; (o) Pan, S. C.; Zhou, J.; List, B. *Angew. Chem. Int. Ed.* **2007**, *46*, 612–614; (p) Huang, J.; Liu, X.; Wen, Y.; Qin, B.; Feng, X. *J. Org. Chem.* **2007**, *72*, 204–208.
- (a) Li, G.; Angert, H. H.; Sharpless, K. B. *Angew. Chem., Int. Ed. Engl.* **1996**, *35*, 2813–2817; (b) Reddy, K. L.; Sharpless, K. B. *J. Am. Chem. Soc.* **1998**, *120*, 1207–1217.
- (a) Shang, G.; Yang, Q.; Zhang, X. *Angew. Chem. Int. Ed.* **2006**, *45*, 6360–6362; (b) Li, G.; Liang, Y.; Antilla, J. C. *J. Am. Chem. Soc.* **2007**, *129*, 5830–5831.
- Mellin-Morliére, C.; Aitken, D. J.; Bull, S. D.; Davies, S. G.; Husson, H.-P. *Tetrahedron: Asymmetry* **2001**, *12*, 149–155.
- Albanese, D.; Landini, D.; Lupi, V.; Penso, M. *Eur. J. Org. Chem.* **2000**, 1443–1449.
- See the supplementary data for experimental details.
- It was found that the presence of an aryl group in silylketene acetals is responsible for the high enantioselectivity. For example, the amination of silylketene acetal derived from methyl cyclohexylacetate (*Z:E* = 64:36, 2 equiv) with NsN=IPh (**2b**) in the presence of Rh₂(S-TCPTTL)₄ afforded the corresponding α -amino ester **6** in 92% yield with 48% ee.
- (a) Evans, D. A.; Sjogren, E. B. *Tetrahedron Lett.* **1985**, *26*, 3783–3786; (b) Nicolás, E.; Russell, K. C.; Hrubby, V. J. *J. Org. Chem.* **1993**, *58*, 766–770.
- Biron, E.; Kessler, H. *J. Org. Chem.* **2005**, *70*, 5183–5189.
- (a) Fukuyama, T.; Jow, C.-K.; Cheung, M. *Tetrahedron Lett.* **1995**, *36*, 6373–6374; For a review on the nitrophenylsulfonamide chemistry, see: (b) Kan, T.; Fukuyama, T. *Chem. Commun.* **2004**, 353–359.